

Class XI : Maths
Chapter 13 : Statistics

Questions and Solutions | Exercise 13.1 - NCERT Books

Question 1:

Find the mean deviation about the mean for the data

4, 7, 8, 9, 10, 12, 13, 17

Answer

The given data is

4, 7, 8, 9, 10, 12, 13, 17

$$\bar{x} = \frac{4+7+8+9+10+12+13+17}{8} = \frac{80}{8} = 10$$

Mean of the data,

The deviations of the respective observations from the mean \bar{x} , i.e. $x_i - \bar{x}$, are
-6, -3, -2, -1, 0, 2, 3, 7

The absolute values of the deviations, i.e. $|x_i - \bar{x}|$, are
6, 3, 2, 1, 0, 2, 3, 7

The required mean deviation about the mean is

$$\text{M.D.}(\bar{x}) = \frac{\sum_{i=1}^8 |x_i - \bar{x}|}{8} = \frac{6+3+2+1+0+2+3+7}{8} = \frac{24}{8} = 3$$

Question 2:

Find the mean deviation about the mean for the data

38, 70, 48, 40, 42, 55, 63, 46, 54, 44

Answer

The given data is

38, 70, 48, 40, 42, 55, 63, 46, 54, 44

Mean of the given data,

$$\bar{x} = \frac{38+70+48+40+42+55+63+46+54+44}{10} = \frac{500}{10} = 50$$

The deviations of the respective observations from the mean \bar{x} , i.e. $x_i - \bar{x}$, are
-12, 20, -2, -10, -8, 5, 13, -4, 4, -6

The absolute values of the deviations, i.e. $|x_i - \bar{x}|$, are
12, 20, 2, 10, 8, 5, 13, 4, 4, 6

The required mean deviation about the mean is

$$\begin{aligned} \text{M.D.}(\bar{x}) &= \frac{\sum_{i=1}^{10} |x_i - \bar{x}|}{10} \\ &= \frac{12 + 20 + 2 + 10 + 8 + 5 + 13 + 4 + 4 + 6}{10} \\ &= \frac{84}{10} \\ &= 8.4 \end{aligned}$$

Question 3:

Find the mean deviation about the median for the data.

13, 17, 16, 14, 11, 13, 10, 16, 11, 18, 12, 17

Answer

The given data is

13, 17, 16, 14, 11, 13, 10, 16, 11, 18, 12, 17

Here, the numbers of observations are 12, which is even.

Arranging the data in ascending order, we obtain

10, 11, 11, 12, 13, 13, 14, 16, 16, 17, 17, 18

$$\begin{aligned} \text{Median, } M &= \frac{\left(\frac{12}{2}\right)^{\text{th}} \text{ observation} + \left(\frac{12}{2} + 1\right)^{\text{th}} \text{ observation}}{2} \\ &= \frac{6^{\text{th}} \text{ observation} + 7^{\text{th}} \text{ observation}}{2} \\ &= \frac{13 + 14}{2} = \frac{27}{2} = 13.5 \end{aligned}$$

The deviations of the respective observations from the median, i.e. $x_i - M$, are
-3.5, -2.5, -2.5, -1.5, -0.5, -0.5, 0.5, 2.5, 2.5, 3.5, 3.5, 4.5

The absolute values of the deviations, $|x_i - M|$, are
3.5, 2.5, 2.5, 1.5, 0.5, 0.5, 0.5, 2.5, 2.5, 3.5, 3.5, 4.5

The required mean deviation about the median is

$$\begin{aligned} \text{M.D.}(M) &= \frac{\sum_{i=1}^{12} |x_i - M|}{12} \\ &= \frac{3.5 + 2.5 + 2.5 + 1.5 + 0.5 + 0.5 + 0.5 + 2.5 + 2.5 + 3.5 + 3.5 + 4.5}{12} \\ &= \frac{28}{12} = 2.33 \end{aligned}$$

Question 4:

Find the mean deviation about the median for the data

36, 72, 46, 42, 60, 45, 53, 46, 51, 49

Answer

The given data is

36, 72, 46, 42, 60, 45, 53, 46, 51, 49

Here, the number of observations is 10, which is even.

Arranging the data in ascending order, we obtain

36, 42, 45, 46, 46, 49, 51, 53, 60, 72

$$\begin{aligned} \text{Median } M &= \frac{\left(\frac{10}{2}\right)^{\text{th}} \text{ observation} + \left(\frac{10}{2} + 1\right)^{\text{th}} \text{ observation}}{2} \\ &= \frac{5^{\text{th}} \text{ observation} + 6^{\text{th}} \text{ observation}}{2} \\ &= \frac{46 + 49}{2} = \frac{95}{2} = 47.5 \end{aligned}$$

The deviations of the respective observations from the median, i.e. $x_i - M$, are
-11.5, -5.5, -2.5, -1.5, -1.5, 1.5, 3.5, 5.5, 12.5, 24.5

The absolute values of the deviations, $|x_i - M|$, are
11.5, 5.5, 2.5, 1.5, 1.5, 1.5, 3.5, 5.5, 12.5, 24.5

Thus, the required mean deviation about the median is

$$\begin{aligned} \text{M.D.}(M) &= \frac{\sum_{i=1}^{10} |x_i - M|}{10} = \frac{11.5 + 5.5 + 2.5 + 1.5 + 1.5 + 1.5 + 3.5 + 5.5 + 12.5 + 24.5}{10} \\ &= \frac{70}{10} = 7 \end{aligned}$$

Question 5:

Find the mean deviation about the mean for the data.

x_i	5	10	15	20	25
f_i	7	4	6	3	5

Answer

x_i	f_i	$f_i x_i$	$ x_i - \bar{x} $	$f_i x_i - \bar{x} $
5	7	35	9	63
10	4	40	4	16
15	6	90	1	6
20	3	60	6	18
25	5	125	11	55
	25	350		158

$$N = \sum_{i=1}^5 f_i = 25$$

$$\sum_{i=1}^5 f_i x_i = 350$$

$$\therefore \bar{x} = \frac{1}{N} \sum_{i=1}^5 f_i x_i = \frac{1}{25} \times 350 = 14$$

$$\therefore MD(\bar{x}) = \frac{1}{N} \sum_{i=1}^5 f_i |x_i - \bar{x}| = \frac{1}{25} \times 158 = 6.32$$

Question 6:

Find the mean deviation about the mean for the data

x_i	10	30	50	70	90
f_i	4	24	28	16	8

Answer

x_i	f_i	$f_i x_i$	$ x_i - \bar{x} $	$f_i x_i - \bar{x} $
10	4	40	40	160
30	24	720	20	480
50	28	1400	0	0
70	16	1120	20	320
90	8	720	40	320
	80	4000		1280

$$N = \sum_{i=1}^5 f_i = 80, \quad \sum_{i=1}^5 f_i x_i = 4000$$

$$\therefore \bar{x} = \frac{1}{N} \sum_{i=1}^5 f_i x_i = \frac{1}{80} \times 4000 = 50$$

$$MD(\bar{x}) = \frac{1}{N} \sum_{i=1}^5 f_i |x_i - \bar{x}| = \frac{1}{80} \times 1280 = 16$$

Question 7:

Find the mean deviation about the median for the data.

x_i	5	7	9	10	12	15
f_i	8	6	2	2	2	6

Answer

The given observations are already in ascending order.

Adding a column corresponding to cumulative frequencies of the given data, we obtain the following table.

x_i	f_i	c.f.
5	8	8
7	6	14
9	2	16
10	2	18
12	2	20
15	6	26

Here, $N = 26$, which is even.

Median is the mean of 13th and 14th observations. Both of these observations lie in the cumulative frequency 14, for which the corresponding observation is 7.

$$\therefore \text{Median} = \frac{13^{\text{th}} \text{ observation} + 14^{\text{th}} \text{ observation}}{2} = \frac{7+7}{2} = 7$$

The absolute values of the deviations from median, i.e. $|x_i - M|$, are

$ x_i - M $	2	0	2	3	5	8
f_i	8	6	2	2	2	6
$f_i x_i - M $	16	0	4	6	10	48

$$\sum_{i=1}^6 f_i = 26 \quad \text{and} \quad \sum_{i=1}^6 f_i |x_i - M| = 84$$

$$\text{M.D.}(M) = \frac{1}{N} \sum_{i=1}^6 f_i |x_i - M| = \frac{1}{26} \times 84 = 3.23$$

Question 8:

Find the mean deviation about the median for the data

x_i	15	21	27	30	35
f_i	3	5	6	7	8

Answer

The given observations are already in ascending order.

Adding a column corresponding to cumulative frequencies of the given data, we obtain the following table.

x_i	f_i	c.f.
15	3	3
21	5	8
27	6	14
30	7	21
35	8	29

Here, $N = 29$, which is odd.

$$\therefore \text{Median} = \left(\frac{29+1}{2} \right)^{\text{th}} \text{ observation} = 15^{\text{th}} \text{ observation}$$

This observation lies in the cumulative frequency 21, for which the corresponding observation is 30.

$$\therefore \text{Median} = 30$$

The absolute values of the deviations from median, i.e. $|x_i - M|$, are

$ x_i - M $	15	9	3	0	5
f_i	3	5	6	7	8
$f_i x_i - M $	45	45	18	0	40

$$\sum_{i=1}^5 f_i = 29, \quad \sum_{i=1}^5 f_i |x_i - M| = 148$$

$$\therefore \text{M.D.}(M) = \frac{1}{N} \sum_{i=1}^5 f_i |x_i - M| = \frac{1}{29} \times 148 = 5.1$$

Question 9:

Find the mean deviation about the mean for the data.

Income per day	Number of persons
0-100	4
100-200	8
200-300	9
300-400	10
400-500	7
500-600	5
600-700	4

700-800	3
---------	---

Answer

The following table is formed.

Income per day	Number of persons f_i	Mid-point x_i	$f_i x_i$	$ x_i - \bar{x} $	$f_i x_i - \bar{x} $
0 - 100	4	50	200	308	1232
100 - 200	8	150	1200	208	1664
200 - 300	9	250	2250	108	972
300 - 400	10	350	3500	8	80
400 - 500	7	450	3150	92	644
500 - 600	5	550	2750	192	960
600 - 700	4	650	2600	292	1168
700 - 800	3	750	2250	392	1176
	50		17900		7896

$$N = \sum_{i=1}^8 f_i = 50, \quad \sum_{i=1}^8 f_i x_i = 17900$$

Here,

$$\therefore \bar{x} = \frac{1}{N} \sum_{i=1}^8 f_i x_i = \frac{1}{50} \times 17900 = 358$$

$$\text{M.D.}(\bar{x}) = \frac{1}{N} \sum_{i=1}^8 f_i |x_i - \bar{x}| = \frac{1}{50} \times 7896 = 157.92$$

Question 10:

Find the mean deviation about the mean for the data

Height in cms	Number of boys
---------------	----------------

95-105	9
105-115	13
115-125	26
125-135	30
135-145	12
145-155	10

Answer

The following table is formed.

Height in cms	Number of boys f_i	Mid-point x_i	$f_i x_i$	$ x_i - \bar{x} $	$f_i x_i - \bar{x} $
95-105	9	100	900	25.3	227.7
105-115	13	110	1430	15.3	198.9
115-125	26	120	3120	5.3	137.8
125-135	30	130	3900	4.7	141
135-145	12	140	1680	14.7	176.4
145-155	10	150	1500	24.7	247

Here, $N = \sum_{i=1}^6 f_i = 100, \sum_{i=1}^6 f_i x_i = 12530$

$$\therefore \bar{x} = \frac{1}{N} \sum_{i=1}^6 f_i x_i = \frac{1}{100} \times 12530 = 125.3$$

$$M.D.(\bar{x}) = \frac{1}{N} \sum_{i=1}^6 f_i |x_i - \bar{x}| = \frac{1}{100} \times 1128.8 = 11.28$$

Question 11:

Find the mean deviation about median for the following data:

Marks	Number of girls
0-10	6
10-20	8
20-30	14
30-40	16
40-50	4
50-60	2

Answer

The following table is formed.

Marks	Number of boys f_i	Cumulative frequency (c.f.)	Mid-point x_i	$ x_i - \text{Med.} $	$f_i x_i - \text{Med.} $
0-10	6	6	5	22.85	137.1
10-20	8	14	15	12.85	102.8
20-30	14	28	25	2.85	39.9
30-40	16	44	35	7.15	114.4
40-50	4	48	45	17.15	68.6
50-60	2	50	55	27.15	54.3
	50				517.1

The class interval containing the $\left(\frac{N}{2}\right)^{\text{th}}$ or 25th item is 20 – 30.

Therefore, 20 – 30 is the median class.

It is known that,

$$\text{Median} = l + \frac{\frac{N}{2} - C}{f} \times h$$

Here, $l = 20$, $C = 14$, $f = 14$, $h = 10$, and $N = 50$

$$\therefore \text{Median} = 20 + \frac{25 - 14}{14} \times 10 = 20 + \frac{110}{14} = 20 + 7.85 = 27.85$$

Thus, mean deviation about the median is given by,

$$\text{M.D.}(M) = \frac{1}{N} \sum_{i=1}^6 f_i |x_i - M| = \frac{1}{50} \times 517.1 = 10.34$$

Question 12:

Calculate the mean deviation about median age for the age distribution of 100 persons given below:

Age	Number
16-20	5
21-25	6
26-30	12
31-35	14
36-40	26
41-45	12
46-50	16
51-55	9

Answer

The given data is not continuous. Therefore, it has to be converted into continuous frequency distribution by subtracting 0.5 from the lower limit and adding 0.5 to the upper limit of each class interval.

The table is formed as follows.

Age	Number f_i	Cumulative frequency (c.f.)	Mid- point x_i	$ x_i -$ Med.	$f_i x_i -$ Med.
15.5- 20.5	5	5	18	20	100
20.5- 25.5	6	11	23	15	90
25.5- 30.5	12	23	28	10	120
30.5- 35.5	14	37	33	5	70
35.5- 40.5	26	63	38	0	0
40.5- 45.5	12	75	43	5	60
45.5- 50.5	16	91	48	10	160
50.5- 55.5	9	100	53	15	135
	100				735

The class interval containing the $\frac{N}{2}$ or 50th item is 35.5 – 40.5.

Therefore, 35.5 – 40.5 is the median class.

It is known that,

$$\text{Median} = l + \frac{\frac{N}{2} - C}{f} \times h$$

Here, $l = 35.5$, $C = 37$, $f = 26$, $h = 5$, and $N = 100$

$$\therefore \text{Median} = 35.5 + \frac{50 - 37}{26} \times 5 = 35.5 + \frac{13 \times 5}{26} = 35.5 + 2.5 = 38$$

Thus, mean deviation about the median is given by,

$$\text{M.D.}(M) = \frac{1}{N} \sum_{i=1}^8 f_i |x_i - M| = \frac{1}{100} \times 735 = 7.35$$



Class XI : Maths
Chapter 13 : Statistics

Questions and Solutions | Exercise 13.2 - NCERT Books

Question 1:

Find the mean and variance for the data 6, 7, 10, 12, 13, 4, 8, 12

Answer

6, 7, 10, 12, 13, 4, 8, 12

$$\bar{x} = \frac{\sum_{i=1}^8 x_i}{n} = \frac{6+7+10+12+13+4+8+12}{8} = \frac{72}{8} = 9$$

Mean,

The following table is obtained.

x_i	$(x_i - \bar{x})$	$(x_i - \bar{x})^2$
6	-3	9
7	-2	4
10	-1	1
12	3	9
13	4	16
4	-5	25
8	-1	1
12	3	9
		74

$$\text{Variance}(\sigma^2) = \frac{1}{n} \sum_{i=1}^8 (x_i - \bar{x})^2 = \frac{1}{8} \times 74 = 9.25$$

Question 2:

Find the mean and variance for the first n natural numbers

Answer

The mean of first n natural numbers is calculated as follows.

$$\text{Mean} = \frac{\text{Sum of all observations}}{\text{Number of observations}}$$

$$\therefore \text{Mean} = \frac{\frac{n(n+1)}{2}}{n} = \frac{n+1}{2}$$

$$\begin{aligned} \text{Variance}(\sigma^2) &= \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 \\ &= \frac{1}{n} \sum_{i=1}^n \left[x_i - \left(\frac{n+1}{2} \right) \right]^2 \\ &= \frac{1}{n} \sum_{i=1}^n x_i^2 - \frac{1}{n} \sum_{i=1}^n 2 \left(\frac{n+1}{2} \right) x_i + \frac{1}{n} \sum_{i=1}^n \left(\frac{n+1}{2} \right)^2 \\ &= \frac{1}{n} \frac{n(n+1)(2n+1)}{6} - \left(\frac{n+1}{n} \right) \left[\frac{n(n+1)}{2} \right] + \frac{(n+1)^2}{4n} \times n \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{2} + \frac{(n+1)^2}{4} \\ &= \frac{(n+1)(2n+1)}{6} - \frac{(n+1)^2}{4} \\ &= (n+1) \left[\frac{4n+2-3n-3}{12} \right] \\ &= \frac{(n+1)(n-1)}{12} \\ &= \frac{n^2-1}{12} \end{aligned}$$

Question 3:

Find the mean and variance for the first 10 multiples of 3

Answer

The first 10 multiples of 3 are

3, 6, 9, 12, 15, 18, 21, 24, 27, 30

Here, number of observations, $n = 10$

$$\text{Mean, } \bar{x} = \frac{\sum_{i=1}^{10} x_i}{10} = \frac{165}{10} = 16.5$$

The following table is obtained.

x_i	$(x_i - \bar{x})$	$(x_i - \bar{x})^2$
3	-13.5	182.25
6	-10.5	110.25
9	-7.5	56.25
12	-4.5	20.25
15	-1.5	2.25
18	1.5	2.25
21	4.5	20.25
24	7.5	56.25
27	10.5	110.25
30	13.5	182.25
		742.5

$$\text{Variance}(\sigma^2) = \frac{1}{n} \sum_{i=1}^{10} (x_i - \bar{x})^2 = \frac{1}{10} \times 742.5 = 74.25$$

Question 4:

Find the mean and variance for the data

x_i	6	10	14	18	24	28	30
f_i	2	4	7	12	8	4	3

Answer

The data is obtained in tabular form as follows.

x_i	f_i	$f_i x_i$	$x_i - \bar{x}$	$(x_i - \bar{x})^2$	$f_i (x_i - \bar{x})^2$
6	2	12	-13	169	338
10	4	40	-9	81	324
14	7	98	-5	25	175
18	12	216	-1	1	12
24	8	192	5	25	200
28	4	112	9	81	324
30	3	90	11	121	363
	40	760			1736

Here, $N = 40$, $\sum_{i=1}^7 f_i x_i = 760$

$$\therefore \bar{x} = \frac{\sum_{i=1}^7 f_i x_i}{N} = \frac{760}{40} = 19$$

$$\text{Variance} = (\sigma^2) = \frac{1}{N} \sum_{i=1}^7 f_i (x_i - \bar{x})^2 = \frac{1}{40} \times 1736 = 43.4$$

Question 5:

Find the mean and variance for the data

x_i	92	93	97	98	102	104	109
f_i	3	2	3	2	6	3	3

Answer

The data is obtained in tabular form as follows.

x_i	f_i	$f_i x_i$	$x_i - \bar{x}$	$(x_i - \bar{x})^2$	$f_i (x_i - \bar{x})^2$
92	3	276	-8	64	192
93	2	186	-7	49	98
97	3	291	-3	9	27
98	2	196	-2	4	8
102	6	612	2	4	24
104	3	312	4	16	48
109	3	327	9	81	243
	22	2200			640

Here, $N = 22$, $\sum_{i=1}^7 f_i x_i = 2200$

$$\therefore \bar{x} = \frac{1}{N} \sum_{i=1}^7 f_i x_i = \frac{1}{22} \times 2200 = 100$$

$$\text{Variance}(\sigma^2) = \frac{1}{N} \sum_{i=1}^7 f_i (x_i - \bar{x})^2 = \frac{1}{22} \times 640 = 29.09$$

Question 6:

Find the mean and standard deviation using short-cut method.

x_i	60	61	62	63	64	65	66	67	68
f_i	2	1	12	29	25	12	10	4	5

Answer

The data is obtained in tabular form as follows.

x_i	f_i	$f_i = \frac{x_i - 64}{1}$	y_i^2	$f_i y_i$	$f_i y_i^2$
60	2	-4	16	-8	32
61	1	-3	9	-3	9
62	12	-2	4	-24	48
63	29	-1	1	-29	29
64	25	0	0	0	0
65	12	1	1	12	12
66	10	2	4	20	40
67	4	3	9	12	36
68	5	4	16	20	80
	100	220		0	286

$$\bar{x} = A + \frac{\sum_{i=1}^9 f_i y_i}{N} \times h = 64 + \frac{0}{100} \times 1 = 64 + 0 = 64$$

Mean,

$$\begin{aligned} \text{Variance, } \sigma^2 &= \frac{h^2}{N^2} \left[N \sum_{i=1}^9 f_i y_i^2 - \left(\sum_{i=1}^9 f_i y_i \right)^2 \right] \\ &= \frac{1}{100^2} [100 \times 286 - 0] \\ &= 2.86 \end{aligned}$$

$$\therefore \text{Standard deviation } (\sigma) = \sqrt{2.86} = 1.69$$

Question 7:

Find the mean and variance for the following frequency distribution.

Classes	0-30	30-60	60-90	90-120	120-150	150-180	180-210
Frequencies	2	3	5	10	3	5	2

Answer

Class	Frequency f_i	Mid-point x_i	$y_i = \frac{x_i - 105}{30}$	y_i^2	$f_i y_i$	$f_i y_i^2$
0-30	2	15	-3	9	-6	18
30-60	3	45	-2	4	-6	12
60-90	5	75	-1	1	-5	5
90-120	10	105	0	0	0	0
120-150	3	135	1	1	3	3
150-180	5	165	2	4	10	20
180-210	2	195	3	9	6	18
	30				2	76

$$\text{Mean, } \bar{x} = A + \frac{\sum_{i=1}^7 f_i y_i}{N} \times h = 105 + \frac{2}{30} \times 30 = 105 + 2 = 107$$

$$\begin{aligned} \text{Variance } (\sigma^2) &= \frac{h^2}{N^2} \left[N \sum_{i=1}^7 f_i y_i^2 - \left(\sum_{i=1}^7 f_i y_i \right)^2 \right] \\ &= \frac{(30)^2}{(30)^2} [30 \times 76 - (2)^2] \\ &= 2280 - 4 \\ &= 2276 \end{aligned}$$

Question 8:

Find the mean and variance for the following frequency distribution.

Classes	0-10	10-20	20-30	30-40	40-50
Frequencies	5	8	15	16	6

Answer

Class	Frequency f_i	Mid-point x_i	$y_i = \frac{x_i - 25}{10}$	y_i^2	$f_i y_i$	$f_i y_i^2$
0-10	5	5	-2	4	-10	20
10-20	8	15	-1	1	-8	8
20-30	15	25	0	0	0	0
30-40	16	35	1	1	16	16
40-50	6	45	2	4	12	24
	50				10	68

$$\text{Mean, } \bar{x} = A + \frac{\sum_{i=1}^5 f_i y_i}{N} \times h = 25 + \frac{10}{50} \times 10 = 25 + 2 = 27$$

$$\begin{aligned} \text{Variance } (\sigma^2) &= \frac{h^2}{N^2} \left[N \sum_{i=1}^5 f_i y_i^2 - \left(\sum_{i=1}^5 f_i y_i \right)^2 \right] \\ &= \frac{(10)^2}{(50)^2} [50 \times 68 - (10)^2] \\ &= \frac{1}{25} [3400 - 100] = \frac{3300}{25} \\ &= 132 \end{aligned}$$

Question 9:

Find the mean, variance and standard deviation using short-cut method

Height in cms	No. of children
70-75	3
75-80	4
80-85	7
85-90	7
90-95	15
95-100	9
100-105	6
105-110	6
110-115	3

Answer

Class Interval	Frequency f_i	Mid-point x_i	$y_i = \frac{x_i - 92.5}{5}$	y_i^2	$f_i y_i$	$f_i y_i^2$
70-75	3	72.5	-4	16	-12	48
75-80	4	77.5	-3	9	-12	36
80-85	7	82.5	-2	4	-14	28
85-90	7	87.5	-1	1	-7	7
90-95	15	92.5	0	0	0	0

95-100	9	97.5	1	1	9	9
100-105	6	102.5	2	4	12	24
105-110	6	107.5	3	9	18	54
110-115	3	112.5	4	16	12	48
	60				6	254

$$\text{Mean, } \bar{x} = A + \frac{\sum_{i=1}^9 f_i y_i}{N} \times h = 92.5 + \frac{6}{60} \times 5 = 92.5 + 0.5 = 93$$

$$\begin{aligned} \text{Variance } (\sigma^2) &= \frac{h^2}{N^2} \left[N \sum_{i=1}^9 f_i y_i^2 - \left(\sum_{i=1}^9 f_i y_i \right)^2 \right] \\ &= \frac{(5)^2}{(60)^2} [60 \times 254 - (6)^2] \\ &= \frac{25}{3600} (15204) = 105.58 \end{aligned}$$

$$\therefore \text{Standard deviation } (\sigma) = \sqrt{105.58} = 10.27$$

Question 10:

The diameters of circles (in mm) drawn in a design are given below:

Diameters	No. of children
33-36	15
37-40	17
41-44	21

45-48	22
49-52	25

Answer

Class Interval	Frequency f_i	Mid-point x_i	$y_i = \frac{x_i - 42.5}{4}$	f_i^2	$f_i y_i$	$f_i y_i^2$
32.5-36.5	15	34.5	-2	4	-30	60
36.5-40.5	17	38.5	-1	1	-17	17
40.5-44.5	21	42.5	0	0	0	0
44.5-48.5	22	46.5	1	1	22	22
48.5-52.5	25	50.5	2	4	50	100
	100				25	199

Here, $N = 100$, $h = 4$

Let the assumed mean, A , be 42.5.

$$\text{Mean, } \bar{x} = A + \frac{\sum_{i=1}^5 f_i y_i}{N} \times h = 42.5 + \frac{25}{100} \times 4 = 43.5$$

$$\begin{aligned} \text{Variance } (\sigma^2) &= \frac{h^2}{N^2} \left[N \sum_{i=1}^5 f_i y_i^2 - \left(\sum_{i=1}^5 f_i y_i \right)^2 \right] \\ &= \frac{16}{10000} [100 \times 199 - (25)^2] \\ &= \frac{16}{10000} [19900 - 625] \\ &= \frac{16}{10000} \times 19275 \\ &= 30.84 \end{aligned}$$

\therefore Standard deviation (σ) = 5.55

Class XI : Maths
Chapter 13 : Statistics

Questions and Solutions | Miscellaneous Exercise 13 - NCERT Books

Question 1:

The mean and variance of eight observations are 9 and 9.25, respectively. If six of the observations are 6, 7, 10, 12, 12 and 13, find the remaining two observations.

Answer

Let the remaining two observations be x and y .

Therefore, the observations are 6, 7, 10, 12, 12, 13, x , y .

$$\text{Mean, } \bar{x} = \frac{6+7+10+12+12+13+x+y}{8} = 9$$

$$\Rightarrow 60 + x + y = 72$$

$$\Rightarrow x + y = 12 \quad \dots(1)$$

$$\text{Variance} = 9.25 = \frac{1}{n} \sum_{i=1}^8 (x_i - \bar{x})^2$$

$$9.25 = \frac{1}{8} [(-3)^2 + (-2)^2 + (1)^2 + (3)^2 + (3)^2 + (4)^2 + x^2 + y^2 - 2 \times 9(x+y) + 2 \times (9)^2]$$

$$9.25 = \frac{1}{8} [9 + 4 + 1 + 9 + 9 + 16 + x^2 + y^2 - 18(12) + 162]$$

...[Using (1)]

$$9.25 = \frac{1}{8} [48 + x^2 + y^2 - 216 + 162]$$

$$9.25 = \frac{1}{8} [x^2 + y^2 - 6]$$

$$\Rightarrow x^2 + y^2 = 80 \quad \dots(2)$$

From (1), we obtain

$$x^2 + y^2 + 2xy = 144 \quad \dots(3)$$

From (2) and (3), we obtain

$$2xy = 64 \quad \dots(4)$$

Subtracting (4) from (2), we obtain

$$x^2 + y^2 - 2xy = 80 - 64 = 16$$

$$\Rightarrow x - y = \pm 4 \quad \dots(5)$$

Therefore, from (1) and (5), we obtain

$$x = 8 \text{ and } y = 4, \text{ when } x - y = 4$$

$$x = 4 \text{ and } y = 8, \text{ when } x - y = -4$$

Thus, the remaining observations are 4 and 8.

Question 2:

The mean and variance of 7 observations are 8 and 16, respectively. If five of the observations are 2, 4, 10, 12 and 14. Find the remaining two observations.

Answer

Let the remaining two observations be x and y .

The observations are 2, 4, 10, 12, 14, x , y .

$$\text{Mean, } \bar{x} = \frac{2+4+10+12+14+x+y}{7} = 8$$

$$\Rightarrow 56 = 42 + x + y$$

$$\Rightarrow x + y = 14 \quad \dots(1)$$

$$\text{Variance} = 16 = \frac{1}{n} \sum_{i=1}^7 (x_i - \bar{x})^2$$

$$16 = \frac{1}{7} [(-6)^2 + (-4)^2 + (2)^2 + (4)^2 + (6)^2 + x^2 + y^2 - 2 \times 8(x+y) + 2 \times (8)^2]$$

$$16 = \frac{1}{7} [36 + 16 + 4 + 16 + 36 + x^2 + y^2 - 16(14) + 2(64)]$$

...[Using (1)]

$$16 = \frac{1}{7} [108 + x^2 + y^2 - 224 + 128]$$

$$16 = \frac{1}{7} [12 + x^2 + y^2]$$

$$\Rightarrow x^2 + y^2 = 112 - 12 = 100$$

$$x^2 + y^2 = 100 \quad \dots(2)$$

From (1), we obtain

$$x^2 + y^2 + 2xy = 196 \quad \dots (3)$$

From (2) and (3), we obtain

$$2xy = 196 - 100$$

$$\Rightarrow 2xy = 96 \quad \dots (4)$$

Subtracting (4) from (2), we obtain

$$x^2 + y^2 - 2xy = 100 - 96$$

$$\Rightarrow (x - y)^2 = 4$$

$$\Rightarrow x - y = \pm 2 \dots (5)$$

Therefore, from (1) and (5), we obtain

$$x = 8 \text{ and } y = 6 \text{ when } x - y = 2$$

$$x = 6 \text{ and } y = 8 \text{ when } x - y = -2$$

Thus, the remaining observations are 6 and 8.

Question 3:

The mean and standard deviation of six observations are 8 and 4, respectively. If each observation is multiplied by 3, find the new mean and new standard deviation of the resulting observations.

Answer

Let the observations be $x_1, x_2, x_3, x_4, x_5,$ and x_6 .

It is given that mean is 8 and standard deviation is 4.

$$\text{Mean, } \bar{x} = \frac{x_1 + x_2 + x_3 + x_4 + x_5 + x_6}{6} = 8 \quad \dots(1)$$

If each observation is multiplied by 3 and the resulting observations are y_i , then

$$y_i = 3x_i \text{ i.e., } x_i = \frac{1}{3}y_i, \text{ for } i = 1 \text{ to } 6$$

$$\begin{aligned} \therefore \text{New mean, } \bar{y} &= \frac{y_1 + y_2 + y_3 + y_4 + y_5 + y_6}{6} \\ &= \frac{3(x_1 + x_2 + x_3 + x_4 + x_5 + x_6)}{6} \\ &= 3 \times 8 \\ &= 24 \end{aligned} \quad \dots[\text{Using (1)}]$$

$$\text{Standard deviation, } \sigma = \sqrt{\frac{1}{n} \sum_{i=1}^6 (x_i - \bar{x})^2}$$

$$\therefore (4)^2 = \frac{1}{6} \sum_{i=1}^6 (x_i - \bar{x})^2$$

$$\sum_{i=1}^6 (x_i - \bar{x})^2 = 96 \quad \dots(2)$$

From (1) and (2), it can be observed that,

$$\bar{y} = 3\bar{x}$$

$$\bar{x} = \frac{1}{3}\bar{y}$$

Substituting the values of x_i and \bar{x} in (2), we obtain

$$\sum_{i=1}^6 \left(\frac{1}{3}y_i - \frac{1}{3}\bar{y} \right)^2 = 96$$

$$\Rightarrow \sum_{i=1}^6 (y_i - \bar{y})^2 = 864$$

Therefore, variance of new observations = $\left(\frac{1}{6} \times 864 \right) = 144$

Hence, the standard deviation of new observations is $\sqrt{144} = 12$

Question 4:

Given that \bar{x} is the mean and σ^2 is the variance of n observations $x_1, x_2 \dots x_n$. Prove that the mean and variance of the observations $ax_1, ax_2, ax_3 \dots ax_n$ are $a\bar{x}$ and $a^2 \sigma^2$, respectively ($a \neq 0$).

Answer

The given n observations are $x_1, x_2 \dots x_n$.

Mean = \bar{x}

Variance = σ^2

$$\therefore \sigma^2 = \frac{1}{n} \sum_{i=1}^n y_i (x_i - \bar{x})^2 \quad \dots(1)$$

If each observation is multiplied by a and the new observations are y_i , then

$$y_i = ax_i \text{ i.e., } x_i = \frac{1}{a} y_i$$

$$\therefore \bar{y} = \frac{1}{n} \sum_{i=1}^n y_i = \frac{1}{n} \sum_{i=1}^n ax_i = \frac{a}{n} \sum_{i=1}^n x_i = a\bar{x} \quad \left(\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i \right)$$

Therefore, mean of the observations, $ax_1, ax_2 \dots ax_n$, is $a\bar{x}$.

Substituting the values of x_i and \bar{x} in (1), we obtain

$$\sigma^2 = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{a} y_i - \frac{1}{a} \bar{y} \right)^2$$

$$\Rightarrow a^2 \sigma^2 = \frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2$$

Thus, the variance of the observations, $ax_1, ax_2 \dots ax_n$, is $a^2 \sigma^2$.

Question 5:

The mean and standard deviation of 20 observations are found to be 10 and 2, respectively. On rechecking, it was found that an observation 8 was incorrect. Calculate the correct mean and standard deviation in each of the following cases:

- (i) If wrong item is omitted.
- (ii) If it is replaced by 12.

Answer

(i) Number of observations (n) = 20

Incorrect mean = 10

Incorrect standard deviation = 2

$$\bar{x} = \frac{1}{n} \sum_{i=1}^{20} x_i$$

$$10 = \frac{1}{20} \sum_{i=1}^{20} x_i$$

$$\Rightarrow \sum_{i=1}^{20} x_i = 200$$

That is, incorrect sum of observations = 200

Correct sum of observations = 200 - 8 = 192

$$\therefore \text{Correct mean} = \frac{\text{Correct sum}}{19} = \frac{192}{19} = 10.1$$

$$\text{Standard deviation } \sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - \frac{1}{n^2} \left(\sum_{i=1}^n x_i \right)^2} = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - (\bar{x})^2}$$

$$\Rightarrow 2 = \sqrt{\frac{1}{20} \text{Incorrect } \sum_{i=1}^n x_i^2 - (10)^2}$$

$$\Rightarrow 4 = \frac{1}{20} \text{Incorrect } \sum_{i=1}^n x_i^2 - 100$$

$$\Rightarrow \text{Incorrect } \sum_{i=1}^n x_i^2 = 2080$$

$$\begin{aligned} \therefore \text{Correct } \sum_{i=1}^n x_i^2 &= \text{Incorrect } \sum_{i=1}^n x_i^2 - (8)^2 \\ &= 2080 - 64 \\ &= 2016 \end{aligned}$$

$$\begin{aligned} \therefore \text{Correct standard deviation} &= \sqrt{\frac{\text{Correct } \sum x_i^2}{n} - (\text{Correct mean})^2} \\ &= \sqrt{\frac{2016}{19} - (10.1)^2} \\ &= \sqrt{106.1 - 102.01} \\ &= \sqrt{4.09} \\ &= 2.02 \end{aligned}$$

(ii) When 8 is replaced by 12,

Incorrect sum of observations = 200

\therefore Correct sum of observations = $200 - 8 + 12 = 204$

$$\therefore \text{Correct mean} = \frac{\text{Correct sum}}{20} = \frac{204}{20} = 10.2$$

$$\text{Standard deviation } \sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - \frac{1}{n^2} \left(\sum_{i=1}^n x_i \right)^2} = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - (\bar{x})^2}$$

$$\Rightarrow 2 = \sqrt{\frac{1}{20} \text{Incorrect } \sum_{i=1}^n x_i^2 - (10)^2}$$

$$\Rightarrow 4 = \frac{1}{20} \text{Incorrect } \sum_{i=1}^n x_i^2 - 100$$

$$\Rightarrow \text{Incorrect } \sum_{i=1}^n x_i^2 = 2080$$

$$\begin{aligned} \therefore \text{Correct } \sum_{i=1}^n x_i^2 &= \text{Incorrect } \sum_{i=1}^n x_i^2 - (8)^2 + (12)^2 \\ &= 2080 - 64 + 144 \\ &= 2160 \end{aligned}$$

$$\begin{aligned} \therefore \text{Correct standard deviation} &= \sqrt{\frac{\text{Correct } \sum_{i=1}^n x_i^2}{n} - (\text{Correct mean})^2} \\ &= \sqrt{\frac{2160}{20} - (10.2)^2} \\ &= \sqrt{108 - 104.04} \\ &= \sqrt{3.96} \\ &= 1.98 \end{aligned}$$

Question 6:

The mean and standard deviation of marks obtained by 50 students of a class in three subjects, Mathematics, Physics and Chemistry are given below:

Subject	Mathematics	Physics	Chemistry
Mean	42	32	40.9
Standard deviation	12	15	20

Which of the three subjects shows the highest variability in marks and which shows the lowest?

Answer

Standard deviation of Mathematics = 12

Standard deviation of Physics = 15

Standard deviation of Chemistry = 20

The coefficient of variation (C.V.) is given by $\frac{\text{Standard deviation}}{\text{Mean}} \times 100$.

$$\text{C.V. (in Mathematics)} = \frac{12}{42} \times 100 = 28.57$$

$$\text{C.V. (in Physics)} = \frac{15}{32} \times 100 = 46.87$$

$$\text{C.V. (in Chemistry)} = \frac{20}{40.9} \times 100 = 48.89$$

The subject with greater C.V. is more variable than others.

Therefore, the highest variability in marks is in Chemistry and the lowest variability in marks is in Mathematics.

Question 7:

The mean and standard deviation of a group of 100 observations were found to be 20 and 3, respectively. Later on it was found that three observations were incorrect, which were recorded as 21, 21 and 18. Find the mean and standard deviation if the incorrect observations are omitted.

Answer

Number of observations (n) = 100

Incorrect mean $(\bar{x}) = 20$

Incorrect standard deviation $(\sigma) = 3$

$$\Rightarrow 20 = \frac{1}{100} \sum_{i=1}^{100} x_i$$

$$\Rightarrow \sum_{i=1}^{100} x_i = 20 \times 100 = 2000$$

\therefore Incorrect sum of observations = 2000

\Rightarrow Correct sum of observations = 2000 - 21 - 21 - 18 = 2000 - 60 = 1940

$$\therefore \text{Correct mean} = \frac{\text{Correct sum}}{100 - 3} = \frac{1940}{97} = 20$$

$$\text{Standard deviation } (\sigma) = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i - \frac{1}{n^2} \left(\sum_{i=1}^n x_i \right)^2} = \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2 - (\bar{x})^2}$$

$$\Rightarrow 3 = \sqrt{\frac{1}{100} \times \text{Incorrect } \sum x_i^2 - (20)^2}$$

$$\Rightarrow \text{Incorrect } \sum x_i^2 = 100(9 + 400) = 40900$$

$$\begin{aligned} \text{Correct } \sum_{i=1}^n x_i^2 &= \text{Incorrect } \sum_{i=1}^n x_i^2 - (21)^2 - (21)^2 - (18)^2 \\ &= 40900 - 441 - 441 - 324 \\ &= 39694 \end{aligned}$$

$$\begin{aligned} \therefore \text{Correct standard deviation} &= \sqrt{\frac{\text{Correct } \sum x_i^2}{n} - (\text{Correct mean})^2} \\ &= \sqrt{\frac{39694}{97} - (20)^2} \\ &= \sqrt{409.216 - 400} \\ &= \sqrt{9.216} \\ &= 3.036 \end{aligned}$$