

CLASS IX: MATHS
Chapter 1: Number System

Questions and Solutions | EXERCISE 1.4 - NCERT Books

Q1. Classify the following numbers as rational or irrational :

(i) $2 - \sqrt{5}$ (ii) $(3 + \sqrt{23}) - \sqrt{23}$ (iii) $\frac{2\sqrt{7}}{7\sqrt{7}}$

(iv) $\frac{1}{\sqrt{2}}$ (v) 2π

Sol. (i) $\because 2$ is a rational number and $\sqrt{5}$ is an irrational number.

$\therefore 2 - \sqrt{5}$ is an irrational number.

(ii) $(3 + \sqrt{23}) - \sqrt{23} \Rightarrow (3 + \sqrt{23}) - \sqrt{23} = 3$ is a rational number.

(iii) $\frac{2\sqrt{7}}{7\sqrt{7}} = \frac{2}{7}$ Rational number.

(iv) $\frac{1}{\sqrt{2}}$

$\because 1$ is a rational number and $\sqrt{2}$ is an irrational number.

So, $\frac{1}{\sqrt{2}}$ is irrational number.

(v) 2π

$\because 2$ is a rational number and π is an irrational number

So, 2π is irrational number.

Q2. Simplify each of the following expressions :

(i) $(3 + \sqrt{3})(2 + \sqrt{2})$

(ii) $(3 + \sqrt{3})(3 - \sqrt{3})$

(iii) $(\sqrt{5} + \sqrt{2})^2$

(iv) $(\sqrt{5} - \sqrt{2})(\sqrt{5} + \sqrt{2})$

Sol. (i) $(3 + \sqrt{3})(2 + \sqrt{2}) = 3(2 + \sqrt{2}) + \sqrt{3}(2 + \sqrt{2})$

$$= 6 + 3\sqrt{2} + 2\sqrt{3} + \sqrt{6}$$

(ii) $(3 + \sqrt{3})(3 - \sqrt{3}) = (3)^2 - (\sqrt{3})^2 = 9 - 3 = 6$

(iii) $(\sqrt{5} + \sqrt{2})^2$

$$= (\sqrt{5})^2 + 2\sqrt{10} + (\sqrt{2})^2$$


$$= 7 + 2\sqrt{10}$$

(iv) $(\sqrt{5} - \sqrt{2})(\sqrt{5} + \sqrt{2}) = 5 - 2 = 3$

Q3. Recall, π is defined as the ratio of the circumference (say c) of a circle to its diameter (say d). That is, $\pi = c/d$. This seems to contradict the fact that π is irrational. How will you resolve this contradiction ?

Sol. There is no contradiction. When we measure a length with a scale or any other device, we only get an approximate rational value. Therefore, we may not realise that c or d is irrational.

Q4. Represent $\sqrt{9.3}$ on the number line.



Let l be the number line.

Draw a line segment $AB = 9.3$ units and $BC = 1$ unit. Find the mid point O of AC .

Draw a semicircle with centre O and radius OA or OC .

Draw $BD \perp AC$ intersecting the semicircle at D . Then, $BD = \sqrt{9.3}$ units. Now, with centre B and radius BD , draw an arc intersecting the number line l at P .

Hence, $BD = BP = \sqrt{9.3}$

Q5. Rationalise the denominators of the following :

(i) $\frac{1}{\sqrt{7}}$

(ii) $\frac{1}{\sqrt{7} - \sqrt{6}}$

(iii) $\frac{1}{\sqrt{5} + \sqrt{2}}$

(iv) $\frac{1}{\sqrt{7} - 2}$

Sol. (i) $\frac{1}{\sqrt{7}} = \frac{1}{\sqrt{7}} \times \frac{\sqrt{7}}{\sqrt{7}} = \frac{\sqrt{7}}{7}$

(ii) $\frac{1}{\sqrt{7} - \sqrt{6}} = \frac{1}{\sqrt{7} - \sqrt{6}} \times \frac{\sqrt{7} + \sqrt{6}}{\sqrt{7} + \sqrt{6}}$



$$= \frac{\sqrt{7} + \sqrt{6}}{7 - 6} = \frac{\sqrt{7} + \sqrt{6}}{1} = \sqrt{7} + \sqrt{6}$$

$$(iii) \frac{1}{\sqrt{5} + \sqrt{2}}$$

$$\frac{1}{\sqrt{5} + \sqrt{2}} \times \frac{\sqrt{5} - \sqrt{2}}{\sqrt{5} - \sqrt{2}} = \frac{\sqrt{5} - \sqrt{2}}{3}$$

$$(iv) \frac{1}{\sqrt{7} - 2} = \frac{1}{\sqrt{7} - 2} \times \frac{\sqrt{7} + 2}{\sqrt{7} + 2}$$

$$= \frac{\sqrt{7} + 2}{7 - 4} = \frac{\sqrt{7} + 2}{3}$$

