

## FINAL JEE-MAIN EXAMINATION – MARCH, 2021

(Held On Thursday 18<sup>th</sup> March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

### MATHEMATICS

#### SECTION-A

1. Let  $y = y(x)$  be the solution of the differential equation  $\frac{dy}{dx} = (y+1)((y+1)e^{x^2/2} - x)$ ,  $0 < x < 2.1$ ,

80

with  $y(2) = 0$ . Then the value of  $\frac{dy}{dx}$  at

$x = 1$  is equal to :

- (1)  $\frac{-e^{3/2}}{(e^2 + 1)^2}$                       (2)  $-\frac{2e^2}{(1 + e^2)^2}$   
 (3)  $\frac{e^{5/2}}{(1 + e^2)^2}$                       (4)  $\frac{5e^{1/2}}{(e^2 + 1)^2}$

**Official Ans. by NTA (1)**

- Sol.** Let  $y + 1 = Y$

$$\therefore \frac{dY}{dx} = Y^2 e^{\frac{x^2}{2}} - xY$$

Put  $-\frac{1}{Y} = k$

$$\Rightarrow \frac{dk}{dx} + k(-x) = e^{\frac{x^2}{2}}$$

I.F. =  $e^{-\frac{x^2}{2}}$

$$\therefore k = (x + c)e^{x^2/2}$$

Put  $k = -\frac{1}{y+1}$

$$\therefore y + 1 = -\frac{1}{(x + c)e^{x^2/2}} \quad \dots(i)$$

when  $x = 2, y = 0$ , then  $c = -2 - \frac{1}{e^2}$

differentiate equation (i) & put  $x = 1$

we get  $\left(\frac{dy}{dx}\right)_{x=1} = -\frac{e^{3/2}}{(1 + e^2)^2}$

### TEST PAPER WITH SOLUTION

2. In a triangle ABC, if  $|\vec{BC}| = 8, |\vec{CA}| = 7,$


$$|\vec{AB}| = 10, \text{ then the projection of the vector } \vec{AB}$$

on  $\vec{AC}$  is equal to :

- (1)  $\frac{25}{4}$                       (2)  $\frac{85}{14}$                       (3)  $\frac{127}{20}$                       (4)  $\frac{115}{16}$

**Official Ans. by NTA (2)**

**Sol.**



$$|\vec{a}| = 8, |\vec{b}| = 7, |\vec{c}| = 10$$

$$\cos \theta = \frac{|\vec{b}|^2 + |\vec{c}|^2 - |\vec{a}|^2}{2|\vec{b}||\vec{c}|} = \frac{17}{28}$$

Projection of  $\vec{c}$  on  $\vec{b}$

$$= |\vec{c}| \cos \theta$$

$$= 10 \times \frac{17}{28}$$

$$= \frac{85}{14}$$

3. Let the system of linear equations

$$4x + \lambda y + 2z = 0$$

$$2x - y + z = 0$$

$$\mu x + 2y + 3z = 0, \lambda, \mu \in \mathbb{R}.$$

has a non-trivial solution. Then which of the following is true ?

- (1)  $\mu = 6, \lambda \in \mathbb{R}$                       (2)  $\lambda = 2, \mu \in \mathbb{R}$   
 (3)  $\lambda = 3, \mu \in \mathbb{R}$                       (4)  $\mu = -6, \lambda \in \mathbb{R}$

**Official Ans. by NTA (1)**

**Sol.** For non-trivial solution

$$\begin{vmatrix} 4 & \lambda & 2 \\ 2 & -1 & 1 \\ \mu & 2 & 3 \end{vmatrix} = 0$$

$$\Rightarrow 2\mu - 6\lambda + \lambda\mu = 12$$

when  $\mu = 6$ ,  $12 - 6\lambda + 6\lambda = 12$   
which is satisfied by all  $\lambda$

4. Let  $f : \mathbb{R} - \{3\} \rightarrow \mathbb{R} - \{1\}$  be defined by

$$f(x) = \frac{x-2}{x-3}$$

Let  $g : \mathbb{R} \rightarrow \mathbb{R}$  be given as

$$g(x) = 2x - 3$$

Then, the sum of all the values of  $x$  for which  $f^{-1}(x) + g^{-1}(x) = \frac{13}{2}$  is equal to

- (1) 7      (2) 2      (3) 5      (4) 3

**Official Ans. by NTA (3)**

**Sol.**  $f(x) = y = \frac{x-2}{x-3}$

$$\therefore x = \frac{3y-2}{y-1}$$

$$\therefore f^{-1}(x) = \frac{3x-2}{x-1}$$

&  $g(x) = y = 2x - 3$

$$\therefore x = \frac{y+3}{2}$$

$$\therefore g^{-1}(x) = \frac{x+3}{2}$$

$$\therefore f^{-1}(x) + g^{-1}(x) = \frac{13}{2}$$

$$\therefore x^2 - 5x + 6 = 0 \begin{cases} x_1 \\ x_2 \end{cases}$$

$\therefore$  sum of roots


$$x_1 + x_2 = 5$$

5. Let the centroid of an equilateral triangle ABC be at the origin. Let one of the sides of the equilateral triangle be along the straight line  $x + y = 3$ . If  $R$  and  $r$  be the radius of circumcircle and incircle respectively of  $\Delta ABC$ , then  $(R + r)$  is equal to :

- (1)  $\frac{9}{\sqrt{2}}$       (2)  $7\sqrt{2}$       (3)  $2\sqrt{2}$       (4)  $3\sqrt{2}$

**Official Ans. by NTA (1)**

**Sol.**



$$r = OM = \frac{3}{\sqrt{2}}$$

$$\& \sin 30^\circ = \frac{1}{2} = \frac{r}{R} \Rightarrow R = \frac{6}{\sqrt{2}}$$


$$\therefore r + R = \frac{9}{\sqrt{2}}$$

6. Consider a hyperbola  $H : x^2 - 2y^2 = 4$ . Let the tangent at a point  $P(4, \sqrt{6})$  meet the  $x$ -axis at  $Q$  and latus rectum at  $R(x_1, y_1)$ ,  $x_1 > 0$ . If  $F$  is a focus of  $H$  which is nearer to the point  $P$ , then the area of  $\Delta QFR$  is equal to

- (1)  $4\sqrt{6}$       (2)  $\sqrt{6} - 1$   
(3)  $\frac{7}{\sqrt{6}} - 2$       (4)  $4\sqrt{6} - 1$

**Official Ans. by NTA (3)**

**Sol.**



$$\frac{x^2}{4} - \frac{y^2}{2} = 1$$

$$e = \sqrt{1 + \frac{b^2}{a^2}} = \sqrt{\frac{3}{2}}$$

$$\therefore \text{Focus } F(ae, 0) \Rightarrow F(\sqrt{6}, 0)$$

equation of tangent at P to the hyperbola is

$$2x - y\sqrt{6} = 2$$

tangent meet x-axis at Q(1, 0)

$$\& \text{ latus rectum } x = \sqrt{6} \text{ at } R\left(\sqrt{6}, \frac{2}{\sqrt{6}}(\sqrt{6}-1)\right)$$

$$\therefore \text{Area of } \Delta_{QFR} = \frac{1}{2}(\sqrt{6}-1) \cdot \frac{2}{\sqrt{6}}(\sqrt{6}-1)$$

$$= \frac{7}{\sqrt{6}} - 2$$

7. If P and Q are two statements, then which of the following compound statement is a tautology ?

- (1)  $(P \Rightarrow Q) \wedge \sim Q \Rightarrow Q$
- (2)  $(P \Rightarrow Q) \wedge \sim Q \Rightarrow \sim P$
- (3)  $(P \Rightarrow Q) \wedge \sim Q \Rightarrow P$
- (4)  $(P \Rightarrow Q) \wedge \sim Q \Rightarrow (P \wedge Q)$


**Official Ans. by NTA (2)**

**Sol.** LHS of all the options are same i.e.

$$\begin{aligned} (P \rightarrow Q) \wedge \sim Q & \\ \equiv (\sim P \vee Q) \wedge \sim Q & \\ \equiv (\sim P \wedge \sim Q) \vee (Q \wedge \sim Q) & \\ \equiv \sim P \wedge \sim Q & \end{aligned}$$

$$\begin{aligned} \text{(A) } (\sim P \wedge \sim Q) \rightarrow Q & \\ \equiv \sim(\sim P \wedge \sim Q) \vee Q & \\ \equiv (P \vee Q) \vee Q \neq \text{tautology} & \end{aligned}$$

$$\begin{aligned} \text{(B) } (\sim P \wedge \sim Q) \rightarrow \sim P & \\ \equiv \sim(\sim P \wedge \sim Q) \vee \sim P & \\ \equiv (P \vee Q) \vee \sim P & \end{aligned}$$



$\Rightarrow$  Tautology

$$\text{(C) } (\sim P \wedge \sim Q) \rightarrow P$$

$$\equiv (P \vee Q) \vee P \neq \text{Tautology}$$

$$\text{(D) } (\sim P \wedge \sim Q) \rightarrow (P \wedge Q)$$

$$\equiv (P \vee Q) \vee (P \wedge Q) \neq \text{Tautology}$$

**Aliter :**

P	Q	$P \vee Q$	$P \wedge Q$	$\sim P$	$(P \vee Q) \vee \sim P$
T	T	T	T	F	T
T	F	T	F	F	T
F	T	T	F	T	T
F	F	F	F	T	T

8. Let  $g(x) = \int_0^x f(t) dt$ , where f is continuous

function in  $[0, 3]$  such that  $\frac{1}{3} \leq f(t) \leq 1$  for all

$t \in [0, 1]$  and  $0 \leq f(t) \leq \frac{1}{2}$  for all  $t \in (1, 3]$ .

The largest possible interval in which g(3) lies is :

- (1)  $\left[-1, -\frac{1}{2}\right]$
- (2)  $\left[-\frac{3}{2}, -1\right]$
- (3)  $\left[\frac{1}{3}, 2\right]$
- (4)  $[1, 3]$

**Official Ans. by NTA (3)**

$$\text{Sol. } \frac{1}{3} \leq f(t) \leq 1 \forall t \in [0, 1]$$

$$0 \leq f(t) \leq \frac{1}{2} \forall t \in (1, 3]$$

$$\text{Now, } g(3) = \int_0^3 f(t) dt = \int_0^1 f(t) dt + \int_1^3 f(t) dt$$

$$\therefore \int_0^1 \frac{1}{3} dt \leq \int_0^1 f(t) dt \leq \int_0^1 1 dt \quad \dots(1)$$

$$\text{and } \int_1^3 0 dt \leq \int_1^3 f(t) dt \leq \int_1^3 \frac{1}{2} dt \quad \dots(2)$$

Adding, we get

$$\frac{1}{3} + 0 \leq g(3) \leq 1 + \frac{1}{2}(3-1)$$

$$\frac{1}{3} \leq g(3) \leq 2$$

9. Let  $S_1$  be the sum of first  $2n$  terms of an arithmetic progression. Let  $S_2$  be the sum of first  $4n$  terms of the same arithmetic progression. If  $(S_2 - S_1)$  is 1000, then the sum of the first  $6n$  terms of the arithmetic progression is equal to:  
 (1) 1000 (2) 7000 (3) 5000 (4) 3000  
**Official Ans. by NTA (4)**

**Sol.**  $S_{2n} = \frac{2n}{2} [2a + (2n - 1)d]$ ,  $S_{4n} = \frac{4n}{2} [2a + (4n - 1)d]$

$$\Rightarrow S_2 - S_1 = \frac{4n}{2} [2a + (4n - 1)d] - \frac{2n}{2} [2a + (2n - 1)d]$$

$$= 4an + (4n - 1)2nd - 2na - (2n - 1)dn$$

$$= 2na + nd[8n - 2 - 2n + 1]$$

$$\Rightarrow 2na + nd[6n - 1] = 1000$$

$$2a + (6n - 1)d = \frac{1000}{n}$$

Now,  $S_{6n} = \frac{6n}{2} [2a + (6n - 1)d]$

$$= 3n \cdot \frac{1000}{n} = 3000$$


10. Let a complex number be  $w = 1 - \sqrt{3}i$ . Let another complex number  $z$  be such that  $|zw| = 1$  and  $\arg(z) - \arg(w) = \frac{\pi}{2}$ . Then the area of the triangle with vertices origin,  $z$  and  $w$  is equal to :  
 (1) 4 (2)  $\frac{1}{2}$  (3)  $\frac{1}{4}$  (4) 2

**Official Ans. by NTA (2)**

**Sol.**  $w = 1 - \sqrt{3}i \Rightarrow |w| = 2$

Now,  $|z| = \frac{1}{|w|} \Rightarrow |z| = \frac{1}{2}$

and  $\arg(z) = \frac{\pi}{2} + \arg(w)$



$$\Rightarrow \text{Area of triangle} = \frac{1}{2} \cdot OP \cdot OQ$$

$$= \frac{1}{2} \cdot 2 \cdot \frac{1}{2} = \frac{1}{2}$$

11. Let in a series of  $2n$  observations, half of them are equal to  $a$  and remaining half are equal to  $-a$ . Also by adding a constant  $b$  in each of these observations, the mean and standard deviation of new set become 5 and 20, respectively. Then the value of  $a^2 + b^2$  is equal to :  
 (1) 425 (2) 650 (3) 250 (4) 925  
**Official Ans. by NTA (1)**

**Sol.** Let observations are denoted by  $x_i$  for  $1 \leq i < 2n$

$$\bar{x} = \frac{\sum x_i}{2n} = \frac{(a + a + \dots + a) - (a + a + \dots + a)}{2n}$$

$$\Rightarrow \bar{x} = 0$$

$$\text{and } \sigma_x^2 = \frac{\sum x_i^2}{2n} - (\bar{x})^2 = \frac{a^2 + a^2 + \dots + a^2}{2n} - 0 = a^2$$

$$\Rightarrow \sigma_x = a$$

Now, adding a constant  $b$  then  $\bar{y} = \bar{x} + b = 5$

$$\Rightarrow b = 5$$

and  $\sigma_y = \sigma_x$  (No change in S.D.)  $\Rightarrow a = 20$

$$\Rightarrow a^2 + b^2 = 425$$

12. Let  $S_1 : x^2 + y^2 = 9$  and  $S_2 : (x - 2)^2 + y^2 = 1$ . Then the locus of center of a variable circle  $S$  which touches  $S_1$  internally and  $S_2$  externally always passes through the points :

(1)  $(0, \pm\sqrt{3})$                       (2)  $\left(\frac{1}{2}, \pm\frac{\sqrt{5}}{2}\right)$


(3)  $\left(2, \pm\frac{3}{2}\right)$                       (4)  $(1, \pm 2)$

**Official Ans. by NTA (3)**

**Sol.**  $S_1 : x^2 + y^2 = 9 \begin{cases} r_1 = 3 \\ A(0, 0) \end{cases}$

$S_2 : (x - 2)^2 + y^2 = 1 \begin{cases} r_2 = 1 \\ B(2, 0) \end{cases}$

$\therefore c_1 c_2 = r_1 - r_2$



$\therefore$  given circle are touching internally

Let a variable circle with centre  $P$  and radius  $r$

$\Rightarrow PA = r_1 - r$  and  $PB = r_2 + r$


$\Rightarrow PA + PB = r_1 + r_2$

$\Rightarrow PA + PB = 4$  ( $> AB$ )

$\Rightarrow$  Locus of  $P$  is an ellipse with foci at  $A(0, 0)$  and  $B(2, 0)$  and length of major axis is  $2a = 4$ ,

$e = \frac{1}{2}$

$\Rightarrow$  centre is at  $(1, 0)$  and  $b^2 = a^2(1 - e^2) = 3$  if  $x$ -ellipse



$\Rightarrow E: \frac{(x-1)^2}{4} + \frac{y^2}{3} = 1$

which is satisfied by  $\left(2, \pm\frac{3}{2}\right)$

13. Let  $\vec{a}$  and  $\vec{b}$  be two non-zero vectors perpendicular to each other and  $|\vec{a}| = |\vec{b}|$ . If

$|\vec{a} \times \vec{b}| = |\vec{a}|$ , then the angle between the vectors

$(\vec{a} + \vec{b} + (\vec{a} \times \vec{b}))$  and  $\vec{a}$  is equal to :

(1)  $\sin^{-1}\left(\frac{1}{\sqrt{3}}\right)$                       (2)  $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$

(3)  $\cos^{-1}\left(\frac{1}{\sqrt{2}}\right)$                       (4)  $\sin^{-1}\left(\frac{1}{\sqrt{6}}\right)$

**Official Ans. by NTA (2)**

**Sol.**  $|\vec{a}| = |\vec{b}|$ ,  $|\vec{a} \times \vec{b}| = |\vec{a}|$ ,  $\vec{a} \perp \vec{b}$

$|\vec{a} \times \vec{b}| = |\vec{a}| \Rightarrow |\vec{a}||\vec{b}|\sin 90^\circ = |\vec{a}| \Rightarrow |\vec{b}| = 1 = |\vec{a}|$

$\vec{a}$  and  $\vec{b}$  are mutually perpendicular unit vectors.

Let  $\vec{a} = \hat{i}$ ,  $\vec{b} = \hat{j} \Rightarrow \vec{a} \times \vec{b} = \hat{k}$

$\cos \theta = \frac{(\hat{i} + \hat{j} + \hat{k}) \cdot \hat{i}}{\sqrt{3} \sqrt{1}} = \frac{1}{\sqrt{3}} \Rightarrow \theta = \cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$

14. Let in a Binomial distribution, consisting of 5 independent trials, probabilities of exactly 1 and 2 successes be 0.4096 and 0.2048 respectively. Then the probability of getting exactly 3 successes is equal to :

(1)  $\frac{32}{625}$                       (2)  $\frac{80}{243}$                       (3)  $\frac{40}{243}$                       (4)  $\frac{128}{625}$

**Official Ans. by NTA (1)**

**Sol.**  $P(X = 1) = {}^5C_1 \cdot p \cdot q^4 = 0.4096$

$P(X = 2) = {}^5C_2 \cdot p^2 \cdot q^3 = 0.2048$

$\Rightarrow \frac{q}{2p} = 2$

$\Rightarrow q = 4p$  and  $p + q = 1$

$\Rightarrow p = \frac{1}{5}$  and  $q = \frac{4}{5}$

Now

$P(X = 3) = {}^5C_3 \cdot \left(\frac{1}{5}\right)^3 \cdot \left(\frac{4}{5}\right)^2 = \frac{10 \times 16}{125 \times 25} = \frac{32}{625}$

15. Let a tangent be drawn to the ellipse  $\frac{x^2}{27} + y^2 = 1$

at  $(3\sqrt{3}\cos\theta, \sin\theta)$  where  $\theta \in \left(0, \frac{\pi}{2}\right)$ . Then the

value of  $\theta$  such that the sum of intercepts on axes made by this tangent is minimum is equal to :

- 80  
 (1)  $\frac{\pi}{8}$       (2)  $\frac{\pi}{4}$       (3)  $\frac{\pi}{6}$       (4)  $\frac{\pi}{3}$

**Official Ans. by NTA (3)**

**Sol.** Equation of tangent be

$$\frac{x \cos \theta}{3\sqrt{3}} + \frac{y \sin \theta}{1} = 1, \quad \theta \in \left(0, \frac{\pi}{2}\right)$$

intercept on x-axis

$$OA = 3\sqrt{3} \sec \theta$$

intercept on y-axis

$$OB = \operatorname{cosec} \theta$$


Now, sum of intercept

$$= 3\sqrt{3} \sec \theta + \operatorname{cosec} \theta = f(\theta) \text{ let}$$

$$f'(\theta) = 3\sqrt{3} \sec \theta \tan \theta - \operatorname{cosec} \theta \cot \theta$$

$$= 3\sqrt{3} \frac{\sin \theta}{\cos^2 \theta} - \frac{\cos \theta}{\sin^2 \theta}$$

$$= \underbrace{\frac{\cos \theta}{\sin^2 \theta}}_{\oplus} \cdot 3\sqrt{3} \left[ \tan^3 \theta - \frac{1}{3\sqrt{3}} \right] = 0 \Rightarrow \theta = \frac{\pi}{6}$$



$\Rightarrow$  at  $\theta = \frac{\pi}{6}$ ,  $f(\theta)$  is minimum

16. Define a relation R over a class of  $n \times n$  real matrices A and B as "ARB iff there exists a non-singular matrix P such that  $PAP^{-1} = B$ ".

Then which of the following is true ?

- (1) R is symmetric, transitive but not reflexive,  
 (2) R is reflexive, symmetric but not transitive  
 (3) R is an equivalence relation  
 (4) R is reflexive, transitive but not symmetric

**Official Ans. by NTA (3)**

**Sol.** A and B are matrices of  $n \times n$  order & ARB iff there exists a non singular matrix P ( $\det(P) \neq 0$ ) such that  $PAP^{-1} = B$

**For reflexive**

$ARA \Rightarrow PAP^{-1} = A$  ... (1) must be true for  $P = I$ , Eq.(1) is true so 'R' is reflexive

**For symmetric**

$ARB \Leftrightarrow PAP^{-1} = B$  ... (1) is true for BRA iff  $PBP^{-1} = A$  ... (2) must be true

$$\therefore PAP^{-1} = B$$

$$P^{-1}PAP^{-1} = P^{-1}B$$

$$IAP^{-1}P = P^{-1}BP$$

$$A = P^{-1}BP \quad \dots (3)$$

$$\text{from (2) \& (3) } PBP^{-1} = P^{-1}BP$$

can be true some  $P = P^{-1} \Rightarrow P^2 = I$  ( $\det(P) \neq 0$ )

So 'R' is symmetric

**For transitive**

$$ARB \Leftrightarrow PAP^{-1} = B \dots \text{ is true}$$

$$BRC \Leftrightarrow PBP^{-1} = C \dots \text{ is true}$$

$$\text{now } PPAP^{-1}P^{-1} = C$$

$$P^2A(P^2)^{-1} = C \Rightarrow ARC$$

So 'R' is transitive relation

$\Rightarrow$  Hence R is equivalence

17. A pole stands vertically inside a triangular park ABC. Let the angle of elevation of the top of

the pole from each corner of the park be  $\frac{\pi}{3}$ .

If the radius of the circumcircle of  $\Delta ABC$  is 2, then the height of the pole is equal to :

- (1)  $\frac{2\sqrt{3}}{3}$       (2)  $2\sqrt{3}$       (3)  $\sqrt{3}$       (4)  $\frac{1}{\sqrt{3}}$

**Official Ans. by NTA (2)**

**Sol.** Let  $PD = h$ ,  $R = 2$


As angle of elevation

of top of pole from

A, B, C are equal So

D must be circumcentre

of  $\Delta ABC$



$$\tan\left(\frac{\pi}{3}\right) = \frac{PD}{R} = \frac{h}{R}$$

$$h = R \tan\left(\frac{\pi}{3}\right) = 2\sqrt{3}$$

18. If  $15\sin^4\alpha + 10\cos^4\alpha = 6$ , for some  $\alpha \in \mathbb{R}$ , then the value of  $27\sec^6\alpha + 8\operatorname{cosec}^6\alpha$  is equal to :  
 (1) 350 (2) 500 (3) 400 (4) 250

**Official Ans. by NTA (4)**

**Sol.**  $15\sin^4\alpha + 10\cos^4\alpha = 6$   
 $15\sin^4\alpha + 10\cos^4\alpha = 6(\sin^2\alpha + \cos^2\alpha)^2$   
 $(3\sin^2\alpha - 2\cos^2\alpha)^2 = 0$

$\tan^2\alpha = \frac{2}{3}$ ,  $\cot^2\alpha = \frac{3}{2}$   
 $\Rightarrow 27\sec^6\alpha + 8\operatorname{cosec}^6\alpha$   
 $= 27(\sec^2\alpha)^3 + 8(\operatorname{cosec}^2\alpha)^3$   
 $= 27(1 + \tan^2\alpha)^3 + 8(1 + \cot^2\alpha)^3$   
 $= 250$

19. The area bounded by the curve  $4y^2 = x^2(4-x)(x-2)$  is equal to :

(1)  $\frac{\pi}{8}$  (2)  $\frac{3\pi}{8}$  (3)  $\frac{3\pi}{2}$  (4)  $\frac{\pi}{16}$

**Official Ans. by NTA (3)**

**Sol.**  $4y^2 = x^2(4-x)(x-2)$

$|y| = \frac{|x|}{2} \sqrt{(4-x)(x-2)}$

$\Rightarrow y_1 = \frac{x}{2} \sqrt{(4-x)(x-2)}$

and  $y_2 = -\frac{x}{2} \sqrt{(4-x)(x-2)}$

D :  $x \in [2, 4]$

Required Area

$= \int_2^4 (y_1 - y_2) dx = \int_2^4 x \sqrt{(4-x)(x-2)} dx \dots(1)$


Applying  $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$

Area =  $\int_2^4 (6-x) \sqrt{(4-x)(x-2)} dx \dots(2)$

(1) + (2)

$2A = 6 \int_2^4 \sqrt{(4-x)(x-2)} dx$

$A = 3 \int_2^4 \sqrt{1-(x-3)^2} dx$



$A = 3 \cdot \frac{\pi}{2} \cdot 1^2 = \frac{3\pi}{2}$

20. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined as

$$f(x) = \begin{cases} \frac{\sin(a+1)x + \sin 2x}{2x} & , \text{if } x < 0 \\ b & , \text{if } x = 0 \\ \frac{\sqrt{x+bx^3} - \sqrt{x}}{bx^{5/2}} & , \text{if } x > 0 \end{cases}$$

If  $f$  is continuous at  $x = 0$ , then the value of  $a + b$  is equal to :

(1)  $-\frac{5}{2}$  (2)  $-2$  (3)  $-3$  (4)  $-\frac{3}{2}$

**Official Ans. by NTA (4)**

**Sol.**  $f(x)$  is continuous at  $x = 0$

$\lim_{x \rightarrow 0^+} f(x) = f(0) = \lim_{x \rightarrow 0^-} f(x) \dots(1)$

$f(0) = b \dots(2)$

$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \left( \frac{\sin(a+1)x}{2x} + \frac{\sin 2x}{2x} \right)$   
 $= \frac{a+1}{2} + 1 \dots(3)$

$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{\sqrt{x+bx^3} - \sqrt{x}}{bx^{5/2}}$   
 $= \lim_{x \rightarrow 0^+} \frac{(x+bx^3-x)}{bx^{5/2}(\sqrt{x+bx^3} + \sqrt{x})}$   
 $= \lim_{x \rightarrow 0^+} \frac{\sqrt{x}}{\sqrt{x}(\sqrt{1+bx^2} + 1)} = \frac{1}{2} \dots(4)$

Use (2), (3) & (4) in (1)

$\frac{1}{2} = b = \frac{a+1}{2} + 1$

$\Rightarrow b = \frac{1}{2}, a = -2$

$a + b = \frac{-3}{2}$

**SECTION-B**

1. If  $f(x)$  and  $g(x)$  are two polynomials such that the polynomial  $P(x) = f(x^3) + xg(x^3)$  is divisible by  $x^2 + x + 1$ , then  $P(1)$  is equal to\_\_\_\_\_.

**Official Ans. by NTA (0)**

**Sol.**  $P(x) = f(x^3) + xg(x^3)$   
 $P(1) = f(1) + g(1) \dots(1)$   
 Now  $P(x)$  is divisible by  $x^2 + x + 1$   
 $\Rightarrow P(x) = Q(x)(x^2 + x + 1)$   
 $P(w) = 0 = P(w^2)$  where  $w, w^2$  are non-real cube roots of units  
 $P(x) = f(x^3) + xg(x^3)$   
 $P(w) = f(w^3) + wg(w^3) = 0$   
 $f(1) + wg(1) = 2 \dots(2)$   
 $P(w^2) = f(w^6) + w^2g(w^6) = 0$   
 $f(1) + w^2g(1) = 0 \dots(3)$   
 $(2) + (3)$   
 $\Rightarrow 2f(1) + (w + w^2)g(1) = 0$   
 $2f(1) = g(1) \dots(4)$   
 $(2) - (3)$   
 $\Rightarrow (w - w^2)g(1) = 0$   
 $g(1) = 0 = f(1)$  from (4)  
 from (1)  $P(1) = f(1) + g(1) = 0$

2. Let  $I$  be an identity matrix of order  $2 \times 2$  and

$P = \begin{bmatrix} 2 & -1 \\ 5 & -3 \end{bmatrix}$ . Then the value of  $n \in \mathbb{N}$  for

which  $P^n = 5I - 8P$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (6)**

**Sol.**  $P = \begin{bmatrix} 2 & -1 \\ 5 & -3 \end{bmatrix}$   
 $5I - 8P = \begin{bmatrix} 5 & 0 \\ 0 & 5 \end{bmatrix} - \begin{bmatrix} 16 & -8 \\ 40 & -24 \end{bmatrix} = \begin{bmatrix} -11 & 8 \\ -40 & 29 \end{bmatrix}$   
 $P^2 = \begin{bmatrix} -1 & 1 \\ -5 & 4 \end{bmatrix}$   
 $P^3 = \begin{bmatrix} 3 & -2 \\ 10 & -7 \end{bmatrix} \Rightarrow P^6 = \begin{bmatrix} -11 & 8 \\ -40 & 29 \end{bmatrix} = P^n$   
 $\Rightarrow n = 6$

3. If  $\sum_{r=1}^{10} r!(r^3 + 6r^2 + 2r + 5) = \alpha(11!)$ , then the value of  $\alpha$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (160)**

**Sol.**  $\sum_{r=1}^{10} r!\{(r+1)(r+2)(r+3) - 9(r+1) + 8\}$   
 $= \sum_{r=1}^{10} [ \{(r+3)! - (r+1)!\} - 8\{(r+1)! - r!\} ]$   
 $= (13! + 12! - 2! - 3!) - 8(11! - 1)$   
 $= (12.13 + 12 - 8).11! - 8 + 8$   
 $= (160)(11)!$   
 Hence  $\alpha = 160$

4. The term independent of  $x$  in the expansion of

$\left[ \frac{x+1}{x^{2/3} - x^{1/3} + 1} - \frac{x-1}{x - x^{1/2}} \right]^{10}$ ,  $x \neq 1$ , is equal to

\_\_\_\_\_.

**Official Ans. by NTA (210)**

**Sol.**  $\left( (x^{1/3} + 1) - \left( \frac{\sqrt{x} + 1}{\sqrt{x}} \right) \right)^{10}$   
 $(x^{1/3} - x^{-1/2})^{10}$   
 $T_{r+1} = {}^{10}C_r (x^{1/3})^{10-r} (-x^{-1/2})^r$   
 $\frac{10-r}{3} - \frac{r}{2} = 0 \Rightarrow 20 - 2r - 3r = 0$   
 $\Rightarrow r = 4$

$T_5 = {}^{10}C_4 = \frac{10 \times 9 \times 8 \times 7}{4 \times 3 \times 2 \times 1} = 210$

5. Let  $P(x)$  be a real polynomial of degree 3 which vanishes at  $x = -3$ . Let  $P(x)$  have local minima at  $x = 1$ , local maxima at  $x = -1$  and

$\int_{-1}^1 P(x) dx = 18$ , then the sum of all the coefficients of the polynomial  $P(x)$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (8)**



**Sol.** Let  $p'(x) = a(x-1)(x+1) = a(x^2-1)$

$$p(x) = a \int (x^2-1) dx + c$$

$$= a \left( \frac{x^3}{3} - x \right) + c$$

Now  $p(-3) = 0$

$$\Rightarrow a \left( -\frac{27}{3} + 3 \right) + c = 0$$

$$\Rightarrow -6a + c = 0 \quad \dots(1)$$

$$\text{Now } \int_{-1}^1 \left( a \left( \frac{x^3}{3} - x \right) + c \right) dx = 18$$

$$= 2c = 18 \Rightarrow c = 9 \quad \dots(2)$$

$$\Rightarrow \text{from (1) \& (2)} \Rightarrow -6a + 9 = 0 \Rightarrow a = \frac{3}{2}$$

$$\Rightarrow p(x) = \frac{3}{2} \left( \frac{x^3}{3} - x \right) + 9$$

sum of coefficient


$$= \frac{1}{2} - \frac{3}{2} + 9$$

$$= 8$$

6. Let the mirror image of the point  $(1, 3, a)$  with respect to the plane  $\vec{r} \cdot (2\hat{i} - \hat{j} + \hat{k}) - b = 0$  be  $(-3, 5, 2)$ . Then the value of  $|a + b|$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (1)**

**Sol.**



$$\text{plane} = 2x - y + z = b$$

$$R \equiv \left( -1, 4, \frac{a+2}{2} \right) \rightarrow \text{on plane}$$

$$\therefore -2 - 4 + \frac{a+2}{2} = b$$

$$\Rightarrow a + 2 = 2b + 12 \Rightarrow a = 2b + 10 \quad \dots(i)$$

$$\langle PQ \rangle = \langle 4, -2, a-2 \rangle$$

$$\therefore \frac{2}{4} = \frac{-1}{-2} = \frac{1}{a-2}$$

$$\Rightarrow a - 2 = 2 \Rightarrow a = 4, b = -3$$

$$\therefore |a + b| = 1$$

7. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  satisfy the equation  $f(x+y) = f(x) \cdot f(y)$  for all  $x, y \in \mathbb{R}$  and  $f(x) \neq 0$  for any  $x \in \mathbb{R}$ . If the function  $f$  is differentiable at  $x = 0$  and  $f'(0) = 3$ , then

$$\lim_{h \rightarrow 0} \frac{1}{h} (f(h) - 1) \text{ is equal to } \underline{\hspace{2cm}}.$$

**Official Ans. by NTA (3)**

**Sol.** If  $f(x+y) = f(x) \cdot f(y)$  &  $f'(0) = 3$  then

$$f(x) = a^x \Rightarrow f'(x) = a^x \cdot \ln a$$

$$\Rightarrow f'(0) = \ln a = 3 \Rightarrow a = e^3$$

$$\Rightarrow f(x) = (e^3)^x = e^{3x}$$

$$\lim_{x \rightarrow 0} \frac{f(x) - 1}{x} = \lim_{x \rightarrow 0} \left( \frac{e^{3x} - 1}{3x} \times 3 \right) = 1 \times 3 = 3$$

8. Let  ${}^n C_r$  denote the binomial coefficient of  $x^r$  in the expansion of  $(1+x)^n$ .

$$\text{If } \sum_{k=0}^{10} (2^2 + 3k) {}^n C_k = \alpha \cdot 3^{10} + \beta \cdot 2^{10}, \alpha, \beta \in \mathbb{R},$$

then  $\alpha + \beta$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (19)**

**Allen Answer (Bonus)**

**Sol. BONUS**

Instead of  ${}^n C_k$  it must be  ${}^{10} C_k$  i.e.

$$\sum_{k=0}^{10} (2^2 + 3k) {}^{10} C_k = \alpha \cdot 3^{10} + \beta \cdot 2^{10}$$

$$\text{LHS} = 4 \sum_{k=0}^{10} {}^{10} C_k + 3 \sum_{k=0}^{10} k \cdot \frac{10}{k} \cdot {}^9 C_{k-1}$$

$$= 4 \cdot 2^{10} + 3 \cdot 10 \cdot 2^9$$

$$= 19 \cdot 2^{10} = \alpha \cdot 3^{10} + \beta \cdot 2^{10}$$

$$\Rightarrow \alpha = 0, \beta = 19 \Rightarrow \alpha + \beta = 19$$

9. Let P be a plane containing the line  $\frac{x-1}{3} = \frac{y+6}{4} = \frac{z+5}{2}$  and parallel to the line  $\frac{x-3}{4} = \frac{y-2}{-3} = \frac{z+5}{7}$ . If the point  $(1, -1, \alpha)$  lies on the plane P, then the value of  $|5\alpha|$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (38)**

**Sol.** Equation of plane is  $\begin{vmatrix} x-1 & y+6 & z+5 \\ 3 & 4 & 2 \\ 4 & -3 & 7 \end{vmatrix} = 0$

Now  $(1, -1, \alpha)$  lies on it so

$$\begin{vmatrix} 0 & 5 & \alpha+5 \\ 3 & 4 & 2 \\ 4 & -3 & 7 \end{vmatrix} = 0 \Rightarrow 5\alpha + 38 = 0 \Rightarrow |5\alpha| = 38$$

10. Let  $y = y(x)$  be the solution of the differential equation  $x dy - y dx = \sqrt{x^2 - y^2} dx$ ,  $x \geq 1$ , with  $y(1) = 0$ . If the area bounded by the line  $x = 1$ ,  $x = e^\pi$ ,  $y = 0$  and  $y = y(x)$  is  $\alpha e^{2\pi} + \beta$ , then the value of  $10(\alpha + \beta)$  is equal to \_\_\_\_\_.

**Official Ans. by NTA (4)**

**Sol.**  $x dy - y dx = \sqrt{x^2 - y^2} dx$   
 $\Rightarrow \frac{x dy - y dx}{x^2} = \frac{1}{x} \sqrt{1 - \frac{y^2}{x^2}} dx$   
 $\Rightarrow \int \frac{d\left(\frac{y}{x}\right)}{\sqrt{1 - \left(\frac{y}{x}\right)^2}} = \int \frac{dx}{x}$

$$\Rightarrow \sin^{-1}\left(\frac{y}{x}\right) = \ln|x| + c$$

at  $x = 1, y = 0 \Rightarrow c = 0$

$$y = x \sin(\ln x)$$

$$A = \int_1^{e^\pi} x \sin(\ln x) dx$$

$$x = e^t, dx = e^t dt \Rightarrow \int_0^\pi e^{2t} \sin(t) dt = A$$

$$\alpha e^{2\pi} + \beta = \left( \frac{e^{2t}}{5} (2 \sin t - \cos t) \right)_0^\pi = \frac{1 + e^{2\pi}}{5}$$

$$\alpha = \frac{1}{5}, \beta = \frac{1}{5} \text{ so } 10(\alpha + \beta) = 4$$