

**FINAL JEE(Advanced) EXAMINATION - 2019**  
(Held On Monday 27<sup>th</sup> MAY, 2019)

PAPER-1

TEST PAPER WITH ANSWER & SOLUTION

**PART-3 : MATHEMATICS**

**SECTION-1 : (Maximum Marks : 12)**

- This section contains **FOUR (04)** questions.
- Each question has **FOUR** options. **ONLY ONE** of these four options is the correct answer.
- For each question, choose the option corresponding to the correct answer.
- Answer to each question will be evaluated according to the following marking scheme :

*Full Marks* : +3 If **ONLY** the correct option is chosen.

*Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered)

*Negative Marks* : -1 In all other cases

1. Let  $M = \begin{bmatrix} \sin^4 \theta & -1 - \sin^2 \theta \\ 1 + \cos^2 \theta & \cos^4 \theta \end{bmatrix} = \alpha I + \beta M^{-1}$ ,

where  $\alpha = \alpha(\theta)$  and  $\beta = \beta(\theta)$  are real number, and I is the  $2 \times 2$  identity matrix. If

$\alpha^*$  is the minimum of the set  $\{\alpha(\theta) : \theta \in [0, 2\pi)\}$  and

$\beta^*$  is the minimum of the set  $\{\beta(\theta) : \theta \in [0, 2\pi)\}$ ,

then the value of  $\alpha^* + \beta^*$  is

- (1)  $-\frac{37}{16}$                       (2)  $-\frac{29}{16}$                       (3)  $-\frac{31}{16}$                       (4)  $-\frac{17}{16}$

**Ans. (2)**

**Sol.** Given  $M = \alpha I + \beta M^{-1}$

$$\Rightarrow M^2 - \alpha M - \beta I = O$$

By putting values of M and  $M^2$ , we get

$$\alpha(\theta) = 1 - 2\sin^2 \theta \cos^2 \theta = 1 - \frac{\sin^2 2\theta}{2} \geq \frac{1}{2}$$

$$\begin{aligned} \text{Also, } \beta(\theta) &= -(\sin^4 \theta \cos^4 \theta + (1 + \cos^2 \theta)(1 + \sin^2 \theta)) \\ &= -(\sin^4 \theta \cos^4 \theta + 1 + \cos^2 \theta + \sin^2 \theta + \sin^2 \theta \cos^2 \theta) \end{aligned}$$

$$= -(t^2 + t + 2), \quad t = \frac{\sin^2 2\theta}{4} \in \left[0, \frac{1}{4}\right]$$


$$\Rightarrow \beta(\theta) \geq -\frac{37}{16}$$

2. A line  $y = mx + 1$  intersects the circle  $(x - 3)^2 + (y + 2)^2 = 25$  at the points P and Q. If the midpoint of the line segment PQ has x-coordinate  $-\frac{3}{5}$ , then which one of the following options is correct ?

- (1)  $6 \leq m < 8$                       (2)  $2 \leq m < 4$                       (3)  $4 \leq m < 6$                       (4)  $-3 \leq m < -1$

Ans. (2)

Sol.



$$R \equiv \left( -\frac{3}{5}, \frac{-3m}{5} + 1 \right)$$

$$\text{So, } m \left( \frac{-\frac{3m}{5} + 3}{-\frac{3}{5} - 3} \right) = -1$$

$$\Rightarrow m^2 - 5m + 6 = 0$$

$$\Rightarrow m = 2, 3$$

3. Let S be the set of all complex numbers z satisfying  $|z - 2 + i| \geq \sqrt{5}$ . If the complex number  $z_0$  is such that

$\frac{1}{|z_0 - 1|}$  is the maximum of the set  $\left\{ \frac{1}{|z - 1|} : z \in S \right\}$ , then the principal argument of  $\frac{4 - z_0 - \bar{z}_0}{z_0 - \bar{z}_0 + 2i}$  is

- (1)  $\frac{\pi}{4}$                       (2)  $-\frac{\pi}{2}$                       (3)  $\frac{3\pi}{4}$                       (4)  $\frac{\pi}{2}$

Ans. (2)

Sol.


$$\arg \left( \frac{4 - (z_0 - \bar{z}_0)}{(z_0 - \bar{z}_0) + 2i} \right)$$

$$= \arg \left( \frac{4 - 2 \operatorname{Re} z_0}{2i \operatorname{Im} z_0 + 2i} \right) = \arg \left( \frac{2 - \operatorname{Re} z_0}{(1 + \operatorname{Im} z_0)i} \right)$$

$$= \arg \left( - \left( \frac{2 - \operatorname{Re} z_0}{1 + \operatorname{Im} z_0} \right) i \right)$$

$$= \arg(-ki) ; k > 0 \quad (\text{as } \operatorname{Re} z_0 < 2 \text{ \& } \operatorname{Im} z_0 > 0)$$


$$= -\frac{\pi}{2}$$



4. The area of the region  $\{(x, y) : xy \leq 8, 1 \leq y \leq x^2\}$  is

- (1)  $8\log_e 2 - \frac{14}{3}$       (2)  $16\log_e 2 - \frac{14}{3}$       (3)  $16\log_e 2 - 6$       (4)  $8\log_e 2 - \frac{7}{3}$

Ans. (2)



For intersection,  $\frac{8}{y} = \sqrt{y} \Rightarrow y = 4$

$$\begin{aligned} \text{Hence, required area} &= \int_1^4 \left( \frac{8}{y} - \sqrt{y} \right) dy \\ &= \left[ 8\ln y - \frac{2}{3}y^{3/2} \right]_1^4 = 16\ln 2 - \frac{14}{3} \end{aligned}$$

**Remark :** The question should contain the phrase "area of the bounded region in the first quadrant". Because, in the 2<sup>nd</sup> quadrant, the region above the line  $y = 1$  and below  $y = x^2$ , satisfies the region, which is unbounded.

### SECTION-2 : (Maximum Marks: 32)

- This section contains **EIGHT (08)** questions.
- Each question has **FOUR** options. **ONE OR MORE THAN ONE** of these four option(s) is (are) correct answer(s).
- For each question, choose the option(s) corresponding to (all ) the correct answer(s)
- Answer to each question will be evaluated according to the following marking scheme:
  - Full Marks* : +4 If only (all) the correct option(s) is (are) chosen.
  - Partial Marks* : +3 If all the four options are correct but **ONLY** three options are chosen.
  - Partial Marks* : +2 If three or more options are correct but **ONLY** two options are chosen and both of which are correct.
  - Partial Marks* : +1 If two or more options are correct but **ONLY** one option is chosen and it is a correct option.
  - Zero Marks* : 0 If none of the options is chosen (i.e. the question is unanswered).
  - Negative Marks* : -1 In all other cases.
- For example, in a question, if (A), (B) and (D) are the **ONLY** three options corresponding to correct answers, then
  - choosing **ONLY** (A), (B) and (D) will get +4 marks;
  - choosing **ONLY** (A) and (B) will get +2 marks;
  - choosing **ONLY** (A) and (D) will get +2 marks;
  - choosing **ONLY** (B) and (D) will get +2 marks;
  - choosing **ONLY** (A) will get +1 marks;
  - choosing **ONLY** (B) will get +1 marks;
  - choosing **ONLY** (D) will get +1 marks;
  - choosing no option (i.e. the question is unanswered) will get 0 marks, and
  - choosing any other combination of options will get -1 mark.

1. There are three bags  $B_1$ ,  $B_2$  and  $B_3$ . The bag  $B_1$  contains 5 red and 5 green balls,  $B_2$  contains 3 red and 5 green balls, and  $B_3$  contains 5 red and 3 green balls, Bags  $B_1$ ,  $B_2$  and  $B_3$  have probabilities  $\frac{3}{10}$ ,  $\frac{3}{10}$  and  $\frac{4}{10}$  respectively of being chosen. A bag is selected at random and a ball is chosen at random from the bag. Then which of the following options is/are correct ?

(1) Probability that the selected bag is  $B_3$  and the chosen ball is green equals  $\frac{3}{10}$

(2) Probability that the chosen ball is green equals  $\frac{39}{80}$

(3) Probability that the chosen ball is green, given that the selected bag is  $B_3$ , equals  $\frac{3}{8}$

(4) Probability that the selected bag is  $B_3$ , given that the chosen balls is green, equals  $\frac{5}{13}$

Ans. (2,3)

Sol.

Ball	Balls composition	$P(B_i)$
$B_1$	5R + 5G	$\frac{3}{10}$
$B_2$	3R + 5G	$\frac{3}{10}$
$B_3$	5R + 3G	$\frac{4}{10}$

$$(1) P(B_3 \cap G) = P\left(\frac{G}{B_3}\right)P(B_3)$$

$$= \frac{3}{8} \times \frac{4}{10} = \frac{3}{20}$$

$$(2) P(G) = P\left(\frac{G}{B_1}\right)P(B_1) + P\left(\frac{G}{B_2}\right)P(B_2) + P\left(\frac{G}{B_3}\right)P(B_3)$$

$$= \frac{3}{20} + \frac{3}{16} + \frac{3}{20} = \frac{39}{80}$$

$$(3) P\left(\frac{G}{B_3}\right) = \frac{3}{8}$$

$$(4) P\left(\frac{B_3}{G}\right) = \frac{P(G \cap B_3)}{P(G)} = \frac{3/20}{39/80} = \frac{4}{13}$$

2. Define the collections  $\{E_1, E_2, E_3, \dots\}$  of ellipses and  $\{R_1, R_2, R_3, \dots\}$  of rectangles as follows :

$$E_1 : \frac{x^2}{9} + \frac{y^2}{4} = 1;$$

$R_1$  : rectangle of largest area, with sides parallel to the axes, inscribed in  $E_1$  ;

$E_n$  : ellipse  $\frac{x^2}{a_n^2} + \frac{y^2}{b_n^2} = 1$  of largest area inscribed in  $R_{n-1}$ ,  $n > 1$  ;

$R_n$  : rectangle of largest area, with sides parallel to the axes, inscribed in  $E_n$ ,  $n > 1$ .

Then which of the following options is/are correct ?


(1) The eccentricities of  $E_{18}$  and  $E_{19}$  are NOT equal

(2) The distance of a focus from the centre in  $E_9$  is  $\frac{\sqrt{5}}{32}$

(3) The length of latus rectum of  $E_9$  is  $\frac{1}{6}$

(4)  $\sum_{n=1}^N (\text{area of } R_n) < 24$ , for each positive integer N

Ans. (3,4)



Area of  $R_1 = 3\sin 2\theta$  ; for this to be maximum

$$\Rightarrow \theta = \frac{\pi}{4} \Rightarrow \left( \frac{3}{\sqrt{2}}, \frac{2}{\sqrt{2}} \right)$$

Hence for subsequent areas of rectangles  $R_n$  to be maximum the coordinates will be in GP with common

$$\text{ratio } r = \frac{1}{\sqrt{2}} \Rightarrow a_n = \frac{3}{(\sqrt{2})^{n-1}} ; b_n = \frac{2}{(\sqrt{2})^{n-1}}$$

Eccentricity of all the ellipses will be same

$$\text{Distance of a focus from the centre in } E_9 = a_9 e_9 = \sqrt{a_9^2 - b_9^2} = \frac{\sqrt{5}}{16}$$

$$\text{Length of latus rectum of } E_9 = \frac{2b_9^2}{a_9} = \frac{1}{6}$$

$$\therefore \sum_{n=1}^{\infty} \text{Area of } R_n = 12 + \frac{12}{2} + \frac{12}{4} + \dots = 24$$

$$\Rightarrow \sum_{n=1}^N (\text{area of } R_n) < 24, \text{ for each positive integer N}$$

3. Let  $M = \begin{bmatrix} 0 & 1 & a \\ 1 & 2 & 3 \\ 3 & b & 1 \end{bmatrix}$  and  $\text{adj}M = \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix}$  where a and b are real numbers. Which of the following

options is/are correct ?

(1)  $a + b = 3$

(2)  $\det(\text{adj}M^2) = 81$

(3)  $(\text{adj}M)^{-1} + \text{adj}M^{-1} = -M$

(4) If  $M \begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ , then  $\alpha - \beta + \gamma = 3$

**Ans. (1,3,4)**

**Sol.**  $(\text{adj}M)_{11} = 2 - 3b = -1 \Rightarrow b = 1$

Also,  $(\text{adj}M)_{22} = -3a = -6 \Rightarrow a = 2$

$$\text{Now, } \det M = \begin{vmatrix} 0 & 1 & 2 \\ 1 & 2 & 3 \\ 3 & 1 & 1 \end{vmatrix} = -2$$

$$\Rightarrow \det(\text{adj}M^2) = (\det M^2)^2 \\ = (\det M)^4 = 16$$

$$\text{Also } M^{-1} = \frac{\text{adj}M}{\det M}$$

$$\Rightarrow \text{adj}M = -2M^{-1}$$

$$\Rightarrow (\text{adj}M)^{-1} = \frac{-1}{2}M$$

$$\text{And, } \text{adj}(M^{-1}) = (M^{-1})^{-1} \det(M^{-1})$$

$$= \frac{1}{\det M} M = \frac{-M}{2}$$

$$\text{Hence, } (\text{adj}M)^{-1} + \text{adj}(M^{-1}) = -M$$

$$\text{Further, } MX = b$$

$$\Rightarrow X = M^{-1}b = \frac{-\text{adj}M}{2}b$$

$$= \frac{-1}{2} \begin{bmatrix} -1 & 1 & -1 \\ 8 & -6 & 2 \\ -5 & 3 & -1 \end{bmatrix} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$= \frac{-1}{2} \begin{bmatrix} -2 \\ 2 \\ -2 \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}$$

$$\Rightarrow (\alpha, \beta, \gamma) = (1, -1, 1)$$

4. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be given by

$$f(x) = \begin{cases} x^5 + 5x^4 + 10x^3 + 10x^2 + 3x + 1, & x < 0; \\ x^2 - x + 1, & 0 \leq x < 1; \\ \frac{2}{3}x^3 - 4x^2 + 7x - \frac{8}{3}, & 1 \leq x < 3; \\ (x-2)\log_e(x-2) - x + \frac{10}{3}, & x \geq 3 \end{cases}$$

Then which of the following options is/are correct ?

- (1)  $f'$  has a local maximum at  $x = 1$
- (2)  $f$  is onto
- (3)  $f$  is increasing on  $(-\infty, 0)$
- (4)  $f'$  is NOT differentiable at  $x = 1$

Ans. (1,2,4)

Sol.  $f(x) = \begin{cases} (x+1)^5 - 2x, & x < 0; \\ x^2 - x + 1, & 0 \leq x < 1; \\ \frac{2}{3}x^3 - 4x^2 + 7x - \frac{8}{3}, & 1 \leq x < 3; \\ (x-2)\log_e(x-2) - x + \frac{10}{3}, & x \geq 3 \end{cases}$

for  $x < 0$ ,  $f(x)$  is continuous

&  $\lim_{x \rightarrow -\infty} f(x) = -\infty$  and  $\lim_{x \rightarrow 0^-} f(x) = 1$

Hence,  $(-\infty, 1) \subset \text{Range of } f(x) \text{ in } (-\infty, 0)$

$f'(x) = 5(x+1)^4 - 2$ , which changes sign in  $(-\infty, 0)$


$\Rightarrow f(x)$  is non-monotonic in  $(-\infty, 0)$

For  $x \geq 3$ ,  $f(x)$  is again continuous and  $\lim_{x \rightarrow \infty} f(x) = \infty$  and  $f(3) = \frac{1}{3}$

$\Rightarrow \left[\frac{1}{3}, \infty\right) \subset \text{Range of } f(x) \text{ in } [3, \infty)$

Hence, range of  $f(x)$  is  $\mathbb{R}$

$f'(x) = \begin{cases} 2x-1, & 0 \leq x < 1 \\ 2x^2 - 8x + 7, & 1 \leq x < 3 \end{cases}$



Hence  $f'$  has a local maximum at  $x = 1$  and  $f'$  is NOT differentiable at  $x = 1$ .

5. Let  $\alpha$  and  $\beta$  be the roots of  $x^2 - x - 1 = 0$ , with  $\alpha > \beta$ . For all positive integers  $n$ , define

$$a_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}, \quad n \geq 1$$

$$b_1 = 1 \text{ and } b_n = a_{n-1} + a_{n+1}, \quad n \geq 2.$$

Then which of the following options is/are correct ?

(1)  $a_1 + a_2 + a_3 + \dots + a_n = a_{n+2} - 1$  for all  $n \geq 1$

(2)  $\sum_{n=1}^{\infty} \frac{a_n}{10^n} = \frac{10}{89}$

(3)  $\sum_{n=1}^{\infty} \frac{b_n}{10^n} = \frac{8}{89}$

(4)  $b_n = \alpha^n + \beta^n$  for all  $n \geq 1$

**Ans. (1,2,4)**

**Sol.**  $\alpha, \beta$  are roots of  $x^2 - x - 1$

$$\begin{aligned} a_{r+2} - a_r &= \frac{(\alpha^{r+2} - \beta^{r+2}) - (\alpha^r - \beta^r)}{\alpha - \beta} = \frac{(\alpha^{r+2} - \alpha^r) - (\beta^{r+2} - \beta^r)}{\alpha - \beta} \\ &= \frac{\alpha^r(\alpha^2 - 1) - \beta^r(\beta^2 - 1)}{\alpha - \beta} = \frac{\alpha^r\alpha - \beta^r\beta}{\alpha - \beta} = \frac{\alpha^{r+1} - \beta^{r+1}}{\alpha - \beta} = a_{r+1} \end{aligned}$$

$$\Rightarrow a_{r+2} - a_{r+1} = a_r$$

$$\Rightarrow \sum_{r=1}^n a_r = a_{n+2} - a_2 = a_{n+2} - \frac{\alpha^2 - \beta^2}{\alpha - \beta} = a_{n+2} - (\alpha + \beta) = a_{n+2} - 1$$

$$\text{Now } \sum_{n=1}^{\infty} \frac{a_n}{10^n} = \frac{\sum_{n=1}^{\infty} \left(\frac{\alpha}{10}\right)^n - \sum_{n=1}^{\infty} \left(\frac{\beta}{10}\right)^n}{\alpha - \beta}$$

$$\begin{aligned} &= \frac{\frac{\frac{\alpha}{10}}{1 - \frac{\alpha}{10}} - \frac{\frac{\beta}{10}}{1 - \frac{\beta}{10}}}{\alpha - \beta} = \frac{\frac{\alpha}{10 - \alpha} - \frac{\beta}{10 - \beta}}{(\alpha - \beta)} = \frac{10}{(10 - \alpha)(10 - \beta)} = \frac{10}{89} \end{aligned}$$

$$\sum_{n=1}^{\infty} \frac{b_n}{10^n} = \sum_{n=1}^{\infty} \frac{a_{n-1} + a_{n+1}}{10^n} = \frac{\frac{\alpha}{10}}{1 - \frac{\alpha}{10}} + \frac{\frac{\beta}{10}}{1 - \frac{\beta}{10}} = \frac{12}{89}$$

Further,  $b_n = a_{n-1} + a_{n+1}$

$$= \frac{(\alpha^{n-1} - \beta^{n-1}) + (\alpha^{n+1} - \beta^{n+1})}{\alpha - \beta}$$

(as  $\alpha\beta = -1 \Rightarrow \alpha^{n-1} = -\alpha^n\beta$  &  $\beta^{n-1} = -\alpha\beta^n$ )

$$= \frac{\alpha^n(\alpha - \beta) + (\alpha - \beta)\beta^n}{\alpha - \beta} = \alpha^n + \beta^n$$



6. Let  $\Gamma$  denote a curve  $y = y(x)$  which is in the first quadrant and let the point  $(1, 0)$  lie on it. Let the tangent to  $\Gamma$  at a point  $P$  intersect the  $y$ -axis at  $Y_p$ . If  $PY_p$  has length 1 for each point  $P$  on  $\Gamma$ , then which of the following options is/are correct ?

(1)  $y = \log_e \left( \frac{1 + \sqrt{1 - x^2}}{x} \right) - \sqrt{1 - x^2}$


(2)  $xy' - \sqrt{1 - x^2} = 0$

(3)  $y = -\log_e \left( \frac{1 + \sqrt{1 - x^2}}{x} \right) + \sqrt{1 - x^2}$

(4)  $xy' + \sqrt{1 - x^2} = 0$

Ans. (1,4)

Sol.



$$Y - y = y'(X - x)$$

So,  $Y_p = (0, y - xy')$

So,  $x^2 + (xy')^2 = 1 \Rightarrow \frac{dy}{dx} = -\sqrt{\frac{1 - x^2}{x^2}}$

[  $\frac{dy}{dx}$  can not be positive i.e.  $f(x)$  can not be increasing in first quadrant, for  $x \in (0, 1)$  ]

Hence,  $\int dy = -\int \frac{\sqrt{1 - x^2}}{x} dx$

$\Rightarrow y = -\int \frac{\cos^2 \theta d\theta}{\sin \theta}$  ; put  $x = \sin \theta$

$\Rightarrow y = -\int \operatorname{cosec} \theta d\theta + \int \sin \theta d\theta$

$\Rightarrow y = \ln(\operatorname{cosec} \theta + \cot \theta) - \cos \theta + C$

$\Rightarrow y = \ln \left( \frac{1 + \sqrt{1 - x^2}}{x} \right) - \sqrt{1 - x^2} + C$

$\Rightarrow y = \ln \left( \frac{1 + \sqrt{1 - x^2}}{x} \right) - \sqrt{1 - x^2}$  (as  $y(1) = 0$ )

7. In a non-right-angled triangle  $\Delta PQR$ , let  $p, q, r$  denote the lengths of the sides opposite to the angles at  $P, Q, R$  respectively. The median from  $R$  meets the side  $PQ$  at  $S$ , the perpendicular from  $P$  meets the side  $QR$  at  $E$ , and  $RS$  and  $PE$  intersect at  $O$ . If  $p = \sqrt{3}, q = 1$ , and the radius of the circumcircle of the  $\Delta PQR$  equals 1, then which of the following options is/are correct ?

(1) Area of  $\Delta SOE = \frac{\sqrt{3}}{12}$

(2) Radius of incircle of  $\Delta PQR = \frac{\sqrt{3}}{2}(2 - \sqrt{3})$

(3) Length of  $RS = \frac{\sqrt{7}}{2}$

(4) Length of  $OE = \frac{1}{6}$

Ans. (2,3,4)

Sol.  $\frac{\sin P}{\sqrt{3}} = \frac{\sin Q}{1} = \frac{1}{2R} = \frac{1}{2}$

$\Rightarrow P = \frac{\pi}{3}$  or  $\frac{2\pi}{3}$  and  $Q = \frac{\pi}{6}$  or  $\frac{5\pi}{6}$

Since  $p > q \Rightarrow P > Q$

So, if  $P = \frac{\pi}{3}$  and  $Q = \frac{\pi}{6} \Rightarrow R = \frac{\pi}{2}$  (not possible)

Hence,  $P = \frac{2\pi}{3}$  and  $Q = R = \frac{\pi}{6}$


$$r = \frac{\Delta}{s} = \frac{\frac{1}{2}(1)(\sqrt{3})\left(\frac{1}{2}\right)}{\left(\frac{\sqrt{3}+2}{2}\right)} = \frac{\sqrt{3}}{2}(2 - \sqrt{3})$$

Now, area of  $\Delta SEF = \frac{1}{4}$  area of  $\Delta PQR$

$\Rightarrow$  area of  $\Delta SOE = \frac{1}{3}$  area of  $\Delta SEF = \frac{1}{12}$  area of  $\Delta PQR = \frac{1}{12} \cdot \frac{\sqrt{3}}{4} = \frac{\sqrt{3}}{48}$

$RS = \frac{1}{2}\sqrt{2(3) + 2(1) - 1} = \frac{\sqrt{7}}{2}$

$OE = \frac{1}{3}PE = \frac{1}{3} \cdot \frac{1}{2}\sqrt{2(1)^2 + 2(1)^2 - 3} = \frac{1}{6}$



8. Let  $L_1$  and  $L_2$  denotes the lines

$$\vec{r} = \hat{i} + \lambda(-\hat{i} + 2\hat{j} + 2\hat{k}), \lambda \in \mathbb{R} \text{ and}$$

$$\vec{r} = \mu(2\hat{i} - \hat{j} + 2\hat{k}), \mu \in \mathbb{R}$$

respectively. If  $L_3$  is a line which is perpendicular to both  $L_1$  and  $L_2$  and cuts both of them, then which of the following options describe(s)  $L_3$  ?

(1)  $\vec{r} = \frac{1}{3}(2\hat{i} + \hat{k}) + t(2\hat{i} + 2\hat{j} - \hat{k}), t \in \mathbb{R}$

(2)  $\vec{r} = \frac{2}{9}(2\hat{i} - \hat{j} + 2\hat{k}) + t(2\hat{i} + 2\hat{j} - \hat{k}), t \in \mathbb{R}$

(3)  $\vec{r} = t(2\hat{i} + 2\hat{j} - \hat{k}), t \in \mathbb{R}$

(4)  $\vec{r} = \frac{2}{9}(4\hat{i} + \hat{j} + \hat{k}) + t(2\hat{i} + 2\hat{j} - \hat{k}), t \in \mathbb{R}$

**Ans. (1,2,4)**

**Sol.** Points on  $L_1$  and  $L_2$  are respectively  $A(1 - \lambda, 2\lambda, 2\lambda)$  and  $B(2\mu, -\mu, 2\mu)$

So,  $\overline{AB} = (2\mu + \lambda - 1)\hat{i} + (-\mu - 2\lambda)\hat{j} + (2\mu - 2\lambda)\hat{k}$

and vector along their shortest distance  $= 2\hat{i} + 2\hat{j} - \hat{k}$ .

Hence,  $\frac{2\mu + \lambda - 1}{2} = \frac{-\mu - 2\lambda}{2} = \frac{2\mu - 2\lambda}{-1}$

$\Rightarrow \lambda = \frac{1}{9} \text{ \& } \mu = \frac{2}{9}$

Hence,  $A \equiv \left(\frac{8}{9}, \frac{2}{9}, \frac{2}{9}\right)$  and  $B \equiv \left(\frac{4}{9}, -\frac{2}{9}, \frac{4}{9}\right)$

$\Rightarrow$  Mid point of AB  $\equiv \left(\frac{2}{3}, 0, \frac{1}{3}\right)$

### SECTION-3 : (Maximum Marks: 18)

- This section contains **SIX (06)** questions. The answer to each question is a **NUMERICAL VALUE**.
- For each question, enter the correct numerical value of the answer using the mouse and the on-screen virtual numeric keypad in the place designated to enter the answer. If the numerical value has more than two decimal places, **truncate/round-off** the value to **Two** decimal places.
- Answer to each question will be evaluated according to the following marking scheme:

*Full Marks* : +3 If **ONLY** the correct numerical value is entered.

*Zero Marks* : 0 In all other cases.

1. If

$$I = \frac{2}{\pi} \int_{-\pi/4}^{\pi/4} \frac{dx}{(1 + e^{\sin x})(2 - \cos 2x)}$$

then  $27I^2$  equals \_\_\_\_\_

**Ans. (4.00)**

**Sol.**  $2I = \frac{2}{\pi} \int_{-\pi/4}^{\pi/4} \left[ \frac{1}{(1 + e^{\sin x})(2 - \cos 2x)} + \frac{1}{(1 + e^{-\sin x})(2 - \cos 2x)} \right] dx$  (using King's Rule)

$$\Rightarrow I = \frac{1}{\pi} \int_{-\pi/4}^{\pi/4} \frac{dx}{2 - \cos 2x}$$

$$\Rightarrow I = \frac{2}{\pi} \int_0^{\pi/4} \frac{dx}{2 - \cos 2x} = \frac{2}{\pi} \int_0^{\pi/4} \frac{\sec^2 x dx}{1 + 3 \tan^2 x}$$

$$= \frac{2}{\sqrt{3}\pi} \left[ \tan^{-1}(\sqrt{3} \tan x) \right]_0^{\pi/4} = \frac{2}{3\sqrt{3}}$$

$$\Rightarrow 27I^2 = 27 \times \frac{4}{27} = 4$$

2. Let the point B be the reflection of the point A(2, 3) with respect to the line  $8x - 6y - 23 = 0$ . Let  $\Gamma_A$  and  $\Gamma_B$  be circles of radii 2 and 1 with centres A and B respectively. Let T be a common tangent to the circles  $\Gamma_A$  and  $\Gamma_B$  such that both the circles are on the same side of T. If C is the point of intersection of T and the line passing through A and B, then the length of the line segment AC is \_\_\_\_\_


**Ans. (10.00)**

**Sol.** Distance of point A from given line =  $\frac{5}{2}$

$$\frac{CA}{CB} = \frac{2}{1}$$

$$\Rightarrow \frac{AC}{AB} = \frac{2}{1}$$

$$\Rightarrow AC = 2 \times 5 = 10$$



3. Let  $AP(a; d)$  denote the set of all the terms of an infinite arithmetic progression with first term  $a$  and common difference  $d > 0$ . If  $AP(1; 3) \cap AP(2; 5) \cap AP(3; 7) = AP(a; d)$  then  $a + d$  equals \_\_\_\_

**Ans. (157.00)**

**Sol.** We equate the general terms of three respective

$$A.P.'s \text{ as } 1 + 3a = 2 + 5b = 3 + 7c$$

$$\Rightarrow 3 \text{ divides } 1 + 2b \text{ and } 5 \text{ divides } 1 + 2c$$

$$\Rightarrow 1 + 2c = 5, 15, 25 \text{ etc.}$$

So, first such terms are possible when  $1 + 2c = 15$  i.e.  $c = 7$

$$\text{Hence, first term} = a = 52$$

$$d = \text{lcm}(3, 5, 7) = 105$$

$$\Rightarrow a + d = 157$$

4. Let  $S$  be the sample space of all  $3 \times 3$  matrices with entries from the set  $\{0, 1\}$ . Let the events  $E_1$  and  $E_2$  be given by

$$E_1 = \{A \in S : \det A = 0\} \text{ and}$$

$$E_2 = \{A \in S : \text{sum of entries of } A \text{ is } 7\}.$$

If a matrix is chosen at random from  $S$ , then the conditional probability  $P(E_1|E_2)$  equals \_\_\_\_

**Ans. (0.50)**

**Sol.**  $n(E_2) = {}^9C_2$  (as exactly two cyphers are there)

Now,  $\det A = 0$ , when two cyphers are in the same column or same row

$$\Rightarrow n(E_1 \cap E_2) = 6 \times {}^3C_2.$$

$$\text{Hence, } P\left(\frac{E_1}{E_2}\right) = \frac{n(E_1 \cap E_2)}{n(E_2)} = \frac{18}{36} = \frac{1}{2} = 0.50$$

5. Three lines are given by

$$\vec{r} = \lambda \hat{i}, \lambda \in \mathbb{R}$$

$$\vec{r} = \mu(\hat{i} + \hat{j}), \mu \in \mathbb{R} \text{ and}$$

$$\vec{r} = \nu(\hat{i} + \hat{j} + \hat{k}), \nu \in \mathbb{R}.$$

Let the lines cut the plane  $x + y + z = 1$  at the points  $A, B$  and  $C$  respectively. If the area of the triangle  $ABC$  is  $\Delta$  then the value of  $(6\Delta)^2$  equals \_\_\_\_

**Ans. (0.75)**

**Sol.**  $A(1, 0, 0)$ ,  $B\left(\frac{1}{2}, \frac{1}{2}, 0\right)$  &  $C\left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\right)$

Hence,  $\overline{AB} = -\frac{1}{2}\hat{i} + \frac{1}{2}\hat{j}$  &  $\overline{AC} = -\frac{2}{3}\hat{i} + \frac{1}{3}\hat{j} + \frac{1}{3}\hat{k}$

So,  $\Delta = \frac{1}{2}|\overline{AB} \times \overline{AC}| = \frac{1}{2}\sqrt{\frac{1}{2} \times \frac{2}{3} - \frac{1}{4}}$

$$= \frac{1}{2 \times 2\sqrt{3}}$$

$$\Rightarrow (6\Delta)^2 = \frac{3}{4} = 0.75$$

6. Let  $\omega \neq 1$  be a cube root of unity. Then the minimum of the set  $\{|a + b\omega + c\omega^2|^2 : a, b, c \text{ distinct non-zero integers}\}$  equals \_\_\_\_\_

**Ans. (3.00)**

**Sol.**  $|a + b\omega + c\omega^2|^2 = (a + b\omega + c\omega^2)(\overline{a + b\omega + c\omega^2})$

$$= (a + b\omega + c\omega^2)(a + b\omega^2 + c\omega)$$

$$= a^2 + b^2 + c^2 - ab - bc - ca$$

$$= \frac{1}{2}[(a-b)^2 + (b-c)^2 + (c-a)^2]$$

$$\geq \frac{1+1+4}{2} = 3 \quad (\text{when } a = 1, b = 2, c = 3)$$