


FINAL JEE–MAIN EXAMINATION – APRIL, 2019
Held On Monday 08th APRIL, 2019
TIME: 09 : 30 AM To 12 : 30 PM

1. The shortest distance between the line $y = x$ and the curve $y^2 = x - 2$ is :

(1) $\frac{7}{4\sqrt{2}}$ (2) $\frac{7}{8}$
 (3) $\frac{11}{4\sqrt{2}}$ (4) 2

Official Ans. by NTA (1)



$$\text{we have, } 2y \cdot \frac{dy}{dx} = 1 \Rightarrow \frac{dy}{dx} \Big|_{P(2+t^2, t)} = \frac{1}{2t} = 1$$

$$\Rightarrow t = \frac{1}{2}$$

$$\therefore P\left(\frac{9}{4}, \frac{1}{2}\right)$$

So, shortest distance

$$= \frac{\left| \frac{9}{4} - \frac{1}{2} \right|}{\sqrt{2}} = \frac{7}{4\sqrt{2}}$$

2. Let $y = y(x)$ be the solution of the differential equation, $(x^2 + 1)^2 \frac{dy}{dx} + 2x(x^2 + 1)y = 1$ such

that $y(0) = 0$. If $\sqrt{a}y(1) = \frac{\pi}{32}$, then the value of 'a' is :

(1) $\frac{1}{2}$ (2) $\frac{1}{16}$
 (3) $\frac{1}{4}$ (4) 1

Official Ans. by NTA (2)

Sol. $\frac{dy}{dx} + \left(\frac{2x}{x^2 + 1} \right) y = \frac{1}{(x^2 + 1)^2}$

(Linear differential equation)

$$\therefore \text{I.F.} = e^{\int \frac{2x}{x^2 + 1} dx} = (x^2 + 1)$$

So, general solution is $y(x^2 + 1) = \tan^{-1}x + c$

$$\text{As } y(0) = 0 \Rightarrow c = 0$$

$$\therefore y(x) = \frac{\tan^{-1}x}{x^2 + 1}$$


$$\text{As, } \sqrt{a} \cdot y(1) = \frac{\pi}{32}$$

$$\Rightarrow \sqrt{a} = \frac{1}{4} \Rightarrow a = \frac{1}{16}$$

3. A point on the straight line, $3x + 5y = 15$ which is equidistant from the coordinate axes will lie only in :

(1) 1st and 2nd quadrants
 (2) 4th quadrant
 (3) 1st, 2nd and 4th quadrant
 (4) 1st quadrant

Official Ans. by NTA (1)



$$\text{Now, } \left| \frac{15-3t}{5} \right| = |t|$$

$$\Rightarrow \frac{15-3t}{5} = t \text{ or } \frac{15-3t}{5} = -t$$

$$\therefore t = \frac{15}{8} \text{ or } t = -\frac{15}{2}$$

So, $P\left(\frac{15}{8}, \frac{15}{8}\right) \in \text{I}^{\text{st}}$ quadrant

or $P\left(\frac{-15}{2}, \frac{15}{2}\right) \in \text{II}^{\text{nd}}$ quadrant

4. If α and β be the roots of the equation $x^2 - 2x + 2 = 0$, then the least value of n for which

$$\left(\frac{\alpha}{\beta}\right)^n = 1 \text{ is :}$$

(1) 2 (2) 3
(3) 4 (4) 5

Official Ans. by NTA (3)

Sol. $(x - 1)^2 + 1 = 0 \Rightarrow x = 1 + i, 1 - i$

$$\therefore \left(\frac{\alpha}{\beta}\right)^n = 1 \Rightarrow (\pm i)^n = 1$$

$\therefore n$ (least natural number) = 4

5. $\lim_{x \rightarrow 0} \frac{\sin^2 x}{\sqrt{2} - \sqrt{1 + \cos x}}$ equals :

(1) $2\sqrt{2}$ (2) $4\sqrt{2}$
(3) $\sqrt{2}$ (4) 4

Official Ans. by NTA (2)

$$\text{Sol. } \lim_{x \rightarrow 0} \frac{\left(\frac{\sin^2 x}{x^2}\right) \left(\sqrt{2} + \sqrt{1 + \cos x}\right)}{\left(\frac{1 - \cos x}{x^2}\right)}$$

$$= \frac{(1)^2 \cdot (2\sqrt{2})}{\frac{1}{2}} = 4\sqrt{2}$$


6. The length of the perpendicular from the point

$(2, -1, 4)$ on the straight line, $\frac{x+3}{10} = \frac{y-2}{-7} = \frac{z}{1}$ is :

(1) less than 2
(2) greater than 3 but less than 4
(3) greater than 4
(4) greater than 2 but less than 3

Official Ans. by NTA (2)

Sol.



$$\text{Now, } \overrightarrow{MP} \cdot (10\hat{i} - 7\hat{j} + \hat{k}) = 0$$

$$\Rightarrow \lambda = \frac{1}{2}$$

\therefore Length of perpendicular

$$(\text{PM}) = \sqrt{0 + \frac{1}{4} + \frac{49}{4}}$$

$$= \sqrt{\frac{50}{4}} = \sqrt{\frac{25}{2}} = \frac{5}{\sqrt{2}},$$

which is greater than 3 but less than 4.

7. The magnitude of the projection of the vector $2\hat{i} + 3\hat{j} + \hat{k}$ on the vector perpendicular to the plane containing the vectors $\hat{i} + \hat{j} + \hat{k}$ and $\hat{i} + 2\hat{j} + 3\hat{k}$, is :

$$(1) \frac{\sqrt{3}}{2} \quad (2) \sqrt{\frac{3}{2}}$$

$$(3) \sqrt{6} \quad (4) 3\sqrt{6}$$

Official Ans. by NTA (2)

Sol. Vector perpendicular to plane containing the vectors $\hat{i} + \hat{j} + \hat{k}$ & $\hat{i} + 2\hat{j} + 3\hat{k}$ is parallel to vector

$$= \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 1 \\ 1 & 2 & 3 \end{vmatrix} = \hat{i} - 2\hat{j} + \hat{k}$$

\therefore Required magnitude of projection

$$= \frac{|(2\hat{i} + 3\hat{j} + \hat{k}) \cdot (\hat{i} - 2\hat{j} + \hat{k})|}{|\hat{i} - 2\hat{j} + \hat{k}|}$$

$$= \frac{|2 - 6 + 1|}{|\sqrt{6}|} = \frac{3}{\sqrt{6}} = \sqrt{\frac{3}{2}}$$

8. The contrapositive of the statement "If you are born in India, then you are a citizen of India", is :

- If you are born in India, then you are not a citizen of India.
- If you are not a citizen of India, then you are not born in India.
- If you are a citizen of India, then you are born in India.
- If you are not born in India, then you are not a citizen of India.

Official Ans. by NTA (2)

Sol. The contrapositive of statement

$$p \rightarrow q \text{ is } \sim q \rightarrow \sim p$$

Here, p : you are born in India.

q : you are citizen of India.

So, contrapositive of above statement is

"If you are not a citizen of India, then you are not born in India".

9. The mean and variance of seven observations are 8 and 16, respectively. If 5 of the observations are 2, 4, 10, 12, 14, then the product of the remaining two observations is :

- 40
- 49
- 48
- 45

Official Ans. by NTA (3)

Sol. Let 7 observations be $x_1, x_2, x_3, x_4, x_5, x_6, x_7$

$$\bar{x} = 8 \Rightarrow \sum_{i=1}^7 x_i = 56 \quad \dots\dots(1)$$

$$\text{Also } \sigma^2 = 16$$

$$\Rightarrow 16 = \frac{1}{7} \left(\sum_{i=1}^7 x_i^2 \right) - (\bar{x})^2$$

$$\Rightarrow 16 = \frac{1}{7} \left(\sum_{i=1}^7 x_i^2 \right) - 64$$

$$\Rightarrow \left(\sum_{i=1}^7 x_i^2 \right) = 560 \quad \dots\dots(2)$$

$$\text{Now, } x_1 = 2, x_2 = 4, x_3 = 10, x_4 = 12, x_5 = 14$$

$$\Rightarrow x_6 + x_7 = 14 \quad (\text{from (1)})$$

$$\text{& } x_6^2 + x_7^2 = 100 \quad (\text{from (2)})$$

$$\therefore x_6^2 + x_7^2 = (x_6 + x_7)^2 - 2x_6 \cdot x_7 \Rightarrow x_6 \cdot x_7 = 48$$

10. If $f(x) = \frac{2-x \cos x}{2+x \cos x}$ and $g(x) = \log_e x, (x > 0)$ then

the value of integral $\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} g(f(x)) dx$ is :

- $\log_e 3$
- $\log_e 2$
- $\log_e e$
- $\log_e 1$

Official Ans. by NTA (4)

$$\text{Sol. } g(f(x)) = \ln(f(x)) = \ln\left(\frac{2-x \cos x}{2+x \cos x}\right)$$

$$\therefore I = \int_{-\pi/4}^{\pi/4} \ln\left(\frac{2-x \cos x}{2+x \cos x}\right) dx$$

$$= \int_0^{\pi/4} \left(\ln\left(\frac{2-x \cos x}{2+x \cos x}\right) + \ln\left(\frac{2+x \cos x}{2-x \cos x}\right) \right) dx$$

$$= \int_0^{\pi/2} (0) dx = 0 = \log_e(1)$$

11. If the tangents on the ellipse $4x^2 + y^2 = 8$ at the points $(1, 2)$ and (a, b) are perpendicular to each other, then a^2 is equal to :

$$(1) \frac{64}{17} \quad (2) \frac{2}{17}$$

$$(3) \frac{128}{17} \quad (4) \frac{4}{17}$$

Official Ans. by NTA (2)

$$\text{Sol. } 4a^2 + b^2 = 8 \quad \dots\dots(1)$$

$$\text{also } \left. \frac{dy}{dx} \right|_{(1,2)} = -\frac{4x}{y} = -2$$

$$\Rightarrow -\frac{4a}{b} = \frac{1}{2}$$

$$b = -8a$$

$$\Rightarrow b^2 = 64a^2$$

$$64a^2 = 8$$

$$a^2 = \frac{2}{17}$$

12. If $\alpha = \cos^{-1}\left(\frac{3}{5}\right)$, $\beta = \tan^{-1}\left(\frac{1}{3}\right)$,

where $0 < \alpha, \beta < \frac{\pi}{2}$, then $\alpha - \beta$ is equal to :

(1) $\sin^{-1}\left(\frac{9}{5\sqrt{10}}\right)$ (2) $\tan^{-1}\left(\frac{9}{14}\right)$
(3) $\cos^{-1}\left(\frac{9}{5\sqrt{10}}\right)$ (4) $\tan^{-1}\left(\frac{9}{5\sqrt{10}}\right)$

Official Ans. by NTA (1)

Sol. $\cos \alpha = \frac{3}{5}, \tan \beta = \frac{1}{3}$

$$\Rightarrow \tan \alpha = \frac{4}{3}$$

$$\Rightarrow \tan(\alpha - \beta) = \frac{\frac{4}{3} - \frac{1}{3}}{1 + \frac{4}{3} \cdot \frac{1}{3}} = \frac{9}{13}$$

$$\Rightarrow \sin(\alpha - \beta) = \frac{9}{5\sqrt{10}}$$

$$\Rightarrow \alpha - \beta = \sin^{-1}\left(\frac{9}{5\sqrt{10}}\right)$$


13. If S_1 and S_2 are respectively the sets of local minimum and local maximum points of the function, $f(x) = 9x^4 + 12x^3 - 36x^2 + 25$, $x \in \mathbb{R}$, then :

(1) $S_1 = \{-2, 1\}$; $S_2 = \{0\}$
(2) $S_1 = \{-2, 0\}$; $S_2 = \{1\}$
(3) $S_1 = \{-2\}$; $S_2 = \{0, 1\}$
(4) $S_1 = \{-1\}$; $S_2 = \{0, 2\}$

Official Ans. by NTA (1)

Sol. $f(x) = 9x^4 + 12x^3 - 36x^2 + 25$

$$f'(x) = 36x^3 + 36x^2 - 72x \\ = 36x(x^2 + x - 2) \\ = 36x(x - 1)(x + 2)$$



Points of minima = $\{-2, 1\} = S_1$


Point of maxima = $\{0\} = S_2$

14. Let $O(0, 0)$ and $A(0, 1)$ be two fixed points. Then the locus of a point P such that the perimeter of ΔAOP is 4, is :

(1) $8x^2 - 9y^2 + 9y = 18$
(2) $9x^2 + 8y^2 - 8y = 16$
(3) $8x^2 + 9y^2 - 9y = 18$
(4) $9x^2 - 8y^2 + 8y = 16$

Official Ans. by NTA (2)

Sol.



$$AP + OP + AO = 4$$

$$\sqrt{h^2 + (k-1)^2} + \sqrt{h^2 + k^2} + 1 = 4$$

$$\sqrt{h^2 + (k-1)^2} + \sqrt{h^2 + k^2} = 3$$

$$h^2 + (k-1)^2 = 9 + h^2 + k^2 - 6\sqrt{h^2 + k^2}$$

$$-2k - 8 = -6\sqrt{h^2 + k^2}$$

$$k + 4 = 3\sqrt{h^2 + k^2}$$

$$k^2 + 16 + 8k = 9(h^2 + k^2)$$

$$9h^2 + 8k^2 - 8k - 16 = 0$$

$$\text{Locus of } P \text{ is } 9x^2 + 8y^2 - 8y - 16 = 0$$

15. Let $A = \begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}$, $(\alpha \in \mathbb{R})$ such that

$$A^{32} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \text{ Then a value of } \alpha \text{ is}$$

(1) $\frac{\pi}{16}$ (2) 0

(3) $\frac{\pi}{32}$ (4) $\frac{\pi}{64}$

Official Ans. by NTA (4)

Sol. $A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$

$$A^2 = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix} \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

Sol. Consider $\cot^{-1} \left(\frac{\frac{\sqrt{3}}{2} \cos x + \frac{1}{2} \sin x}{\frac{1}{2} \sin x - \frac{\sqrt{3}}{2} \cos x} \right)$

$$= \cot^{-1} \left(\frac{\sin \left(x + \frac{\pi}{3} \right)}{\cos \left(x + \frac{\pi}{3} \right)} \right)$$

$$= \cot^{-1} \left(\tan \left(x + \frac{\pi}{3} \right) \right) = \frac{\pi}{2} - \tan^{-1} \left(\tan \left(x + \frac{\pi}{3} \right) \right)$$

$$\begin{cases} \frac{\pi}{2} - \left(x + \frac{\pi}{3} \right) = \left(\frac{\pi}{6} - x \right); \quad 0 < x < \frac{\pi}{6} \\ \frac{\pi}{2} - \left(\left(x - \frac{\pi}{3} \right) - \pi \right) = \left(\frac{7\pi}{6} - x \right); \quad \frac{\pi}{6} < x < \frac{\pi}{2} \end{cases}$$

$$\therefore 2y = \begin{cases} \left(\frac{\pi}{6} - x \right)^2; \quad 0 < x < \frac{\pi}{6} \\ \left(\frac{7\pi}{6} - x \right)^2; \quad \frac{\pi}{6} < x < \frac{\pi}{2} \end{cases}$$

$$\therefore 2 \frac{dy}{dx} = \begin{cases} 2 \left(\frac{\pi}{6} - x \right) \cdot (-1); \quad 0 < x < \frac{\pi}{6} \\ 2 \left(\frac{7\pi}{6} - x \right) \cdot (-1); \quad \frac{\pi}{6} < x < \frac{\pi}{2} \end{cases}$$

29. The greatest value of $c \in \mathbb{R}$ for which the system of linear equations

$$x - cy - cz = 0$$

$$cx - y + cz = 0$$

$$cx + cy - z = 0$$

has a non-trivial solution, is :

(1) $\frac{1}{2}$

(2) -1

(3) 0

(4) 2

Official Ans. by NTA (1)

Sol. For non-trivial solution

$$D = 0$$

$$\begin{vmatrix} 1 & -c & -c \\ c & -1 & c \\ c & c & -1 \end{vmatrix} = 0 \Rightarrow 2c^3 - 3c^2 - 1 = 0$$

$$\Rightarrow (c+1)^2(2c-1) = 0$$

$$\therefore \text{Greatest value of } c \text{ is } \frac{1}{2}$$

30. If $\cos(\alpha + \beta) = \frac{3}{5}$, $\sin(\alpha - \beta) = \frac{5}{13}$ and

$0 < \alpha, \beta < \frac{\pi}{4}$, then $\tan(2\alpha)$ is equal to :

(1) $\frac{21}{16}$

(2) $\frac{63}{52}$

(3) $\frac{33}{52}$

(4) $\frac{63}{16}$

Official Ans. by NTA (4)

Sol. $0 < \alpha + \beta = \frac{\pi}{2}$ and $-\frac{\pi}{4} < \alpha - \beta < \frac{\pi}{4}$

$$\text{if } \cos(\alpha + \beta) = \frac{3}{5} \text{ then } \tan(\alpha + \beta) = \frac{4}{3}$$

$$\text{and if } \sin(\alpha - \beta) = \frac{5}{13} \text{ then } \tan(\alpha - \beta) = \frac{5}{12}$$

(since $\alpha - \beta$ here lies in the first quadrant)

$$\text{Now } \tan(2\alpha) = \tan \{(\alpha + \beta) + (\alpha - \beta)\}$$

$$= \frac{\tan(\alpha + \beta) + \tan(\alpha - \beta)}{1 - \tan(\alpha + \beta) \cdot \tan(\alpha - \beta)} = \frac{\frac{4}{3} + \frac{5}{12}}{1 - \frac{4}{3} \cdot \frac{5}{12}} = \frac{63}{16}$$