

## FINAL JEE-MAIN EXAMINATION – APRIL, 2019

Held On Monday 08th APRIL, 2019

TIME: 2 : 30 PM To 5 : 30 PM

1. Calculate the standard cell potential in(V) of the cell in which following reaction takes place :  
 $\text{Fe}^{2+}(\text{aq}) + \text{Ag}^+(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + \text{Ag}(\text{s})$   
 Given that

$$E_{\text{Ag}^+/\text{Ag}}^{\circ} = x \text{ V}$$

$$E_{\text{Fe}^{2+}/\text{Fe}}^{\circ} = y \text{ V}$$

$$E_{\text{Fe}^{3+}/\text{Fe}}^{\circ} = z \text{ V}$$

(1)  $x + 2y - 3z$       (2)  $x - z$   
 (3)  $x - y$       (4)  $x + y - z$

**Official Ans. by NTA (1)**

**Sol.**  $\text{Fe}^{2+}(\text{aq}) + \text{Ag}^+(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + \text{Ag}(\text{s})$

Cell reaction

anode :  $\text{Fe}^{2+}(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + e^{\ominus}$  ;

$$E_{\text{Fe}^{2+}/\text{Fe}^{3+}}^{\circ} = m \text{ V}$$

cathode :  $\text{Ag}^+(\text{aq}) + e^{\ominus} \rightarrow \text{Ag}(\text{s})$  ;

$$E_{\text{Ag}^+/\text{Ag}}^{\circ} = x \text{ V}$$

$\Rightarrow$  cell standard potential =  $(m + x) \text{ V}$

$\therefore$  to find 'm' ;

$\text{Fe}^{2+} + 2e^{\ominus} \rightarrow \text{Fe}$  ;

$$E_1^{\circ} = y \text{ V} \Rightarrow \Delta_1^{\circ}G = -(2Fy)$$

$\text{Fe}^{3+} + 3e^{\ominus} \rightarrow \text{Fe}$  ;

$$E_2^{\circ} = z \text{ V} \Rightarrow \Delta_2^{\circ}G = -(3Fz)$$

$\text{Fe}^{2+}(\text{aq}) \rightarrow \text{Fe}^{3+}(\text{aq}) + e^{\ominus}$  ;






$$E_3^{\circ} = m \text{ V} \Rightarrow \Delta_3^{\circ}G = -(1Fm)$$

$$\Delta_3^{\circ}G = \Delta G_1^{\circ} - \Delta G_2^{\circ} = (-2Fy + 3Fz) = -Fm$$

$$\Rightarrow m = (2y - 3z)$$


$$\Rightarrow E_{\text{cell}}^{\circ} = (x + 2y - 3z) \text{ V}$$

2. The major product in the following reaction is :



**Official Ans. by NTA (2)**
**Ans. (Bonus)**

**Sol.** because one double bond is missing in all given option. So aromaticity is lost in both the ring.



3. For the following reactions, equilibrium constants are given :



The equilibrium constant for the reaction,




(1)  $10^{181}$    (2)  $10^{154}$    (3)  $10^{25}$    (4)  $10^{77}$

**Official Ans. by NTA (3)**


multiplying equation (1) by 2;



$\Rightarrow$  Subtracting (3) from (2); we get



$$K_{\text{eq}} = 10^{(129 - 104)} = 10^{25}$$

4. The ion that has  $sp^3d^2$  hybridization for the central atom, is :

- (1)  $[ICl_2]^-$
- (2)  $[IF_6]^-$
- (3)  $[ICl_4]^-$
- (4)  $[BrF_2]^-$


**Official Ans. by NTA (3)**

Sol.	Chemical species	Hybridisation of central atom
	$ICl_2^-$	$sp^3d$
	$IF_6^-$	$sp^3d^3$
	$ICl_4^-$	$sp^3d^2$
	$BrF_2^-$	$sp^3d$

5. The structure of Nylon-6 is :


- (1)  $\left[ -\left( CH_2 \right)_6 - C(=O) - N - \right]_n$
- (2)  $\left[ -\left( CH_2 \right)_4 - C(=O) - N - \right]_n$
- (3)  $\left[ -C(=O) - \left( CH_2 \right)_5 - N - \right]_n$
- (4)  $\left[ -C(=O) - \left( CH_2 \right)_6 - N - \right]_n$

**Official Ans. by NTA (3)**



Nylon-6


6. The major product of the following reaction is:




- (1)
- (2)
- (3)
- (4)

**Official Ans. by NTA (4)**

**Sol.**




7. The major product of the following reaction is:



- (1)
- (2)
- (3)
- (4)

**Official Ans. by NTA (4)**



8. The percentage composition of carbon by mole in methane is :  
 (1) 80% (2) 25% (3) 75% (4) 20%  
**Official Ans. by NTA (4)**

**Sol.**  $\text{CH}_4$

$$\% \text{ by mole of carbon} = \frac{1 \text{ mol atom}}{5 \text{ mol atom}} \times 100 \\ = 20\%$$

9. The IUPAC symbol for the element with atomic number 119 would be :  
 (1) unh (2) uun (3) une (4) uue  
**Official Ans. by NTA (4)**

Symbol	Atomic number
unh	106
uun	110
une	109
uue	119




10. The compound that inhibits the growth of tumors is :  
 (1) cis-[Pd(Cl)<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>]  
 (2) cis-[Pt(Cl)<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>]  
 (3) trans-[Pt(Cl)<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>]  
 (4) trans-[Pd(Cl)<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>]  
**Official Ans. by NTA (2)**

**Sol.** cis-[PtCl<sub>2</sub>(NH<sub>3</sub>)<sub>2</sub>] is used in chemotherapy to inhibits the growth of tumors.


11. The covalent alkaline earth metal halide ( $X = \text{Cl, Br, I}$ ) is :  
 (1)  $\text{CaX}_2$  (2)  $\text{SrX}_2$  (3)  $\text{BeX}_2$  (4)  $\text{MgX}_2$   
**Official Ans. by NTA (3)**

**Sol.** All halides of Be are predominantly covalent in nature.

12. The major product obtained in the following reaction is :



**Official Ans. by NTA (1)**



13. The statement that is **INCORRECT** about the interstitial compounds is :

- They have high melting points
- They are chemically reactive
- They have metallic conductivity
- They are very hard

**Official Ans. by NTA (2)**

**Sol.** Generally interstitial compounds are chemically inert.

14. The maximum prescribed concentration of copper in drinking water is:

- 5 ppm
- 0.5 ppm
- 0.05 ppm
- 3 ppm

**Official Ans. by NTA (4)**

**Sol.** The maximum prescribed concentration of Cu in drinking water is 3 ppm.

15. The calculated spin-only magnetic moments (BM) of the anionic and cationic species of  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$  and  $[\text{Fe}(\text{CN})_6]^{4-}$ , respectively, are :

- 4.9 and 0
- 2.84 and 5.92
- 0 and 4.9
- 0 and 5.92

**Official Ans. by NTA (3)**

**Sol.** Complex is  $[\text{Fe}(\text{H}_2\text{O})_6]^{2+} [\text{Fe}(\text{CN})_6]^{4-}$


Complex ion	Configuration	No. of unpaired electrons	Magnetic moment
$[\text{Fe}(\text{H}_2\text{O})_6]^{2+}$	$t_{2g}^4 e_g^2$	4	4.9 BM
$[\text{Fe}(\text{CN})_6]^{4-}$	$t_{2g}^6 e_g^0$	0	0



20. For a reaction scheme  $A \xrightarrow{k_1} B \xrightarrow{k_2} C$ , if the rate of formation of B is set to be zero then the concentration of B is given by :

(1)  $\left(\frac{k_1}{k_2}\right)[A]$  (2)  $(k_1 + k_2)[A]$   
 (3)  $k_1 k_2 [A]$  (4)  $(k_1 - k_2)[A]$

**Official Ans. by NTA (1)**



$$\frac{d[B]}{dt} = 0 = K_1[A] - K_2[B]$$


$$\Rightarrow [B] = \frac{K_1}{K_2}[A]$$

21. Which of the following compounds will show the maximum enol content?

(1)  $\text{CH}_3\text{COCH}_2\text{COCH}_3$   
 (2)  $\text{CH}_3\text{COCH}_3$   
 (3)  $\text{CH}_3\text{COCH}_2\text{CONH}_2$   
 (4)  $\text{CH}_3\text{COCH}_2\text{COOC}_2\text{H}_5$

**Official Ans. by NTA (1)**

Sol. Solution



Due to intramolecular H-bonding and resonance stabilisation enol content is maximum


22. The correct statement about  $\text{ICl}_5$  and  $\text{ICl}_4^-$  is

(1)  $\text{ICl}_5$  is trigonal bipyramidal and  $\text{ICl}_4^-$  is tetrahedral.  
 (2)  $\text{ICl}_5$  is square pyramidal and  $\text{ICl}_4^-$  is tetrahedral.  
 (3)  $\text{ICl}_5$  is square pyramidal and  $\text{ICl}_4^-$  is square planar.  
 (4) Both are isostructural.






**Official Ans. by NTA (3)**

**Sol.**

Chemical species	Hybridisation	Shape
$\text{ICl}_5$	$\text{sp}^3\text{d}^2$	Square pyramidal
$\text{ICl}_4^-$	$\text{sp}^3\text{d}^2$	Square planar




23. The major product obtained in the following reaction is



**Official Ans. by NTA (4)**

**Sol.**



24. Fructose and glucose can be distinguished by :

- Fehling's test
- Barfoed's test
- Benedict's test
- Seliwanoff's test

**Official Ans. by NTA (4)**

**Sol.** Seliwanoff's test is used to distinguish aldose and ketose group.

25. If  $p$  is the momentum of the fastest electron ejected from a metal surface after the irradiation of light having wavelength  $\lambda$ , then for 1.5  $p$  momentum of the photoelectron, the wavelength of the light should be:

(Assume kinetic energy of ejected photoelectron to be very high in comparison to work function)

$$(1) \frac{1}{2}\lambda \quad (2) \frac{3}{4}\lambda$$

$$(3) \frac{2}{3}\lambda \quad (4) \frac{4}{9}\lambda$$

**Official Ans. by NTA (4)**

**Sol.**  $h\nu - \phi = KE$

$$\Rightarrow \left( \frac{hc}{\lambda} \right)_{\text{incident}} = KE + \phi$$

$$\left( \frac{hc}{\lambda} \right)_{\text{incident}} \approx KE$$

$$KE = \frac{p^2}{2m} = \frac{hc}{\lambda_{\text{incident}}} = \frac{hc}{\lambda} \quad \dots(1)$$


$$\Rightarrow \frac{p^2 \times (1.5)^2}{2m} = \frac{hc}{\lambda'} \quad \dots(2)$$

divide (1) and (2)

$$(1.5)^2 = \frac{\lambda}{\lambda'}$$

$$\Rightarrow \lambda' = \frac{4\lambda}{9}$$

26. Consider the bcc unit cells of the solids 1 and 2 with the position of atoms as shown below. The radius of atom B is twice that of atom A. The unit cell edge length is 50% more in solid 2 than in 1. What is the approximate packing efficiency in solid 2?



(1) 45% (2) 65% (3) 90% (4) 75%

**Official Ans. by NTA (3)**

$$\text{Sol. p.f.} = \frac{\left( z_{\text{eff}} \times \frac{4}{3} \pi r_A^3 \right)_A + \left( z_{\text{eff}} \times \frac{4}{3} \pi r_B^3 \right)_B}{a^3}$$

$$2(r_A + r_B) = \sqrt{3}a$$

$$\Rightarrow 2(r_A + 2r_A) = \sqrt{3}a$$

$$\Rightarrow 2\sqrt{3}r_A = a$$

$$\Rightarrow \text{p.f.} = \frac{1 \times \frac{4}{3} \pi r_A^3 + \frac{4}{3} \pi (8r_A^3)}{8 \times 3\sqrt{3}r_A^3} = \frac{9 \times \frac{4}{3} \pi}{8 \times 3\sqrt{3}} = \frac{\pi}{2\sqrt{3}}$$

$$\text{p. efficiency} = \frac{\pi}{2\sqrt{3}} \times 100 \approx 90\%$$

27. Polysubstitution is a major drawback in:

- Reimer Tiemann reaction
- Friedel Craft's acylation
- Friedel Craft's alkylation
- Acetylation of aniline

**Official Ans. by NTA (3)**

**Sol.** In Friedel crafts alkylation product obtained is more activated and hence polysubstitution will take place.

28. The Mond process is used for the

- extraction of Mo
- Purification of Ni
- Purification of Zr and Ti
- Extraction of Zn

**Official Ans. by NTA (2)**

**Sol.** Mond's process is used for the purification of Nickel.

29. The strength of 11.2 volume solution of  $\text{H}_2\text{O}_2$  is : [Given that molar mass of H = 1 g  $\text{mol}^{-1}$  and O = 16 g  $\text{mol}^{-1}$ ]

- 13.6%
- 3.4%
- 34%
- 1.7%

**Official Ans. by NTA (2)**





**Sol.** Volume strength =  $11.2 \times$  molarity = 11.2

$$\Rightarrow \text{molarity} = 1 \text{ M}$$

$$\Rightarrow \text{strength} = 34 \text{ g/L}$$

$$\Rightarrow \% \text{ w/w} = \frac{34}{1000} \times 100 = 3.4\%$$

30. For the solution of the gases w, x, y and z in water at 298K, the Henry's law constants ( $K_H$ ) are 0.5, 2, 35 and 40 kbar, respectively. The correct plot for the given data is :-



**Official Ans. by NTA (3)**

**Sol.** 
$$p = k_H \times \left( \frac{n_{\text{gas}}}{n_{\text{H}_2\text{O}} + n_{\text{gas}}} \right)$$

$$= k_H \left( 1 - \frac{n_{\text{H}_2\text{O}}}{n_{\text{H}_2\text{O}} + n_{\text{gas}}} \right)$$

$$\Rightarrow p = k_H - k_H \times \chi_{\text{H}_2\text{O}}$$

$$p = (-k_H) \times \chi_{\text{H}_2\text{O}} + k_H$$