



FINAL JEE–MAIN EXAMINATION – JULY, 2021

Held On Tuesday 27th July, 2021

TIME: 9:00 AM to 12:00 NOON

SECTION-A

1. If the mean and variance of the following data:
6, 10, 7, 13, a, 12, b, 12
are 9 and $\frac{37}{4}$ respectively, then $(a - b)^2$ is equal to:

- (1) 24 (2) 12 (3) 32 (4) 16

Official Ans. by NTA (4)

2. The value of $\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \frac{(2j-1) + 8n}{(2j-1) + 4n}$ is equal to :

- (1) $5 + \log_e \left(\frac{3}{2}\right)$ (2) $2 - \log_e \left(\frac{2}{3}\right)$
(3) $3 + 2 \log_e \left(\frac{2}{3}\right)$ (4) $1 + 2 \log_e \left(\frac{3}{2}\right)$

Official Ans. by NTA (4)

3. Let $\vec{a} = \hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = -\hat{i} + 2\hat{j} + 3\hat{k}$. Then the vector product $(\vec{a} + \vec{b}) \times \left((\vec{a} \times ((\vec{a} - \vec{b}) \times \vec{b})) \times \vec{b} \right)$ is equal to :

- (1) $5(34\hat{i} - 5\hat{j} + 3\hat{k})$ (2) $7(34\hat{i} - 5\hat{j} + 3\hat{k})$
(3) $7(30\hat{i} - 5\hat{j} + 7\hat{k})$ (4) $5(30\hat{i} - 5\hat{j} + 7\hat{k})$

Official Ans. by NTA (2)

4. The value of the definite integral

$$\int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{dx}{(1 + e^{x \cos x})(\sin^4 x + \cos^4 x)}$$

is equal to :

- (1) $-\frac{\pi}{2}$ (2) $\frac{\pi}{2\sqrt{2}}$ (3) $-\frac{\pi}{4}$ (4) $\frac{\pi}{\sqrt{2}}$

Official Ans. by NTA (2)

5. Let C be the set of all complex numbers. Let
 $S_1 = \{z \in C \mid |z - 3 - 2i|^2 = 8\}$,
 $S_2 = \{z \in C \mid \text{Re}(z) \geq 5\}$ and
 $S_3 = \{z \in C \mid |z - \bar{z}| \geq 8\}$.

Then the number of elements in $S_1 \cap S_2 \cap S_3$ is equal to

- (1) 1 (2) 0 (3) 2 (4) Infinite

Official Ans. by NTA (1)

6. If the area of the bounded region

$$R = \left\{ (x, y) : \max\{0, \log_e x\} \leq y \leq 2^x, \frac{1}{2} \leq x \leq 2 \right\}$$

is, $\alpha(\log_e 2)^{-1} + \beta(\log_e 2) + \gamma$, then the value of $(\alpha + \beta - 2\gamma)^2$ is equal to :

- (1) 8 (2) 2 (3) 4 (4) 1

Official Ans. by NTA (2)

7. A ray of light through (2,1) is reflected at a point P on the y-axis and then passes through the point (5, 3). If this reflected ray is the directrix of an ellipse with eccentricity $\frac{1}{3}$ and the distance of the

nearer focus from this directrix is $\frac{8}{\sqrt{53}}$, then the

equation of the other directrix can be:

- (1) $11x + 7y + 8 = 0$ or $11x + 7y - 15 = 0$
(2) $11x - 7y - 8 = 0$ or $11x + 7y + 15 = 0$
(3) $2x - 7y + 29 = 0$ or $2x - 7y - 7 = 0$
(4) $2x - 7y - 39 = 0$ or $2x - 7y - 7 = 0$

Official Ans. by NTA (3)

8. If the coefficients of x^7 in $\left(x^2 + \frac{1}{bx}\right)^{11}$ and x^{-7} in

$$\left(x - \frac{1}{bx^2}\right)^{11}, b \neq 0, \text{ are equal, then the value of } b$$

is equal to:

- (1) 2 (2) -1 (3) 1 (4) -2

Official Ans. by NTA (3)

9. The compound statement $(P \vee Q) \wedge (\sim P) \Rightarrow Q$ is equivalent to:

- (1) $P \vee Q$ (2) $P \wedge \sim Q$
(3) $\sim(P \Rightarrow Q)$ (4) $\sim(P \Rightarrow Q) \Leftrightarrow P \wedge \sim Q$

Official Ans. by NTA (4)

10. If $\sin \theta + \cos \theta = \frac{1}{2}$, then

$16(\sin(2\theta) + \cos(4\theta) + \sin(6\theta))$ is equal to:

- (1) 23 (2) -27 (3) -23 (4) 27

Official Ans. by NTA (3)



SECTION-B

1. For real numbers α and β , consider the following system of linear equations :

$$x + y - z = 2, x + 2y + \alpha z = 1, 2x - y + z = \beta.$$

If the system has infinite solutions, then $\alpha + \beta$ is equal to _____.

Official Ans. by NTA (5)

2. Let $\vec{a} = \hat{i} + \hat{j} + \hat{k}$, \vec{b} and $\vec{c} = \hat{j} - \hat{k}$ be three vectors such that $\vec{a} \times \vec{b} = \vec{c}$ and $\vec{a} \cdot \vec{b} = 1$. If the length of projection vector of the vector \vec{b} on the vector $\vec{a} \times \vec{c}$ is l , then the value of $3l^2$ is equal to _____.

Official Ans. by NTA (2)

3. If $\log_3 2, \log_3(2^x - 5), \log_3\left(2^x - \frac{7}{2}\right)$ are in an arithmetic progression, then the value of x is equal to _____.

Official Ans. by NTA (3)

4. Let the domain of the function $f(x) = \log_4\left(\log_5\left(\log_3(18x - x^2 - 77)\right)\right)$ be (a, b) .

Then the value of the integral

$$\int_a^b \frac{\sin^3 x}{(\sin^3 x + \sin^3(a + b - x))} dx \text{ is equal to } \underline{\hspace{2cm}}.$$

Official Ans. by NTA (1)

5. Let

$$f(x) = \begin{vmatrix} \sin^2 x & -2 + \cos^2 x & \cos 2x \\ 2 + \sin^2 x & \cos^2 x & \cos 2x \\ \sin^2 x & \cos^2 x & 1 + \cos 2x \end{vmatrix}, x \in [0, \pi]$$

Then the maximum value of $f(x)$ is equal to _____.

Official Ans. by NTA (6)

6. Let $F : [3, 5] \rightarrow \mathbf{R}$ be a twice differentiable function on $(3, 5)$ such that

$$F(x) = e^{-x} \int_3^x (3t^2 + 2t + 4F'(t)) dt.$$

If $F'(4) = \frac{\alpha e^\beta - 224}{(e^\beta - 4)^2}$, then $\alpha + \beta$ is equal to _____.

Official Ans. by NTA (16)

7. Let a plane P pass through the point $(3, 7, -7)$ and contain the line, $\frac{x-2}{-3} = \frac{y-3}{2} = \frac{z+2}{1}$. If distance of the plane P from the origin is d , then d^2 is equal to _____.

Official Ans. by NTA (3)

8. Let $S = \{1, 2, 3, 4, 5, 6, 7\}$. Then the number of possible functions $f : S \rightarrow S$ such that $f(m \cdot n) = f(m) \cdot f(n)$ for every $m, n \in S$ and $m \cdot n \in S$ is equal to _____.

Official Ans. by NTA (490)

9. If $y = y(x)$, $y \in \left[0, \frac{\pi}{2}\right)$ is the solution of the differential equation

$$\sec y \frac{dy}{dx} - \sin(x+y) - \sin(x-y) = 0, \text{ with } y(0) = 0,$$

then $5y'\left(\frac{\pi}{2}\right)$ is equal to _____.

Official Ans. by NTA (2)

10. Let $f : [0, 3] \rightarrow \mathbf{R}$ be defined by

$$f(x) = \min \{x - [x], 1 + [x] - x\}$$

where $[x]$ is the greatest integer less than or equal to x . Let P denote the set containing all $x \in [0, 3]$ where f is discontinuous, and Q denote the set containing all $x \in (0, 3)$ where f is not differentiable. Then the sum of number of elements in P and Q is equal to _____.

Official Ans. by NTA (5)