



FINAL JEE–MAIN EXAMINATION – AUGUST, 2021

Held On Friday 27th August, 2021

TIME: 3:00 PM to 06:00 PM

SECTION-A

1. The angle between the straight lines, whose direction cosines are given by the equations $2l + 2m - n = 0$ and $mn + nl + lm = 0$, is :

- (1) $\frac{\pi}{2}$ (2) $\pi - \cos^{-1}\left(\frac{4}{9}\right)$
 (3) $\cos^{-1}\left(\frac{8}{9}\right)$ (4) $\frac{\pi}{3}$

Official Ans. by NTA (1)

Sol. $n = 2(l + m)$

$$lm + n(l + m) = 0$$

$$lm + 2(l + m)^2 = 0$$

$$2l^2 + 2m^2 + 5ml = 0$$

$$2\left(\frac{l}{m}\right)^2 + 2 + 5\left(\frac{l}{m}\right) = 0.$$

$$2t^2 + 5t + 2 = 0$$

$$(t + 2)(2t + 1) = 0$$

$$\Rightarrow t = -2; -\frac{1}{2}$$

(i) $\frac{l}{m} = -2$	(ii) $\frac{l}{m} = -\frac{1}{2}$
$\frac{n}{m} = -2$	
$(-2m, m, -2m)$	
$(-2, 1, -2)$	
$n = -2l$	
	$(l, -2l, -2l)$
	$(1, -2, -2)$

$$\cos\theta = \frac{-2 - 2 + 4}{\sqrt{9} \sqrt{9}} = 0 \Rightarrow \theta = \frac{\pi}{2}$$

2. Let $A = \begin{bmatrix} [x+1] & [x+2] & [x+3] \\ [x] & [x+3] & [x+3] \\ [x] & [x+2] & [x+4] \end{bmatrix}$, where $[t]$

denotes the greatest integer less than or equal to t . If $\det(A) = 192$, then the set of values of x is the interval:

- (1) [68, 69] (2) [62, 63]
 (3) [65, 66] (4) [60, 61]

Official Ans. by NTA (2)

Sol. $\begin{vmatrix} [x+1] & [x+2] & [x+3] \\ [x] & [x+3] & [x+3] \\ [x] & [x+2] & [x+4] \end{vmatrix} = 192$

$$R_1 \rightarrow R_1 - R_3 \text{ \& } R_2 \rightarrow R_2 - R_3$$

$$\begin{vmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \\ [x] & [x]+2 & [x]+4 \end{vmatrix} = 192$$

$$2[x] + 6 + [x] = 192 \Rightarrow [x] = 62$$

3. Let M and m respectively be the maximum and minimum values of the function $f(x) = \tan^{-1}(\sin x + \cos x)$ in $\left[0, \frac{\pi}{2}\right]$, Then the value of $\tan(M - m)$ is equal to:

- (1) $2 + \sqrt{3}$ (2) $2 - \sqrt{3}$
 (3) $3 + 2\sqrt{2}$ (4) $3 - 2\sqrt{2}$

Official Ans. by NTA (4)

Sol. Let $g(x) = \sin x + \cos x = \sqrt{2} \sin\left(x + \frac{\pi}{4}\right)$

$$g(x) \in [1, \sqrt{2}] \text{ for } x \in [0, \pi/2]$$

$$f(x) = \tan^{-1}(\sin x + \cos x) \in \left[\frac{\pi}{4}, \tan^{-1}\sqrt{2}\right]$$

$$\tan\left(\tan^{-1}\sqrt{2} - \frac{\pi}{4}\right) = \frac{\sqrt{2}-1}{1+\sqrt{2}} \times \frac{\sqrt{2}-1}{\sqrt{2}-1} = 3 - 2\sqrt{2}$$

4. Each of the persons A and B independently tosses three fair coins. The probability that both of them get the same number of heads is :

- (1) $\frac{1}{8}$ (2) $\frac{5}{8}$ (3) $\frac{5}{16}$ (4) 1

Official Ans. by NTA (3)



Sol. C - I '0' Head

$$T T T \quad \left(\frac{1}{2}\right)^3 \left(\frac{1}{2}\right)^3 = \frac{1}{64}$$

C - II '1' head

$$H T T \quad \left(\frac{3}{8}\right)\left(\frac{3}{8}\right) = \frac{9}{64}$$

C - III '2' Head

$$H H T \quad \left(\frac{3}{8}\right)\left(\frac{3}{8}\right) = \frac{9}{64}$$

C-IV '3' Heads

$$H H H \quad \left(\frac{1}{8}\right)\left(\frac{1}{8}\right) = \frac{1}{64}$$

$$\text{Total probability} = \frac{5}{16}.$$

5. A differential equation representing the family of parabolas with axis parallel to y-axis and whose length of latus rectum is the distance of the point (2, -3) from the line $3x + 4y = 5$, is given by :

$$(1) 10 \frac{d^2y}{dx^2} = 11 \quad (2) 11 \frac{d^2x}{dy^2} = 10$$

$$(3) 10 \frac{d^2x}{dy^2} = 11 \quad (4) 11 \frac{d^2y}{dx^2} = 10$$

Official Ans. by NTA (4)

Sol. $\alpha. R = \frac{|3(2) + 4(-3) - 5|}{5} = \frac{11}{5}$

$$(x-h)^2 = \frac{11}{5}(y-k)$$

differentiate w.r.t 'x' :-

$$2(x-h) = \frac{11}{5} \frac{dy}{dx}$$

again differentiate

$$2 = \frac{11}{5} \frac{d^2y}{dx^2}$$

$$\frac{11d^2y}{dx^2} = 10.$$

6. If two tangents drawn from a point P to the parabola $y^2 = 16(x - 3)$ are at right angles, then the locus of point P is :

$$(1) x + 3 = 0 \quad (2) x + 1 = 0$$

$$(3) x + 2 = 0 \quad (4) x + 4 = 0$$

Official Ans. by NTA (2)

Sol. Locus is directrix of parabola

$$x - 3 + 4 = 0 \Rightarrow x + 1 = 0.$$

7. The equation of the plane passing through the line of intersection of the planes $\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) = 1$ and

$$\vec{r} \cdot (2\hat{i} + 3\hat{j} - \hat{k}) + 4 = 0$$

$$(1) \vec{r} \cdot (\hat{j} - 3\hat{k}) + 6 = 0 \quad (2) \vec{r} \cdot (\hat{i} + 3\hat{k}) + 6 = 0$$

$$(3) \vec{r} \cdot (\hat{i} - 3\hat{k}) + 6 = 0 \quad (4) \vec{r} \cdot (\hat{j} - 3\hat{k}) - 6 = 0$$

Official Ans. by NTA (1)

Sol. Equation of planes are

$$\vec{r} \cdot (\hat{i} + \hat{j} + \hat{k}) - 1 = 0 \Rightarrow x + y + z - 1 = 0$$

$$\text{and } \vec{r} \cdot (2\hat{i} + 3\hat{j} - \hat{k}) + 4 = 0 \Rightarrow 2x + 3y - z + 4 = 0$$

equation of planes through line of intersection of these planes is :-

$$(x + y + z - 1) + \lambda (2x + 3y - z + 4) = 0$$

$$\Rightarrow (1 + 2\lambda)x + (1 + 3\lambda)y + (1 - \lambda)z - 1 + 4\lambda = 0$$

But this plane is parallel to x-axis whose direction are (1, 0, 0)

$$\therefore (1 + 2\lambda)1 + (1 + 3\lambda)0 + (1 - \lambda)0 = 0$$

$$\lambda = -\frac{1}{2}$$

\therefore Required plane is

$$0x + \left(1 - \frac{3}{2}\right)y + \left(1 + \frac{1}{2}\right)z - 1 + 4\left(\frac{-1}{2}\right) = 0$$

$$\Rightarrow \frac{-y}{2} + \frac{3}{2}z - 3 = 0$$

$$\Rightarrow y - 3z + 6 = 0$$

$$\Rightarrow \boxed{\vec{r} \cdot (\hat{j} - 3\hat{k}) + 6 = 0} \text{ Ans.}$$



8. If the solution curve of the differential equation $(2x - 10y^3) dy + y dx = 0$, passes through the points $(0, 1)$ and $(2, \beta)$, then β is a root of the equation:

- (1) $y^5 - 2y - 2 = 0$ (2) $2y^5 - 2y - 1 = 0$
 (3) $2y^5 - y^2 - 2 = 0$ (4) $y^5 - y^2 - 1 = 0$

Official Ans. by NTA (4)

Sol. $(2x - 10y^3) dy + y dx = 0$

$$\Rightarrow \frac{dx}{dy} + \left(\frac{2}{y}\right)x = 10y^2$$

$$\text{I. F.} = e^{\int \frac{2}{y} dy} = e^{2 \ln(y)} = y^2$$

Solution of D.E. is

$$\therefore x \cdot y = \int (10y^2) y^2 \cdot dy$$

$$xy^2 = \frac{10y^5}{5} + C \Rightarrow xy^2 = 2y^5 + C$$

It passes through $(0, 1) \rightarrow 0 = 2 + C \Rightarrow C = -2$

$$\therefore \text{Curve is } \boxed{xy^2 = 2y^5 - 2}$$

Now, it passes through $(2, \beta)$

$$2\beta^2 = 2\beta^5 - 2 \Rightarrow \beta^5 - \beta^2 - 1 = 0$$

$$\therefore \beta \text{ is root of an equation } \boxed{y^5 - y^2 - 1 = 0} \text{ Ans.}$$

9. Let $A(a, 0)$, $B(b, 2b + 1)$ and $C(0, b)$, $b \neq 0$, $|b| \neq 1$, be points such that the area of triangle ABC is 1 sq. unit, then the sum of all possible values of a is :

- (1) $\frac{-2b}{b+1}$ (2) $\frac{2b}{b+1}$
 (3) $\frac{2b^2}{b+1}$ (4) $\frac{-2b^2}{b+1}$

Official Ans. by NTA (4)

$$\text{Sol. } \left| \begin{array}{ccc} a & 0 & 1 \\ b & 2b+1 & 1 \\ 0 & b & 1 \end{array} \right| = 1$$

$$\Rightarrow \left| \begin{array}{ccc} a & 0 & 1 \\ b & 2b+1 & 1 \\ 0 & b & 1 \end{array} \right| = \pm 2$$

$$\Rightarrow a(2b + 1 - b) - 0 + 1(b^2 - 0) = \pm 2$$

$$\Rightarrow a = \frac{\pm 2 - b^2}{b + 1}$$

$$\therefore a = \frac{2 - b^2}{b + 1} \text{ and } a = \frac{-2 - b^2}{b + 1}$$

sum of possible values of 'a' is

$$= \frac{-2b^2}{a + 1} \text{ Ans.}$$

10. Let $[\lambda]$ be the greatest integer less than or equal to λ . The set of all values of λ for which the system of linear equations $x + y + z = 4$, $3x + 2y + 5z = 3$, $9x + 4y + (28 + [\lambda])z = [\lambda]$ has a solution is:

- (1) \mathbf{R}
 (2) $(-\infty, -9) \cup (-9, \infty)$
 (3) $[-9, -8]$
 (4) $(-\infty, -9) \cup [-8, \infty)$

Official Ans. by NTA (1)

$$\text{Sol. } D = \begin{vmatrix} 1 & 1 & 1 \\ 3 & 2 & 5 \\ 9 & 4 & 28 + [\lambda] \end{vmatrix} = -24 - [\lambda] + 15 = -[\lambda] - 9$$

if $[\lambda] + 9 \neq 0$ then unique solution

if $[\lambda] + 9 = 0$ then $D_1 = D_2 = D_3 = 0$

so infinite solutions

Hence λ can be any real number.

11. The set of all values of $k > -1$, for which the equation $(3x^2 + 4x + 3)^2 - (k + 1)(3x^2 + 4x + 3)(3x^2 + 4x + 2) + k(3x^2 + 4x + 2)^2 = 0$ has real roots, is :

- (1) $\left[1, \frac{5}{2}\right]$ (2) $[2, 3]$
 (3) $\left[-\frac{1}{2}, 1\right]$ (4) $\left[\frac{1}{2}, \frac{3}{2}\right] - \{1\}$

Official Ans. by NTA (1)

$$\text{Sol. } (3x^2 + 4x + 3)^2 - (k + 1)(3x^2 + 4x + 3)(3x^2 + 4x + 2) + k(3x^2 + 4x + 2)^2 = 0$$

$$\text{Let } 3x^2 + 4x + 3 = a$$

$$\text{and } 3x^2 + 4x + 2 = b \Rightarrow b = a - 1$$

Given equation becomes

$$\Rightarrow a^2 - (k + 1)ab + kb^2 = 0$$

$$\Rightarrow a(a - kb) - b(a - kb) = 0$$

$$\Rightarrow (a - kb)(a - b) = 0 \Rightarrow a = kb \text{ or } a = b \text{ (reject)}$$

$$\therefore a = kb$$

$$\Rightarrow 3x^2 + 4x + 3 = k(3x^2 + 4x + 2)$$

$$\Rightarrow 3(k - 1)x^2 + 4(k - 1)x + (2k - 3) = 0$$

for real roots

$$D \geq 0$$


$$\Rightarrow 16(k - 1)^2 - 4(3(k - 1))(2k - 3) \geq 0$$

$$\Rightarrow 4(k - 1)\{4(k - 1) - 3(2k - 3)\} \geq 0$$

$$\Rightarrow 4(k - 1)\{-2k + 5\} \geq 0$$

$$\Rightarrow -4(k - 1)\{2k - 5\} \geq 0$$

$$\Rightarrow (k - 1)(2k - 5) \leq 0$$



$$\therefore k \in \left[1, \frac{5}{2}\right]$$

$$\therefore k \neq 1$$


$$\therefore k \in \left(1, \frac{5}{2}\right] \text{ Ans.}$$



12. A box open from top is made from a rectangular sheet of dimension $a \times b$ by cutting squares each of side x from each of the four corners and folding up the flaps. If the volume of the box is maximum, then x is equal to :

- (1) $\frac{a+b-\sqrt{a^2+b^2-ab}}{12}$
- (2) $\frac{a+b-\sqrt{a^2+b^2+ab}}{6}$
- (3) $\frac{a+b-\sqrt{a^2+b^2-ab}}{6}$
- (4) $\frac{a+b+\sqrt{a^2+b^2-ab}}{6}$

Official Ans. by NTA (3)



$$V = \ell. b. h = (a - 2x)(b - 2x)x$$

$$\Rightarrow V(x) = (2x - a)(2x - b)x$$

$$\Rightarrow V(x) = 4x^3 - 2(a + b)x^2 + abx$$

$$\Rightarrow \frac{d}{dx} v(x) = 12x^2 - 4(a + b)x + ab$$

$$\frac{d}{dx} (v(x)) = 0 \Rightarrow 12x^2 - 4(a + b)x + ab = 0 \dots \alpha$$

$$\Rightarrow x = \frac{4(a + b) \pm \sqrt{16(a + b)^2 - 48ab}}{2(12)}$$

$$= \frac{(a + b) \pm \sqrt{a^2 + b^2 - ab}}{6}$$

Let $x = \alpha = \frac{(a + b) + \sqrt{a^2 + b^2 - ab}}{6}$

$$\beta = \frac{(a + b) - \sqrt{a^2 + b^2 - ab}}{6}$$

Now, $12(x - \alpha)(x - \beta) = 0$

$\leftarrow \begin{array}{ccc} + & - & + \\ | & & | \\ \beta & & \alpha \\ \downarrow & & \downarrow \\ \text{Maxima} & & \text{minima} \end{array} \rightarrow x$

$$\therefore x = \beta$$

$$= \frac{a + b - \sqrt{a^2 + b^2 - ab}}{6}$$

13. The Boolean expression $(p \wedge q) \Rightarrow ((r \wedge q) \wedge p)$ is equivalent to :

- (1) $(p \wedge q) \Rightarrow (r \wedge q)$
- (2) $(q \wedge r) \Rightarrow (p \wedge q)$
- (3) $(p \wedge q) \Rightarrow (r \vee q)$
- (4) $(p \wedge r) \Rightarrow (p \wedge q)$

Official Ans. by NTA (1)

Sol.

$$(p \wedge q) \Rightarrow ((r \wedge q) \wedge p)$$

$$\sim (p \wedge q) \vee ((r \wedge q) \wedge p)$$

$$\sim (p \wedge q) \vee ((r \wedge p) \wedge (p \wedge q))$$

$$\Rightarrow [\sim (p \wedge q) \vee (p \wedge q)] \wedge (\sim (p \wedge q) \vee (r \wedge p))$$

$$\Rightarrow t \wedge [\sim (p \wedge q) \vee (r \wedge p)]$$

$$\Rightarrow \sim (p \wedge q) \vee (r \wedge p)$$

$$\Rightarrow (p \wedge q) \Rightarrow (r \wedge p)$$

Aliter :

given statement says

" if p and q both happen then p and q and r will happen"

it Simply implies

" If p and q both happen then 'r' too will happen "

i.e.

" if p and q both happen then r and p too will happen

i.e.

$$(p \wedge q) \Rightarrow (r \wedge p)$$

14. Let \mathbb{Z} be the set of all integers,

$$A = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} : (x - 2)^2 + y^2 \leq 4\},$$

$$B = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} : x^2 + y^2 \leq 4\} \text{ and}$$

$$C = \{(x, y) \in \mathbb{Z} \times \mathbb{Z} : (x - 2)^2 + (y - 2)^2 \leq 4\}$$

If the total number of relation from $A \cap B$ to


$A \cap C$ is 2^p , then the value of p is :

- (1) 16
- (2) 25
- (3) 49
- (4) 9

Official Ans. by NTA (2)



Sol.



$$(x - 2)^2 + y^2 \leq 4$$

$$x^2 + y^2 \leq 4$$

No. of points common in C_1 & C_2 is 5.

$(0, 0), (1, 0), (2, 0), (1, 1), (1, -1)$

Similarly in C_2 & C_3 is 5.

No. of relations = $2^{5 \times 5} = 2^{25}$.

15. The area of the region bounded by the parabola $(y-2)^2 = (x-1)$, the tangent to it at the point whose ordinate is 3 and the x-axis is :

- (1) 9 (2) 10 (3) 4 (4) 6

Official Ans. by NTA (1)

Sol. $y = 3 \Rightarrow x = 2$

Point is $(2, 3)$

Diff. w.r.t x

$$2(y - 2)y' = 1$$


$$\Rightarrow y' = \frac{1}{2(y-2)}$$

$$\Rightarrow y'_{(2,3)} = \frac{1}{2}$$

$$\Rightarrow \frac{y-3}{x-2} = \frac{1}{2} \Rightarrow x - 2y + 4 = 0$$

$$\text{Area} = \int_0^3 \left((y-2)^2 + 1 - (2y-4) \right) dy$$

$$= 9 \text{ sq. units}$$



16. If $y(x) = \cot^{-1} \left(\frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}} \right)$, $x \in \left(\frac{\pi}{2}, \pi \right)$,

then $\frac{dy}{dx}$ at $x = \frac{5\pi}{6}$ is:

- (1) $-\frac{1}{2}$ (2) -1 (3) $\frac{1}{2}$ (4) 0

Official Ans. by NTA (1)

Sol. $y(x) = \cot^{-1} \left[\frac{\cos \frac{x}{2} + \sin \frac{x}{2} + \sin \frac{x}{2} - \cos \frac{x}{2}}{\cos \frac{x}{2} + \sin \frac{x}{2} - \sin \frac{x}{2} + \cos \frac{x}{2}} \right]$

$$y(x) = \cot^{-1} \left(\tan \frac{x}{2} \right) = \frac{\pi}{2} - \frac{x}{2}$$


$$y'(x) = \frac{-1}{2}$$

17. Two poles, AB of length a metres and CD of length a + b ($b \neq a$) metres are erected at the same horizontal level with bases at B and D. If $BD = x$

and $\tan \angle ACB = \frac{1}{2}$, then:

- (1) $x^2 + 2(a + 2b)x - b(a + b) = 0$
 (2) $x^2 + 2(a + 2b)x + a(a + b) = 0$
 (3) $x^2 - 2ax + b(a + b) = 0$
 (4) $x^2 - 2ax + a(a + b) = 0$

Official Ans. by NTA (3)



$$\tan \theta = \frac{1}{2}$$

$$\tan(\theta + \alpha) = \frac{x}{b}, \quad \tan \alpha = \frac{x}{a+b}$$

$$\Rightarrow \frac{1}{2} + \frac{x}{a+b}$$

$$\Rightarrow \frac{\frac{1}{2} + \frac{x}{a+b}}{1 - \frac{1}{2} \times \frac{x}{a+b}} = \frac{x}{b}$$

$$\Rightarrow x^2 - 2ax + ab + b^2 = 0$$



18. If $0 < x < 1$ and $y = \frac{1}{2}x^2 + \frac{2}{3}x^3 + \frac{3}{4}x^4 + \dots$, then

the value of e^{1+y} at $x = \frac{1}{2}$ is:

- (1) $\frac{1}{2}e^2$ (2) $2e$
 (3) $\frac{1}{2}\sqrt{e}$ (4) $2e^2$

Official Ans. by NTA (1)

Sol. $y = \left(1 - \frac{1}{2}\right)x^2 + \left(1 - \frac{1}{3}\right)x^3 + \dots$
 $= (x^2 + x^3 + x^4 + \dots) - \left(\frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots\right)$
 $= \frac{x^2}{1-x} + x - \left(x + \frac{x^2}{2} + \frac{x^2}{3} + \dots\right)$
 $= \frac{x}{1-x} + \ln(1-x)$
 $x = \frac{1}{2} \Rightarrow y = 1 - \ln 2$
 $e^{1+y} = e^{1+1-\ln 2}$
 $= e^{2-\ln 2} = \frac{e^2}{2}$

19. The value of the integral $\int_0^1 \frac{\sqrt{x} dx}{(1+x)(1+3x)(3+x)}$

is:

- (1) $\frac{\pi}{8} \left(1 - \frac{\sqrt{3}}{2}\right)$ (2) $\frac{\pi}{4} \left(1 - \frac{\sqrt{3}}{6}\right)$
 (3) $\frac{\pi}{8} \left(1 - \frac{\sqrt{3}}{6}\right)$ (4) $\frac{\pi}{4} \left(1 - \frac{\sqrt{3}}{2}\right)$

Official Ans. by NTA (1)

Sol. $I = \int_0^1 \frac{\sqrt{x}}{(1+x)(1+3x)(3+x)} dx$

Let $x = t^2 \Rightarrow dx = 2t dt$

$$I = \int_0^1 \frac{t(2t)}{(t^2+1)(1+3t^2)(3+t^2)} dt$$

$$I = \int_0^1 \frac{(3t^2+1) - (t^2+1)}{(3t^2+1)(t^2+1)(3+t^2)} dt$$

$$I = \int_0^1 \frac{dt}{(t^2+1)(3+t^2)} - \int_0^1 \frac{dt}{(1+3t^2)(3+t^2)}$$

$$= \frac{1}{2} \int_0^1 \frac{(3+t^2) - (t^2+1)}{(t^2+1)(3+t^2)} dt + \frac{1}{8} \int_0^1 \frac{(1+3t^2) - 3(3+t^2)}{(1+3t^2)(3+t^2)} dt$$

$$= \frac{1}{2} \int_0^1 \frac{dt}{1+t^2} - \frac{1}{2} \int_0^1 \frac{dt}{t^2+3} + \frac{1}{8} \int_0^1 \frac{dt}{t^2+3} - \frac{3}{8} \int_0^1 \frac{dt}{(1+3t^2)}$$

$$= \frac{1}{2} \int_0^1 \frac{dt}{t^2+1} - \frac{3}{8} \int_0^1 \frac{dt}{t^2+3} - \frac{3}{8} \int_0^1 \frac{dt}{1+3t^2}$$

$$= \frac{1}{2} (\tan^{-1}(t))_0^1 - \frac{3}{8\sqrt{3}} \left(\tan^{-1} \left(\frac{t}{\sqrt{3}} \right) \right)_0^1$$

$$- \frac{3}{8\sqrt{3}} (\tan^{-1}(\sqrt{3}t))_0^1$$

$$= \frac{1}{2} \left(\frac{\pi}{4} \right) - \frac{\sqrt{3}}{8} \left(\frac{\pi}{6} \right) - \frac{\sqrt{3}}{8} \left(\frac{\pi}{3} \right)$$

$$= \frac{\pi}{8} - \frac{\sqrt{3}}{16} \pi$$

$$= \frac{\pi}{8} \left(1 - \frac{\sqrt{3}}{2} \right)$$

20. If $\lim_{x \rightarrow \infty} (\sqrt{x^2 - x + 1} - ax) = b$, then the ordered pair (a, b) is:

(1) $\left(1, \frac{1}{2}\right)$ (2) $\left(1, -\frac{1}{2}\right)$

(3) $\left(-1, \frac{1}{2}\right)$ (4) $\left(-1, -\frac{1}{2}\right)$

Official Ans. by NTA (2)



Sol. (2)
 $\lim_{x \rightarrow \infty} (\sqrt{x^2 - x + 1}) - ax = b \quad (\infty - \infty)$
 $\Rightarrow a > 0$

Now, $\lim_{x \rightarrow \infty} \frac{(x^2 - x + 1 - a^2 x^2)}{\sqrt{x^2 - x + 1} + ax} = b$

$\Rightarrow \lim_{x \rightarrow \infty} \frac{(1 - a^2)x^2 - x + 1}{\sqrt{x^2 - x + 1} + ax} = b$

$\Rightarrow \lim_{x \rightarrow \infty} \frac{(1 - a^2)x^2 - x + 1}{x \left(\sqrt{1 - \frac{1}{x} + \frac{1}{x^2}} + a \right)} = b$

$\Rightarrow 1 - a^2 = 0 \Rightarrow a = 1$

Now, $\lim_{x \rightarrow \infty} \frac{-x + 1}{x \left(\sqrt{1 - \frac{1}{x} + \frac{1}{x^2}} + a \right)} = b$

$\Rightarrow \frac{-1}{1 + a} = b \Rightarrow b = -\frac{1}{2}$

$(a, b) = \left(1, -\frac{1}{2} \right)$

SECTION-B

1. Let S be the sum of all solutions (in radians) of the equation $\sin^4 \theta + \cos^4 \theta - \sin \theta \cos \theta = 0$ in $[0, 4\pi]$.

Then $\frac{8S}{\pi}$ is equal to _____.

Official Ans. by NTA (56)

Sol. Given equation

$\sin^4 \theta + \cos^4 \theta - \sin \theta \cos \theta = 0$

$\Rightarrow 1 - \sin^2 \theta \cos^2 \theta - \sin \theta \cos \theta = 0$

$\Rightarrow 2 - (\sin 2\theta)^2 - \sin 2\theta = 0$

$\Rightarrow (\sin 2\theta)^2 + (\sin 2\theta) - 2 = 0$

$\Rightarrow (\sin 2\theta + 2)(\sin 2\theta - 1) = 0$

$\Rightarrow \sin 2\theta = 1$ or $\boxed{\sin 2\theta = -2}$
(not possible)

$\Rightarrow 2\theta = \frac{\pi}{2}, \frac{5\pi}{2}, \frac{9\pi}{2}, \frac{13\pi}{2}$

$\Rightarrow \theta = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}, \frac{13\pi}{4}$

$\Rightarrow S = \frac{\pi}{4} + \frac{5\pi}{4} + \frac{9\pi}{4} + \frac{13\pi}{4} = 7\pi$

$\Rightarrow \frac{8S}{\pi} = \frac{8 \times 7\pi}{\pi} = 56.00$

2. Let S be the mirror image of the point Q(1, 3, 4) with respect to the plane $2x - y + z + 3 = 0$ and let R(3, 5, γ) be a point of this plane. Then the square of the length of the line segment SR is _____.

Official Ans. by NTA (72)

Sol. Since R(3, 5, γ) lies on the plane $2x - y + z + 3 = 0$.

Therefore, $6 - 5 + \gamma + 3 = 0$

$\Rightarrow \gamma = -4$

Now,

dir's of line QS

are 2, -1, 1

equation of line QS is

$\frac{x-1}{2} = \frac{y-3}{-1} = \frac{z-4}{1} = \lambda$ (say)

$\Rightarrow F(2\lambda + 1, -\lambda + 3, \lambda + 4)$

F lies in the plane

$\Rightarrow 2(2\lambda + 1) - (-\lambda + 3) + (\lambda + 4) + 3 = 0$

$\Rightarrow 4\lambda + 2 + \lambda - 3 + \lambda + 7 = 0$

$\Rightarrow 6\lambda + 6 = 0 \Rightarrow \lambda = -1$.

$\Rightarrow F(-1, 4, 3)$

Since, F is mid-point of QS.

Therefore, co-ordinates of S are (-3, 5, 2).

So, $SR = \sqrt{36 + 0 + 36} = \sqrt{72}$


$SR^2 = 72$.

3. The probability distribution of random variable X is given by:

X	1	2	3	4	5
P(X)	K	2K	2K	3K	K

Let $p = P(1 < X < 4 \mid X < 3)$. If $5p = \lambda K$, then λ is equal to _____.

Official Ans. by NTA (30)





$$\sum P(X) = 1 \Rightarrow k + 2k + 2k + 3k + k = 1$$

$$\Rightarrow k = \frac{1}{9}$$

$$\text{Now, } p = P\left(\frac{kX < 4}{X < 3}\right) = \frac{P(X=2)}{P(X < 3)} = \frac{\frac{2k}{9k}}{\frac{k}{9k} + \frac{2k}{9k}} = \frac{2}{3}$$

$$\Rightarrow p = \frac{2}{3}$$

$$\text{Now, } 5p = \lambda k$$

$$\Rightarrow (5)\left(\frac{2}{3}\right) = \lambda(1/9)$$

$$\Rightarrow \lambda = 30$$

4. Let z_1 and z_2 be two complex numbers such that $\arg(z_1 - z_2) = \frac{\pi}{4}$ and z_1, z_2 satisfy the equation $|z - 3| = \text{Re}(z)$. Then the imaginary part of $z_1 + z_2$ is equal to _____.

Official Ans. by NTA (6)

Sol. $|z - 3| = \text{Re}(z)$

$$\text{let } Z = x + iy$$

$$\Rightarrow (x - 3)^2 + y^2 = x^2$$

$$\Rightarrow x^2 + 9 - 6x + y^2 = x^2$$


$$\Rightarrow y^2 = 6x - 9$$

$$\Rightarrow y^2 = 6\left(x - \frac{3}{2}\right)$$

$\Rightarrow z_1$ and z_2 lie on the parabola mentioned in eq.(1)

$$\arg(z_1 - z_2) = \frac{\pi}{4}$$

\Rightarrow Slope of PQ = 1.



$$\text{Let } P\left(\frac{3}{2} + \frac{3}{2}t_1^2, 3t_1\right) \text{ and } Q\left(\frac{3}{2} + \frac{3}{2}t_2^2, 3t_2\right)$$

$$\text{Slope of PQ} = \frac{3(t_2 - t_1)}{\frac{3}{2}(t_2^2 - t_1^2)} = 1$$

$$\Rightarrow \frac{2}{t_2 + t_1} = 1$$

$$\Rightarrow t_2 + t_1 = 2$$

$$\text{Im}(z_1 + z_2) = 3t_1 + 3t_2 = 3(t_1 + t_2) = 3(2)$$

Ans. 6.00

Aliter :

$$\text{Let } z_1 = x_1 + iy_1; z_2 = x_2 + iy_2$$

$$z_1 - z_2 = (x_1 - x_2) + i(y_1 - y_2)$$

$$\therefore \arg(z_1 - z_2) = \frac{\pi}{4} \Rightarrow \tan^{-1}\left(\frac{y_1 - y_2}{x_1 - x_2}\right) = \frac{\pi}{4}$$

$$y_1 - y_2 = x_1 - x_2 \text{ (1)}$$

$$|z_1 - 3| = \text{Re}(z_1) \Rightarrow (x_1 - 3)^2 + y_1^2 = x_1^2 \text{ (2)}$$

$$|z_2 - 3| = \text{Re}(z_2) \Rightarrow (x_2 - 3)^2 + y_2^2 = x_2^2 \text{ (2)}$$

sub (2) & (3)

$$(x_1 - 3)^2 - (x_2 - 3)^2 + y_1^2 - y_2^2 = x_1^2 - x_2^2$$

$$(x_1 - x_2)(x_1 + x_2 - 6) + (y_1 - y_2)(y_1 + y_2)$$

$$= (x_1 - x_2)(x_1 + x_2)$$

$$x_1 + x_2 - 6 + y_1 + y_2 = x_1 + x_2 \Rightarrow y_1 + y_2 = 6.$$

5. Let $S = \{1, 2, 3, 4, 5, 6, 9\}$. Then the number of elements in the set $T = \{A \subseteq S : A \neq \phi \text{ and the sum of all the elements of } A \text{ is not a multiple of } 3\}$ is _____.

Official Ans. by NTA (80)

Sol. $3n$ type $\rightarrow 3, 6, 9 = P$

$3n - 1$ type $\rightarrow 2, 5 = Q$

$3n - 2$ type $\rightarrow 1, 4 = R$

number of subset of S containing one element which are not divisible by 3 = ${}^2C_1 + {}^2C_1 = 4$

number of subset of S containing two numbers whose some is not divisible by 3

$$= {}^3C_1 \times {}^2C_1 + {}^3C_1 \times {}^2C_1 + {}^2C_2 + {}^2C_2 = 14$$

number of subsets containing 3 elements whose sum is not divisible by 3

$$= {}^3C_2 \times {}^4C_1 + ({}^2C_2 \times {}^2C_1) \times 2 + {}^3C_1 ({}^2C_2 + {}^2C_2) = 22$$

number of subsets containing 4 elements whose sum is not divisible by 3

$$= {}^3C_3 \times {}^4C_1 + {}^3C_2 ({}^2C_2 + {}^2C_2) + ({}^3C_1 {}^2C_1 \times {}^2C_2) \times 2$$

$$= 4 + 6 + 12 = 22.$$

number of subsets of S containing 5 elements whose sum is not divisible by 3.

$$= {}^3C_3 ({}^2C_2 + {}^2C_2) + ({}^3C_2 {}^2C_1 \times {}^2C_2) \times 2 = 2 + 12 = 14$$

number of subsets of S containing 6 elements whose sum is not divisible by 3 = 4

\Rightarrow Total subsets of Set A whose sum of digits is not divisible by 3 = $4 + 14 + 22 + 22 + 14 + 4 = 80.$



6. Let A (secθ, 2tanθ) and B (secφ, 2tanφ), where θ + φ = π/2, be two points on the hyperbola 2x² - y² = 2. If (α, β) is the point of the intersection of the normals to the hyperbola at A and B, then (2β)² is equal to _____.

Official Ans. by NTA (36)

ALLEN Ans. (Bonus)

Sol. Since, point A (sec θ, 2 tan θ) lies on the hyperbola
2x² - y² = 2

$$\text{Therefore, } 2 \sec^2 \theta - 4 \tan^2 \theta = 2$$

$$\Rightarrow 2 + 2 \tan^2 \theta - 4 \tan^2 \theta = 2$$

$$\Rightarrow \tan \theta = 0 \Rightarrow \theta = 0$$

Similarly, for point B, we will get φ = 0.

$$\text{but according to question } \theta + \phi = \frac{\pi}{2}$$


which is not possible.

Hence it must be a 'BONUS'.

7. Two circles each of radius 5 units touch each other at the point (1, 2). If the equation of their common tangent is 4x + 3y = 10, and C₁(α, β) and C₂(γ, δ), C₁ ≠ C₂ are their centres, then |(α + β) (γ + δ)| is equal to _____.

Official Ans. by NTA (40)

Sol. Slope of line joining centres of circles = $\frac{4}{3} = \tan \theta$



$$\Rightarrow \cos \theta = \frac{3}{5}, \sin \theta = \frac{4}{5}$$

Now using parametric form

$$\frac{x-1}{\cos \theta} = \frac{y-2}{\sin \theta} = \pm 5$$

$$\oplus (x, y) = (1 + 5 \cos \theta, 2 + 5 \sin \theta)$$

$$(\alpha, \beta) = (4, 6)$$

$$\ominus (x, y) = (\gamma, \delta) = (1 - 5 \cos \theta, 2 - 5 \sin \theta)$$

$$(\gamma, \delta) = (-2, -2)$$

$$\Rightarrow |(\alpha + \beta) (\gamma + \delta)| = |10 \times -4| = 40$$

8. $3 \times 7^{22} + 2 \times 10^{22} - 44$ when divided by 18 leaves the remainder _____.

Official Ans. by NTA (15)

Sol. $3(1 + 6)^{22} + 2 \cdot (1 + 9)^{22} - 44 = (3 + 2 - 44) = 18 \cdot I$

$$= -39 + 18 \cdot I$$

$$= (54 - 39) + 18(I - 3)$$

$$= 15 + 18 I_1$$

⇒ Remainder = 15.

9. An online exam is attempted by 50 candidates out of which 20 are boys. The average marks obtained by boys is 12 with a variance 2. The variance of marks obtained by 30 girls is also 2. The average marks of all 50 candidates is 15. If μ is the average marks of girls and σ² is the variance of marks of 50 candidates, then μ + σ² is equal to _____.

Official Ans. by NTA (25)

Sol. σ_b² = 2 (variance of boys) n₁ = no. of boys

$$\bar{x}_b = 12$$

$$n_2 = \text{no. of girls}$$

$$\sigma_g^2 = 2$$

$$\bar{x}_g = \frac{50 \times 15 - 12 \times \sigma_b}{30} = \frac{750 - 12 \times 20}{30} = 17 = \mu$$

variance of combined series

$$\sigma^2 = \frac{n_1 \sigma_b^2 + n_2 \sigma_g^2}{n_1 + n_2} + \frac{n_1 \cdot n_2}{(n_1 + n_2)^2} (\bar{x}_b - \bar{x}_g)^2$$

$$\sigma^2 = \frac{20 \times 2 + 30 \times 2}{20 + 30} + \frac{20 \times 30}{(20 + 30)^2} (12 - 17)^2$$

$$\sigma^2 = 8.$$

$$\Rightarrow \mu + \sigma^2 = 17 + 8 = 25$$



10. If $\int \frac{2e^x + 3e^{-x}}{4e^x + 7e^{-x}} dx = \frac{1}{14}(ux + v \log_e(4e^x + 7e^{-x})) + C$,

where C is a constant of integration, then u + v is equal to _____.

Official Ans. by NTA (7)

Sol. $\int \frac{2e^x}{4e^x + 7e^{-x}} dx + 3 \int \frac{e^{-x}}{4e^x + 7e^{-x}} dx$

$= \int \frac{2e^{2x}}{4e^{2x} + 7} dx + 3 \int \frac{e^{-2x}}{4 + 7e^{-2x}} dx$

Let $4e^{2x} + 7 = T$ Let $4 + 7e^{-2x} = t$

$8e^{2x} dx = dT$ $-14e^{-2x} dx = dt$

$2e^{2x} dx = \frac{dT}{4}$ $e^{-2x} dx = -\frac{dt}{14}$

$\int \frac{dT}{4T} - \frac{3}{14} \int \frac{dt}{t}$

$= \frac{1}{4} \log T - \frac{3}{14} \log t + C$

$= \frac{1}{4} \log(4e^{2x} + 7) - \frac{3}{14} \log(4 + 7e^{-2x}) + C$

$= \frac{1}{14} \left[\frac{1}{2} \log(4e^x + 7e^{-x}) + \frac{13}{2} x \right] + C$

$u = \frac{13}{2}, v = \frac{1}{2} \Rightarrow u + v = 7$

Aliter :

$2e^x + 3e^{-x} = A(4e^x + 7e^{-x}) + B(4e^x - 7e^{-x}) + \lambda$

$2 = 4A + 4B \quad ; \quad 3 = 7A - 7B \quad ; \quad \lambda = 0$

$A + B = \frac{1}{2}$

$A - B = \frac{3}{7}$

$A = \frac{1}{2} \left(\frac{1}{2} + \frac{3}{7} \right) = \frac{7+6}{28} = \frac{13}{28}$

$B = A - \frac{3}{7} = \frac{13}{28} - \frac{3}{7} = \frac{13-12}{28} = \frac{1}{28}$

$\int \frac{13}{28} dx + \frac{1}{28} \int \frac{4e^x - 7e^{-x}}{4e^x + 7e^{-x}} dx$

$\frac{13}{28} x + \frac{1}{28} \ln |4e^x + 7e^{-x}| + C$

$u = \frac{13}{2}; v = \frac{1}{2}$

$\Rightarrow u + v = 7$