

Class XII : Maths  
Chapter 5 : Continuity And Differentiability

Questions and Solutions | Exercise 5.1 - NCERT Books

**Question 1:**

Prove that the function  $f(x) = 5x - 3$  is continuous at  $x = 0$ , at  $x = -3$  and at  $x = 5$ .

Answer

The given function is  $f(x) = 5x - 3$

At  $x = 0$ ,  $f(0) = 5 \times 0 - 3 = 3$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} (5x - 3) = 5 \times 0 - 3 = -3$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

At  $x = -3$ ,  $f(-3) = 5 \times (-3) - 3 = -18$

$$\lim_{x \rightarrow -3} f(x) = \lim_{x \rightarrow -3} (5x - 3) = 5 \times (-3) - 3 = -18$$

$$\therefore \lim_{x \rightarrow -3} f(x) = f(-3)$$

Therefore,  $f$  is continuous at  $x = -3$

At  $x = 5$ ,  $f(5) = 5 \times 5 - 3 = 25 - 3 = 22$

$$\lim_{x \rightarrow 5} f(x) = \lim_{x \rightarrow 5} (5x - 3) = 5 \times 5 - 3 = 22$$

$$\therefore \lim_{x \rightarrow 5} f(x) = f(5)$$

Therefore,  $f$  is continuous at  $x = 5$

**Question 2:**

Examine the continuity of the function  $f(x) = 2x^2 - 1$  at  $x = 3$ .

Answer

The given function is  $f(x) = 2x^2 - 1$

At  $x = 3$ ,  $f(3) = 2 \times 3^2 - 1 = 17$

$$\lim_{x \rightarrow 3} f(x) = \lim_{x \rightarrow 3} (2x^2 - 1) = 2 \times 3^2 - 1 = 17$$

$$\therefore \lim_{x \rightarrow 3} f(x) = f(3)$$

Thus,  $f$  is continuous at  $x = 3$

**Question 3:**

Examine the following functions for continuity.

$$(a) f(x) = x - 5 \quad (b) f(x) = \frac{1}{x - 5}, x \neq 5$$

$$(c) f(x) = \frac{x^2 - 25}{x + 5}, x \neq -5 \quad (d) f(x) = |x - 5|$$

Answer

(a) The given function is  $f(x) = x - 5$

It is evident that  $f$  is defined at every real number  $k$  and its value at  $k$  is  $k - 5$ .

It is also observed that,  $\lim_{x \rightarrow k} f(x) = \lim_{x \rightarrow k} (x - 5) = k - 5 = f(k)$

$$\therefore \lim_{x \rightarrow k} f(x) = f(k)$$

Hence,  $f$  is continuous at every real number and therefore, it is a continuous function.

(b) The given function is  $f(x) = \frac{1}{x - 5}, x \neq 5$

For any real number  $k \neq 5$ , we obtain

$$\lim_{x \rightarrow k} f(x) = \lim_{x \rightarrow k} \frac{1}{x - 5} = \frac{1}{k - 5}$$

$$\text{Also, } f(k) = \frac{1}{k - 5} \quad (\text{As } k \neq 5)$$

$$\therefore \lim_{x \rightarrow k} f(x) = f(k)$$

Hence,  $f$  is continuous at every point in the domain of  $f$  and therefore, it is a continuous function.

(c) The given function is  $f(x) = \frac{x^2 - 25}{x + 5}, x \neq -5$

For any real number  $c \neq -5$ , we obtain

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \frac{x^2 - 25}{x + 5} = \lim_{x \rightarrow c} \frac{(x+5)(x-5)}{x+5} = \lim_{x \rightarrow c} (x-5) = (c-5)$$

$$\text{Also, } f(c) = \frac{(c+5)(c-5)}{c+5} = (c-5) \quad (\text{as } c \neq -5)$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Hence,  $f$  is continuous at every point in the domain of  $f$  and therefore, it is a continuous function.

$$f(x) = |x-5| = \begin{cases} 5-x, & \text{if } x < 5 \\ x-5, & \text{if } x \geq 5 \end{cases}$$

(d) The given function is

This function  $f$  is defined at all points of the real line.

Let  $c$  be a point on a real line. Then,  $c < 5$  or  $c = 5$  or  $c > 5$

Case I:  $c < 5$

Then,  $f(c) = 5 - c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (5 - x) = 5 - c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all real numbers less than 5.

Case II :  $c = 5$

Then,  $f(c) = f(5) = (5-5) = 0$

$$\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5} (5 - x) = (5-5) = 0$$

$$\lim_{x \rightarrow 5^+} f(x) = \lim_{x \rightarrow 5} (x - 5) = 0$$

$$\therefore \lim_{x \rightarrow c^-} f(x) = \lim_{x \rightarrow c^+} f(x) = f(c)$$

Therefore,  $f$  is continuous at  $x = 5$

Case III:  $c > 5$

Then,  $f(c) = f(5) = c - 5$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x - 5) = c - 5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all real numbers greater than 5.

Hence,  $f$  is continuous at every real number and therefore, it is a continuous function.

**Question 4:**

Prove that the function  $f(x) = x^n$  is continuous at  $x = n$ , where  $n$  is a positive integer.

Answer

The given function is  $f(x) = x^n$

It is evident that  $f$  is defined at all positive integers,  $n$ , and its value at  $n$  is  $n^n$ .

$$\text{Then, } \lim_{x \rightarrow n} f(n) = \lim_{x \rightarrow n} (x^n) = n^n$$

$$\therefore \lim_{x \rightarrow n} f(x) = f(n)$$

Therefore,  $f$  is continuous at  $n$ , where  $n$  is a positive integer.

**Question 5:**

Is the function  $f$  defined by

$$f(x) = \begin{cases} x, & \text{if } x \leq 1 \\ 5, & \text{if } x > 1 \end{cases}$$

continuous at  $x = 0$ ? At  $x = 1$ ? At  $x = 2$ ?

Answer

$$f(x) = \begin{cases} x, & \text{if } x \leq 1 \\ 5, & \text{if } x > 1 \end{cases}$$

The given function  $f$  is

At  $x = 0$ ,

It is evident that  $f$  is defined at 0 and its value at 0 is 0.

$$\text{Then, } \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} x = 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

At  $x = 1$ ,

$f$  is defined at 1 and its value at 1 is 1.

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} x = 1$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (5) = 5$$

$$\therefore \lim_{x \rightarrow 1^-} f(x) \neq \lim_{x \rightarrow 1^+} f(x)$$

Therefore,  $f$  is not continuous at  $x = 1$

At  $x = 2$ ,

$f$  is defined at 2 and its value at 2 is 5.

$$\text{Then, } \lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} (5) = 5$$

$$\therefore \lim_{x \rightarrow 2} f(x) = f(2)$$

Therefore,  $f$  is continuous at  $x = 2$

#### Question 6:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} 2x+3, & \text{if } x \leq 2 \\ 2x-3, & \text{if } x > 2 \end{cases}$$

Answer

$$\text{The given function } f \text{ is } f(x) = \begin{cases} 2x+3, & \text{if } x \leq 2 \\ 2x-3, & \text{if } x > 2 \end{cases}$$

It is evident that the given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line. Then, three cases arise.

(i)  $c < 2$

(ii)  $c > 2$

(iii)  $c = 2$

Case (i)  $c < 2$

$$\text{Then, } f(c) = 2c+3$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x+3) = 2c+3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 2$

Case (ii)  $c > 2$

Then,  $f(c) = 2c - 3$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x - 3) = 2c - 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 2$

Case (iii)  $c = 2$

Then, the left hand limit of  $f$  at  $x = 2$  is,

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (2x + 3) = 2 \times 2 + 3 = 7$$

The right hand limit of  $f$  at  $x = 2$  is,

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (2x - 3) = 2 \times 2 - 3 = 1$$

It is observed that the left and right hand limit of  $f$  at  $x = 2$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 2$

Hence,  $x = 2$  is the only point of discontinuity of  $f$ .

#### Question 7:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} |x| + 3, & \text{if } x \leq -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x + 2, & \text{if } x \geq 3 \end{cases}$$

Answer

$$f(x) = \begin{cases} |x| + 3 = -x + 3, & \text{if } x \leq -3 \\ -2x, & \text{if } -3 < x < 3 \\ 6x + 2, & \text{if } x \geq 3 \end{cases}$$

The given function  $f$  is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < -3$ , then  $f(c) = -c + 3$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-x + 3) = -c + 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < -3$

Case II:

If  $c = -3$ , then  $f(-3) = -(-3) + 3 = 6$

$$\lim_{x \rightarrow -3^-} f(x) = \lim_{x \rightarrow -3^-} (-x + 3) = -(-3) + 3 = 6$$

$$\lim_{x \rightarrow -3^+} f(x) = \lim_{x \rightarrow -3^+} (-2x) = -2 \times (-3) = 6$$

$$\therefore \lim_{x \rightarrow -3} f(x) = f(-3)$$

Therefore,  $f$  is continuous at  $x = -3$

Case III:

If  $-3 < c < 3$ , then  $f(c) = -2c$  and  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-2x) = -2c$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous in  $(-3, 3)$ .

Case IV:

If  $c = 3$ , then the left hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (-2x) = -2 \times 3 = -6$$

The right hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (6x + 2) = 6 \times 3 + 2 = 20$$

It is observed that the left and right hand limit of  $f$  at  $x = 3$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 3$

Case V:

If  $c > 3$ , then  $f(c) = 6c + 2$  and  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (6x + 2) = 6c + 2$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 3$

Hence,  $x = 3$  is the only point of discontinuity of  $f$ .

### Question 8:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} \frac{|x|}{x} & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

Answer

$$f(x) = \begin{cases} \frac{|x|}{x} & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

The given function  $f$  is

It is known that,  $x < 0 \Rightarrow |x| = -x$  and  $x > 0 \Rightarrow |x| = x$

Therefore, the given function can be rewritten as

$$f(x) = \begin{cases} \frac{|x|}{x} = \frac{-x}{x} = -1 & \text{if } x < 0 \\ 0, & \text{if } x = 0 \\ \frac{|x|}{x} = \frac{x}{x} = 1, & \text{if } x > 0 \end{cases}$$

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < 0$ , then  $f(c) = -1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-1) = -1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x < 0$

Case II:

If  $c = 0$ , then the left hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (-1) = -1$$

The right hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (1) = 1$$

It is observed that the left and right hand limit of  $f$  at  $x = 0$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 0$

Case III:



If  $c > 0$ , then  $f(c) = 1$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (1) = 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 0$

Hence,  $x = 0$  is the only point of discontinuity of  $f$ .

### Question 9:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ -1, & \text{if } x \geq 0 \end{cases}$$

Answer

$$f(x) = \begin{cases} \frac{x}{|x|}, & \text{if } x < 0 \\ -1, & \text{if } x \geq 0 \end{cases}$$

The given function  $f$  is

It is known that,  $x < 0 \Rightarrow |x| = -x$

Therefore, the given function can be rewritten as

$$f(x) = \begin{cases} \frac{x}{|x|} = \frac{x}{-x} = -1, & \text{if } x < 0 \\ -1, & \text{if } x \geq 0 \end{cases}$$

$$\Rightarrow f(x) = -1 \text{ for all } x \in \mathbf{R}$$

Let  $c$  be any real number. Then,  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-1) = -1$

Also,  $f(c) = -1 = \lim_{x \rightarrow c} f(x)$

Therefore, the given function is a continuous function.

Hence, the given function has no point of discontinuity.

### Question 10:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} x+1, & \text{if } x \geq 1 \\ x^2+1, & \text{if } x < 1 \end{cases}$$

Answer

$$f(x) = \begin{cases} x+1, & \text{if } x \geq 1 \\ x^2+1, & \text{if } x < 1 \end{cases}$$

The given function  $f$  is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

$$\text{If } c < 1, \text{ then } f(c) = c^2 + 1 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^2 + 1) = c^2 + 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 1$

Case II:

$$\text{If } c = 1, \text{ then } f(c) = f(1) = 1 + 1 = 2$$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^2 + 1) = 1^2 + 1 = 2$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x + 1) = 1 + 1 = 2$$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(1)$$

Therefore,  $f$  is continuous at  $x = 1$

Case III:

$$\text{If } c > 1, \text{ then } f(c) = c + 1$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x + 1) = c + 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$

Hence, the given function  $f$  has no point of discontinuity.

#### Question 11:

Find all points of discontinuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} x^3 - 3, & \text{if } x \leq 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}$$

Answer

$$f(x) = \begin{cases} x^3 - 3, & \text{if } x \leq 2 \\ x^2 + 1, & \text{if } x > 2 \end{cases}$$

The given function  $f$  is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

$$\text{If } c < 2, \text{ then } f(c) = c^3 - 3 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^3 - 3) = c^3 - 3$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 2$

Case II:

$$\text{If } c = 2, \text{ then } f(c) = f(2) = 2^3 - 3 = 5$$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} (x^3 - 3) = 2^3 - 3 = 5$$

$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} (x^2 + 1) = 2^2 + 1 = 5$$

$$\therefore \lim_{x \rightarrow 2} f(x) = f(2)$$

Therefore,  $f$  is continuous at  $x = 2$

Case III:

$$\text{If } c > 2, \text{ then } f(c) = c^2 + 1$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^2 + 1) = c^2 + 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 2$

Thus, the given function  $f$  is continuous at every point on the real line.

Hence,  $f$  has no point of discontinuity.

### Question 12:

Find all points of discontinuity of  $f$ , where  $f$  is defined by



$$f(x) = \begin{cases} x^{10} - 1, & \text{if } x \leq 1 \\ x^2, & \text{if } x > 1 \end{cases}$$

Answer

$$f(x) = \begin{cases} x^{10} - 1, & \text{if } x \leq 1 \\ x^2, & \text{if } x > 1 \end{cases}$$

The given function  $f$  is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

$$\text{If } c < 1, \text{ then } f(c) = c^{10} - 1 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^{10} - 1) = c^{10} - 1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 1$

Case II:

If  $c = 1$ , then the left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x^{10} - 1) = 1^{10} - 1 = 1 - 1 = 0$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x^2) = 1^2 = 1$$

It is observed that the left and right hand limit of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$

Case III:

$$\text{If } c > 1, \text{ then } f(c) = c^2$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x^2) = c^2$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$

Thus, from the above observation, it can be concluded that  $x = 1$  is the only point of discontinuity of  $f$ .

### Question 13:

Is the function defined by

$$f(x) = \begin{cases} x+5, & \text{if } x \leq 1 \\ x-5, & \text{if } x > 1 \end{cases}$$

a continuous function?

Answer

$$f(x) = \begin{cases} x+5, & \text{if } x \leq 1 \\ x-5, & \text{if } x > 1 \end{cases}$$

The given function is

The given function  $f$  is defined at all the points of the real line.

Let  $c$  be a point on the real line.

Case I:

$$\text{If } c < 1, \text{ then } f(c) = c+5 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x+5) = c+5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 1$

Case II:

$$\text{If } c = 1, \text{ then } f(1) = 1+5 = 6$$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (x+5) = 1+5 = 6$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (x-5) = 1-5 = -4$$

It is observed that the left and right hand limit of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$

Case III:

$$\text{If } c > 1, \text{ then } f(c) = c-5 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x-5) = c-5$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$

Thus, from the above observation, it can be concluded that  $x = 1$  is the only point of discontinuity of  $f$ .

## Question 14:

Discuss the continuity of the function  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$

Answer

$$f(x) = \begin{cases} 3, & \text{if } 0 \leq x \leq 1 \\ 4, & \text{if } 1 < x < 3 \\ 5, & \text{if } 3 \leq x \leq 10 \end{cases}$$

The given function is

The given function is defined at all points of the interval  $[0, 10]$ .

Let  $c$  be a point in the interval  $[0, 10]$ .

Case I:

If  $0 \leq c < 1$ , then  $f(c) = 3$  and  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (3) = 3$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous in the interval  $[0, 1)$ .

Case II:

If  $c = 1$ , then  $f(1) = 3$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (3) = 3$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (4) = 4$$

It is observed that the left and right hand limits of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$

Case III:

If  $1 < c < 3$ , then  $f(c) = 4$  and  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (4) = 4$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points of the interval  $(1, 3)$ .

Case IV:

If  $c = 3$ , then  $f(c) = 5$

The left hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (4) = 4$$

The right hand limit of  $f$  at  $x = 3$  is,

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (5) = 5$$

It is observed that the left and right hand limits of  $f$  at  $x = 3$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 3$

Case V:

$$\text{If } 3 < c \leq 10, \text{ then } f(c) = 5 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (5) = 5$$

$$\lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points of the interval  $(3, 10]$ .

Hence,  $f$  is not continuous at  $x = 1$  and  $x = 3$

#### Question 15:

Discuss the continuity of the function  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \leq x \leq 1 \\ 4x, & \text{if } x > 1 \end{cases}$$

Answer

$$f(x) = \begin{cases} 2x, & \text{if } x < 0 \\ 0, & \text{if } 0 \leq x \leq 1 \\ 4x, & \text{if } x > 1 \end{cases}$$

The given function is

The given function is defined at all points of the real line.

Let  $c$  be a point on the real line.

Case I:

$$\text{If } c < 0, \text{ then } f(c) = 2c$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x) = 2c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 0$

Case II:

$$\text{If } c = 0, \text{ then } f(c) = f(0) = 0$$

The left hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} (2x) = 2 \times 0 = 0$$

The right hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (0) = 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

Case III:

$$\text{If } 0 < c < 1, \text{ then } f(x) = 0 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (0) = 0$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points of the interval  $(0, 1)$ .

Case IV:

$$\text{If } c = 1, \text{ then } f(c) = f(1) = 0$$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (0) = 0$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} (4x) = 4 \times 1 = 4$$

It is observed that the left and right hand limits of  $f$  at  $x = 1$  do not coincide.

Therefore,  $f$  is not continuous at  $x = 1$

Case V:

$$\text{If } c < 1, \text{ then } f(c) = 4c \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (4x) = 4c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$

Hence,  $f$  is not continuous only at  $x = 1$

#### Question 16:

Discuss the continuity of the function  $f$ , where  $f$  is defined by



$$f(x) = \begin{cases} -2, & \text{if } x \leq -1 \\ 2x, & \text{if } -1 < x \leq 1 \\ 2, & \text{if } x > 1 \end{cases}$$

Answer

$$f(x) = \begin{cases} -2, & \text{if } x \leq -1 \\ 2x, & \text{if } -1 < x \leq 1 \\ 2, & \text{if } x > 1 \end{cases}$$

The given function  $f$  is

The given function is defined at all points of the real line.

Let  $c$  be a point on the real line.

Case I:

If  $c < -1$ , then  $f(c) = -2$  and  $\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (-2) = -2$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < -1$

Case II:

If  $c = -1$ , then  $f(c) = f(-1) = -2$

The left hand limit of  $f$  at  $x = -1$  is,

$$\lim_{x \rightarrow -1^-} f(x) = \lim_{x \rightarrow -1^-} (-2) = -2$$

The right hand limit of  $f$  at  $x = -1$  is,

$$\lim_{x \rightarrow -1^+} f(x) = \lim_{x \rightarrow -1^+} (2x) = 2 \times (-1) = -2$$

$$\therefore \lim_{x \rightarrow -1} f(x) = f(-1)$$

Therefore,  $f$  is continuous at  $x = -1$

Case III:

If  $-1 < c < 1$ , then  $f(c) = 2c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2x) = 2c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points of the interval  $(-1, 1)$ .

Case IV:

$$\text{If } c = 1, \text{ then } f(c) = f(1) = 2 \times 1 = 2$$

The left hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^-} f(x) = \lim_{x \rightarrow 1^-} (2x) = 2 \times 1 = 2$$

The right hand limit of  $f$  at  $x = 1$  is,

$$\lim_{x \rightarrow 1^+} f(x) = \lim_{x \rightarrow 1^+} 2 = 2$$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(c)$$

Therefore,  $f$  is continuous at  $x = 2$

Case V:

$$\text{If } c > 1, \text{ then } f(c) = 2 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (2) = 2$$

$$\lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 1$

Thus, from the above observations, it can be concluded that  $f$  is continuous at all points of the real line.

#### Question 17:

Find the relationship between  $a$  and  $b$  so that the function  $f$  defined by

$$f(x) = \begin{cases} ax+1, & \text{if } x \leq 3 \\ bx+3, & \text{if } x > 3 \end{cases}$$

is continuous at  $x = 3$ .

Answer

$$f(x) = \begin{cases} ax+1, & \text{if } x \leq 3 \\ bx+3, & \text{if } x > 3 \end{cases}$$

The given function  $f$  is

If  $f$  is continuous at  $x = 3$ , then

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^+} f(x) = f(3) \quad \dots(1)$$

Also,

$$\lim_{x \rightarrow 3^-} f(x) = \lim_{x \rightarrow 3^-} (ax+1) = 3a+1$$

$$\lim_{x \rightarrow 3^+} f(x) = \lim_{x \rightarrow 3^+} (bx+3) = 3b+3$$

$$f(3) = 3a+1$$

Therefore, from (1), we obtain

$$3a+1 = 3b+3 = 3a+1$$

$$\Rightarrow 3a+1 = 3b+3$$

$$\Rightarrow 3a = 3b+2$$

$$\Rightarrow a = b + \frac{2}{3}$$

Therefore, the required relationship is given by,  $a = b + \frac{2}{3}$

#### Question 18:

For what value of  $\lambda$  is the function defined by

$$f(x) = \begin{cases} \lambda(x^2 - 2x), & \text{if } x \leq 0 \\ 4x+1, & \text{if } x > 0 \end{cases}$$

continuous at  $x = 0$ ? What about continuity at  $x = 1$ ?

Answer

The given function  $f$  is 
$$f(x) = \begin{cases} \lambda(x^2 - 2x), & \text{if } x \leq 0 \\ 4x+1, & \text{if } x > 0 \end{cases}$$

If  $f$  is continuous at  $x = 0$ , then

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

$$\Rightarrow \lim_{x \rightarrow 0^-} \lambda(x^2 - 2x) = \lim_{x \rightarrow 0^+} (4x+1) = \lambda(0^2 - 2 \times 0)$$

$$\Rightarrow \lambda(0^2 - 2 \times 0) = 4 \times 0 + 1 = 1$$

$$\Rightarrow 0 = 1 = 0, \text{ which is not possible}$$

Therefore, there is no value of  $\lambda$  for which  $f$  is continuous at  $x = 0$

At  $x = 1$ ,

$$f(1) = 4x + 1 = 4 \times 1 + 1 = 5$$

$$\lim_{x \rightarrow 1} (4x + 1) = 4 \times 1 + 1 = 5$$

$$\therefore \lim_{x \rightarrow 1} f(x) = f(1)$$

Therefore, for any values of  $\lambda$ ,  $f$  is continuous at  $x = 1$

### Question 19:

Show that the function defined by  $g(x) = x - [x]$  is discontinuous at all integral point.

Here  $[x]$  denotes the greatest integer less than or equal to  $x$ .

Answer

The given function is  $g(x) = x - [x]$

It is evident that  $g$  is defined at all integral points.

Let  $n$  be an integer.

Then,

$$g(n) = n - [n] = n - n = 0$$

The left hand limit of  $f$  at  $x = n$  is,

$$\lim_{x \rightarrow n^-} g(x) = \lim_{x \rightarrow n^-} (x - [x]) = \lim_{x \rightarrow n^-} (x) - \lim_{x \rightarrow n^-} [x] = n - (n-1) = 1$$

The right hand limit of  $f$  at  $x = n$  is,

$$\lim_{x \rightarrow n^+} g(x) = \lim_{x \rightarrow n^+} (x - [x]) = \lim_{x \rightarrow n^+} (x) - \lim_{x \rightarrow n^+} [x] = n - n = 0$$

It is observed that the left and right hand limits of  $f$  at  $x = n$  do not coincide.

Therefore,  $f$  is not continuous at  $x = n$

Hence,  $g$  is discontinuous at all integral points.

### Question 20:

Is the function defined by  $f(x) = x^2 - \sin x + 5$  continuous at  $x = p$ ?

Answer

The given function is  $f(x) = x^2 - \sin x + 5$

It is evident that  $f$  is defined at  $x = p$

$$\text{At } x = \pi, f(x) = f(\pi) = \pi^2 - \sin \pi + 5 = \pi^2 - 0 + 5 = \pi^2 + 5$$

$$\text{Consider } \lim_{x \rightarrow \pi} f(x) = \lim_{x \rightarrow \pi} (x^2 - \sin x + 5)$$

Put  $x = \pi + h$

If  $x \rightarrow \pi$ , then it is evident that  $h \rightarrow 0$

$$\begin{aligned} \therefore \lim_{x \rightarrow \pi} f(x) &= \lim_{x \rightarrow \pi} (x^2 - \sin x + 5) \\ &= \lim_{h \rightarrow 0} [(\pi + h)^2 - \sin(\pi + h) + 5] \\ &= \lim_{h \rightarrow 0} (\pi + h)^2 - \lim_{h \rightarrow 0} \sin(\pi + h) + \lim_{h \rightarrow 0} 5 \\ &= (\pi + 0)^2 - \lim_{h \rightarrow 0} [\sin \pi \cosh + \cos \pi \sinh] + 5 \\ &= \pi^2 - \lim_{h \rightarrow 0} \sin \pi \cosh - \lim_{h \rightarrow 0} \cos \pi \sinh + 5 \\ &= \pi^2 - \sin \pi \cos 0 - \cos \pi \sin 0 + 5 \\ &= \pi^2 - 0 \times 1 - (-1) \times 0 + 5 \\ &= \pi^2 + 5 \end{aligned}$$

$$\therefore \lim_{x \rightarrow \pi} f(x) = f(\pi)$$

Therefore, the given function  $f$  is continuous at  $x = \pi$

### Question 21:

Discuss the continuity of the following functions.

(a)  $f(x) = \sin x + \cos x$

(b)  $f(x) = \sin x - \cos x$

(c)  $f(x) = \sin x \times \cos x$

Answer

It is known that if  $g$  and  $h$  are two continuous functions, then

$g + h$ ,  $g - h$ , and  $g \cdot h$  are also continuous.

It has to be proved first that  $g(x) = \sin x$  and  $h(x) = \cos x$  are continuous functions.

Let  $g(x) = \sin x$

It is evident that  $g(x) = \sin x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$g(c) = \sin c$$

$$\begin{aligned}\lim_{x \rightarrow c} g(x) &= \lim_{x \rightarrow c} \sin x \\ &= \lim_{h \rightarrow 0} \sin(c+h) \\ &= \lim_{h \rightarrow 0} [\sin c \cos h + \cos c \sin h] \\ &= \lim_{h \rightarrow 0} (\sin c \cos h) + \lim_{h \rightarrow 0} (\cos c \sin h) \\ &= \sin c \cos 0 + \cos c \sin 0 \\ &= \sin c + 0 \\ &= \sin c\end{aligned}$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is a continuous function.

$$\text{Let } h(x) = \cos x$$

It is evident that  $h(x) = \cos x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$h(c) = \cos c$$

$$\begin{aligned}\lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \cos x \\ &= \lim_{h \rightarrow 0} \cos(c+h) \\ &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\ &= \lim_{h \rightarrow 0} \cos c \cos h - \lim_{h \rightarrow 0} \sin c \sin h \\ &= \cos c \cos 0 - \sin c \sin 0 \\ &= \cos c \times 1 - \sin c \times 0 \\ &= \cos c\end{aligned}$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h$  is a continuous function.

Therefore, it can be concluded that

- (a)  $f(x) = g(x) + h(x) = \sin x + \cos x$  is a continuous function
- (b)  $f(x) = g(x) - h(x) = \sin x - \cos x$  is a continuous function
- (c)  $f(x) = g(x) \times h(x) = \sin x \times \cos x$  is a continuous function

**Question 22:**

Discuss the continuity of the cosine, cosecant, secant and cotangent functions,

Answer

It is known that if  $g$  and  $h$  are two continuous functions, then

$$(i) \frac{h(x)}{g(x)}, g(x) \neq 0 \text{ is continuous}$$

$$(ii) \frac{1}{g(x)}, g(x) \neq 0 \text{ is continuous}$$

$$(iii) \frac{1}{h(x)}, h(x) \neq 0 \text{ is continuous}$$

It has to be proved first that  $g(x) = \sin x$  and  $h(x) = \cos x$  are continuous functions.

Let  $g(x) = \sin x$

It is evident that  $g(x) = \sin x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$g(c) = \sin c$$

$$\begin{aligned} \lim_{x \rightarrow c} g(x) &= \lim_{x \rightarrow c} \sin x \\ &= \lim_{h \rightarrow 0} \sin(c+h) \\ &= \lim_{h \rightarrow 0} [\sin c \cos h + \cos c \sin h] \\ &= \lim_{h \rightarrow 0} (\sin c \cos h) + \lim_{h \rightarrow 0} (\cos c \sin h) \\ &= \sin c \cos 0 + \cos c \sin 0 \\ &= \sin c + 0 \\ &= \sin c \end{aligned}$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is a continuous function.

Let  $h(x) = \cos x$

It is evident that  $h(x) = \cos x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$h(c) = \cos c$$

$$\begin{aligned}
 \lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \cos x \\
 &= \lim_{h \rightarrow 0} \cos(c+h) \\
 &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\
 &= \lim_{h \rightarrow 0} \cos c \cos h - \lim_{h \rightarrow 0} \sin c \sin h \\
 &= \cos c \cos 0 - \sin c \sin 0 \\
 &= \cos c \times 1 - \sin c \times 0 \\
 &= \cos c
 \end{aligned}$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h(x) = \cos x$  is continuous function.

It can be concluded that,

$$\begin{aligned}
 \operatorname{cosec} x &= \frac{1}{\sin x}, \sin x \neq 0 \text{ is continuous} \\
 \Rightarrow \operatorname{cosec} x, x \neq n\pi \ (n \in \mathbf{Z}) \text{ is continuous}
 \end{aligned}$$

Therefore, cosecant is continuous except at  $x = n\pi, n \in \mathbf{Z}$

$$\begin{aligned}
 \sec x &= \frac{1}{\cos x}, \cos x \neq 0 \text{ is continuous} \\
 \Rightarrow \sec x, x \neq (2n+1)\frac{\pi}{2} \ (n \in \mathbf{Z}) \text{ is continuous}
 \end{aligned}$$

Therefore, secant is continuous except at  $x = (2n+1)\frac{\pi}{2} \ (n \in \mathbf{Z})$

$$\begin{aligned}
 \cot x &= \frac{\cos x}{\sin x}, \sin x \neq 0 \text{ is continuous} \\
 \Rightarrow \cot x, x \neq n\pi \ (n \in \mathbf{Z}) \text{ is continuous}
 \end{aligned}$$

Therefore, cotangent is continuous except at  $x = n\pi, n \in \mathbf{Z}$

### Question 23:

Find the points of discontinuity of  $f$ , where

$$f(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x < 0 \\ x+1, & \text{if } x \geq 0 \end{cases}$$



Answer

$$f(x) = \begin{cases} \frac{\sin x}{x}, & \text{if } x < 0 \\ x+1, & \text{if } x \geq 0 \end{cases}$$

The given function  $f$  is

It is evident that  $f$  is defined at all points of the real line.

Let  $c$  be a real number.

Case I:

$$\text{If } c < 0, \text{ then } f(c) = \frac{\sin c}{c} \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \left( \frac{\sin x}{x} \right) = \frac{\sin c}{c}$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x < 0$

Case II:

$$\text{If } c > 0, \text{ then } f(c) = c+1 \text{ and } \lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (x+1) = c+1$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x > 0$

Case III:

$$\text{If } c = 0, \text{ then } f(c) = f(0) = 0+1 = 1$$

The left hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \frac{\sin x}{x} = 1$$

The right hand limit of  $f$  at  $x = 0$  is,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} (x+1) = 1$$

$$\therefore \lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

From the above observations, it can be concluded that  $f$  is continuous at all points of the real line.

Thus,  $f$  has no point of discontinuity.

**Question 24:**

Determine if  $f$  defined by

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

is a continuous function?

Answer

$$f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & \text{if } x \neq 0 \\ 0, & \text{if } x = 0 \end{cases}$$

The given function  $f$  is

It is evident that  $f$  is defined at all points of the real line.

Let  $c$  be a real number.

Case I:

$$\text{If } c \neq 0, \text{ then } f(c) = c^2 \sin \frac{1}{c}$$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} \left( x^2 \sin \frac{1}{x} \right) = \left( \lim_{x \rightarrow c} x^2 \right) \left( \lim_{x \rightarrow c} \sin \frac{1}{x} \right) = c^2 \sin \frac{1}{c}$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x \neq 0$

Case II:

$$\text{If } c = 0, \text{ then } f(0) = 0$$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^-} \left( x^2 \sin \frac{1}{x} \right) = \lim_{x \rightarrow 0} \left( x^2 \sin \frac{1}{x} \right)$$

It is known that,  $-1 \leq \sin \frac{1}{x} \leq 1$ ,  $x \neq 0$

$$\Rightarrow -x^2 \leq \sin \frac{1}{x} \leq x^2$$

$$\Rightarrow \lim_{x \rightarrow 0} (-x^2) \leq \lim_{x \rightarrow 0} \left( x^2 \sin \frac{1}{x} \right) \leq \lim_{x \rightarrow 0} x^2$$

$$\Rightarrow 0 \leq \lim_{x \rightarrow 0} \left( x^2 \sin \frac{1}{x} \right) \leq 0$$

$$\Rightarrow \lim_{x \rightarrow 0} \left( x^2 \sin \frac{1}{x} \right) = 0$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = 0$$

$$\text{Similarly, } \lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \left( x^2 \sin \frac{1}{x} \right) = \lim_{x \rightarrow 0} \left( x^2 \sin \frac{1}{x} \right) = 0$$

$$\therefore \lim_{x \rightarrow 0} f(x) = f(0) = \lim_{x \rightarrow 0} f(x)$$

Therefore,  $f$  is continuous at  $x = 0$

From the above observations, it can be concluded that  $f$  is continuous at every point of the real line.

Thus,  $f$  is a continuous function.

### Question 25:

Examine the continuity of  $f$ , where  $f$  is defined by

$$f(x) = \begin{cases} \sin x - \cos x, & \text{if } x \neq 0 \\ -1 & \text{if } x = 0 \end{cases}$$

Answer

$$f(x) = \begin{cases} \sin x - \cos x, & \text{if } x \neq 0 \\ -1 & \text{if } x = 0 \end{cases}$$

The given function  $f$  is

It is evident that  $f$  is defined at all points of the real line.

Let  $c$  be a real number.

Case I:



If  $c \neq 0$ , then  $f(c) = \sin c - \cos c$

$$\lim_{x \rightarrow c} f(x) = \lim_{x \rightarrow c} (\sin x - \cos x) = \sin c - \cos c$$

$$\therefore \lim_{x \rightarrow c} f(x) = f(c)$$

Therefore,  $f$  is continuous at all points  $x$ , such that  $x \neq 0$

Case II:

If  $c = 0$ , then  $f(0) = -1$

$$\lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} (\sin x - \cos x) = \sin 0 - \cos 0 = 0 - 1 = -1$$

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} (\sin x - \cos x) = \sin 0 - \cos 0 = 0 - 1 = -1$$

$$\therefore \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0^+} f(x) = f(0)$$

Therefore,  $f$  is continuous at  $x = 0$

From the above observations, it can be concluded that  $f$  is continuous at every point of the real line.

Thus,  $f$  is a continuous function.

#### Question 26:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point.

$$f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases} \quad \text{at } x = \frac{\pi}{2}$$

Answer

$$f(x) = \begin{cases} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 3, & \text{if } x = \frac{\pi}{2} \end{cases}$$

The given function  $f$  is

The given function  $f$  is continuous at  $x = \frac{\pi}{2}$ , if  $f$  is defined at  $x = \frac{\pi}{2}$  and if the value of the  $f$

at  $x = \frac{\pi}{2}$  equals the limit of  $f$  at  $x = \frac{\pi}{2}$ .

It is evident that  $f$  is defined at  $x = \frac{\pi}{2}$  and  $f\left(\frac{\pi}{2}\right) = 3$

$$\lim_{x \rightarrow \frac{\pi}{2}} f(x) = \lim_{x \rightarrow \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x}$$

$$\text{Put } x = \frac{\pi}{2} + h$$

$$\text{Then, } x \rightarrow \frac{\pi}{2} \Rightarrow h \rightarrow 0$$

$$\begin{aligned} \therefore \lim_{x \rightarrow \frac{\pi}{2}} f(x) &= \lim_{x \rightarrow \frac{\pi}{2}} \frac{k \cos x}{\pi - 2x} = \lim_{h \rightarrow 0} \frac{k \cos\left(\frac{\pi}{2} + h\right)}{\pi - 2\left(\frac{\pi}{2} + h\right)} \\ &= k \lim_{h \rightarrow 0} \frac{-\sin h}{-2h} = \frac{k}{2} \lim_{h \rightarrow 0} \frac{\sin h}{h} = \frac{k}{2} \cdot 1 = \frac{k}{2} \end{aligned}$$

$$\therefore \lim_{x \rightarrow \frac{\pi}{2}} f(x) = f\left(\frac{\pi}{2}\right)$$

$$\Rightarrow \frac{k}{2} = 3$$

$$\Rightarrow k = 6$$

Therefore, the required value of  $k$  is 6.

#### Question 27:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point.

$$f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases} \quad \text{at } x = 2$$

Answer

$$f(x) = \begin{cases} kx^2, & \text{if } x \leq 2 \\ 3, & \text{if } x > 2 \end{cases}$$

The given function is

The given function  $f$  is continuous at  $x = 2$ , if  $f$  is defined at  $x = 2$  and if the value of  $f$  at  $x = 2$  equals the limit of  $f$  at  $x = 2$

It is evident that  $f$  is defined at  $x = 2$  and  $f(2) = k(2)^2 = 4k$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^+} f(x) = f(2)$$

$$\Rightarrow \lim_{x \rightarrow 2^-} (kx^2) = \lim_{x \rightarrow 2^+} (3) = 4k$$

$$\Rightarrow k \times 2^2 = 3 = 4k$$

$$\Rightarrow 4k = 3 = 4k$$

$$\Rightarrow 4k = 3$$

$$\Rightarrow k = \frac{3}{4}$$

Therefore, the required value of  $k$  is  $\frac{3}{4}$ .

#### Question 28:

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point.

$$f(x) = \begin{cases} kx+1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases} \quad \text{at } x = \pi$$

Answer

$$f(x) = \begin{cases} kx+1, & \text{if } x \leq \pi \\ \cos x, & \text{if } x > \pi \end{cases}$$

The given function is

The given function  $f$  is continuous at  $x = p$ , if  $f$  is defined at  $x = p$  and if the value of  $f$  at  $x = p$  equals the limit of  $f$  at  $x = p$

It is evident that  $f$  is defined at  $x = p$  and  $f(\pi) = k\pi + 1$

$$\lim_{x \rightarrow \pi^-} f(x) = \lim_{x \rightarrow \pi^+} f(x) = f(\pi)$$

$$\Rightarrow \lim_{x \rightarrow \pi^-} (kx+1) = \lim_{x \rightarrow \pi^+} \cos x = k\pi + 1$$

$$\Rightarrow k\pi + 1 = \cos \pi = k\pi + 1$$

$$\Rightarrow k\pi + 1 = -1 = k\pi + 1$$

$$\Rightarrow k = -\frac{2}{\pi}$$

Therefore, the required value of  $k$  is  $-\frac{2}{\pi}$ .

**Question 29:**

Find the values of  $k$  so that the function  $f$  is continuous at the indicated point.

$$f(x) = \begin{cases} kx+1, & \text{if } x \leq 5 \\ 3x-5, & \text{if } x > 5 \end{cases} \quad \text{at } x = 5$$

Answer

$$f(x) = \begin{cases} kx+1, & \text{if } x \leq 5 \\ 3x-5, & \text{if } x > 5 \end{cases}$$

The given function  $f$  is

The given function  $f$  is continuous at  $x = 5$ , if  $f$  is defined at  $x = 5$  and if the value of  $f$  at  $x = 5$  equals the limit of  $f$  at  $x = 5$

It is evident that  $f$  is defined at  $x = 5$  and  $f(5) = kx+1 = 5k+1$

$$\lim_{x \rightarrow 5^-} f(x) = \lim_{x \rightarrow 5^+} f(x) = f(5)$$

$$\Rightarrow \lim_{x \rightarrow 5^-} (kx+1) = \lim_{x \rightarrow 5^+} (3x-5) = 5k+1$$

$$\Rightarrow 5k+1 = 15-5 = 5k+1$$

$$\Rightarrow 5k+1 = 10$$

$$\Rightarrow 5k = 9$$

$$\Rightarrow k = \frac{9}{5}$$

Therefore, the required value of  $k$  is  $\frac{9}{5}$ .

**Question 30:**

Find the values of  $a$  and  $b$  such that the function defined by

$$f(x) = \begin{cases} 5, & \text{if } x \leq 2 \\ ax+b, & \text{if } 2 < x < 10 \\ 21, & \text{if } x \geq 10 \end{cases}$$

is a continuous function.

Answer

$$f(x) = \begin{cases} 5, & \text{if } x \leq 2 \\ ax + b, & \text{if } 2 < x < 10 \\ 21, & \text{if } x \geq 10 \end{cases}$$

The given function  $f$  is

It is evident that the given function  $f$  is defined at all points of the real line.

If  $f$  is a continuous function, then  $f$  is continuous at all real numbers.

In particular,  $f$  is continuous at  $x = 2$  and  $x = 10$

Since  $f$  is continuous at  $x = 2$ , we obtain

$$\begin{aligned} \lim_{x \rightarrow 2^-} f(x) &= \lim_{x \rightarrow 2^+} f(x) = f(2) \\ \Rightarrow \lim_{x \rightarrow 2^-} (5) &= \lim_{x \rightarrow 2^+} (ax + b) = 5 \\ \Rightarrow 5 &= 2a + b = 5 \\ \Rightarrow 2a + b &= 5 \quad \dots(1) \end{aligned}$$

Since  $f$  is continuous at  $x = 10$ , we obtain

$$\begin{aligned} \lim_{x \rightarrow 10^-} f(x) &= \lim_{x \rightarrow 10^+} f(x) = f(10) \\ \Rightarrow \lim_{x \rightarrow 10^-} (ax + b) &= \lim_{x \rightarrow 10^+} (21) = 21 \\ \Rightarrow 10a + b &= 21 = 21 \\ \Rightarrow 10a + b &= 21 \quad \dots(2) \end{aligned}$$

On subtracting equation (1) from equation (2), we obtain

$$\begin{aligned} 8a &= 16 \\ \Rightarrow a &= 2 \end{aligned}$$

By putting  $a = 2$  in equation (1), we obtain

$$\begin{aligned} 2 \times 2 + b &= 5 \\ \Rightarrow 4 + b &= 5 \\ \Rightarrow b &= 1 \end{aligned}$$

Therefore, the values of  $a$  and  $b$  for which  $f$  is a continuous function are 2 and 1 respectively.



**Question 31:**

Show that the function defined by  $f(x) = \cos(x^2)$  is a continuous function.

Answer

The given function is  $f(x) = \cos(x^2)$

This function  $f$  is defined for every real number and  $f$  can be written as the composition of two functions as,

$f = g \circ h$ , where  $g(x) = \cos x$  and  $h(x) = x^2$

$$\left[ \because (g \circ h)(x) = g(h(x)) = g(x^2) = \cos(x^2) = f(x) \right]$$

It has to be first proved that  $g(x) = \cos x$  and  $h(x) = x^2$  are continuous functions.

It is evident that  $g$  is defined for every real number.

Let  $c$  be a real number.

Then,  $g(c) = \cos c$

Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$\begin{aligned} \lim_{x \rightarrow c} g(x) &= \lim_{x \rightarrow c} \cos x \\ &= \lim_{h \rightarrow 0} \cos(c + h) \\ &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\ &= \lim_{h \rightarrow 0} \cos c \cos h - \lim_{h \rightarrow 0} \sin c \sin h \\ &= \cos c \cos 0 - \sin c \sin 0 \\ &= \cos c \times 1 - \sin c \times 0 \\ &= \cos c \end{aligned}$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g(x) = \cos x$  is continuous function.

$$h(x) = x^2$$

Clearly,  $h$  is defined for every real number.

Let  $k$  be a real number, then  $h(k) = k^2$

$$\lim_{x \rightarrow k} h(x) = \lim_{x \rightarrow k} x^2 = k^2$$

$$\therefore \lim_{x \rightarrow k} h(x) = h(k)$$

Therefore,  $h$  is a continuous function.

It is known that for real valued functions  $g$  and  $h$ , such that  $(g \circ h)$  is defined at  $c$ , if  $g$  is continuous at  $c$  and if  $h$  is continuous at  $g(c)$ , then  $(g \circ h)$  is continuous at  $c$ .

Therefore,  $f(x) = (g \circ h)(x) = \cos(x^2)$  is a continuous function.

### Question 32:

Show that the function defined by  $f(x) = |\cos x|$  is a continuous function.

Answer

The given function is  $f(x) = |\cos x|$

This function  $f$  is defined for every real number and  $f$  can be written as the composition of two functions as,

$$f = g \circ h, \text{ where } g(x) = |x| \text{ and } h(x) = \cos x$$

$$[\because (g \circ h)(x) = g(h(x)) = g(\cos x) = |\cos x| = f(x)]$$

It has to be first proved that  $g(x) = |x|$  and  $h(x) = \cos x$  are continuous functions.

$g(x) = |x|$  can be written as

$$g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$$

Clearly,  $g$  is defined for all real numbers.

Let  $c$  be a real number.

Case I:

$$\text{If } c < 0, \text{ then } g(c) = -c \text{ and } \lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x < 0$

Case II:

If  $c > 0$ , then  $g(c) = c$  and  $\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} x = c$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x > 0$

Case III:

If  $c = 0$ , then  $g(c) = g(0) = 0$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} (x) = g(0)$$

Therefore,  $g$  is continuous at  $x = 0$

From the above three observations, it can be concluded that  $g$  is continuous at all points.

$$h(x) = \cos x$$

It is evident that  $h(x) = \cos x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + h$

If  $x \rightarrow c$ , then  $h \rightarrow 0$

$$h(c) = \cos c$$

$$\begin{aligned} \lim_{x \rightarrow c} h(x) &= \lim_{x \rightarrow c} \cos x \\ &= \lim_{h \rightarrow 0} \cos(c + h) \\ &= \lim_{h \rightarrow 0} [\cos c \cos h - \sin c \sin h] \\ &= \lim_{h \rightarrow 0} \cos c \cos h - \lim_{h \rightarrow 0} \sin c \sin h \\ &= \cos c \cos 0 - \sin c \sin 0 \\ &= \cos c \times 1 - \sin c \times 0 \\ &= \cos c \end{aligned}$$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h(x) = \cos x$  is a continuous function.

It is known that for real valued functions  $g$  and  $h$ , such that  $(g \circ h)$  is defined at  $c$ , if  $g$  is continuous at  $c$  and if  $f$  is continuous at  $g(c)$ , then  $(f \circ g)$  is continuous at  $c$ .

Therefore,  $f(x) = (goh)(x) = g(h(x)) = g(\cos x) = |\cos x|$  is a continuous function.

**Question 33:**

Examine that  $\sin|x|$  is a continuous function.

Answer

$$\text{Let } f(x) = \sin|x|$$

This function  $f$  is defined for every real number and  $f$  can be written as the composition of two functions as,

$$f = g \circ h, \text{ where } g(x) = |x| \text{ and } h(x) = \sin x$$

$$[\because (goh)(x) = g(h(x)) = g(\sin x) = |\sin x| = f(x)]$$

It has to be proved first that  $g(x) = |x|$  and  $h(x) = \sin x$  are continuous functions.

$g(x) = |x|$  can be written as

$$g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$$

Clearly,  $g$  is defined for all real numbers.

Let  $c$  be a real number.

Case I:

$$\text{If } c < 0, \text{ then } g(c) = -c \text{ and } \lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x < 0$

Case II:

$$\text{If } c > 0, \text{ then } g(c) = c \text{ and } \lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} x = c$$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x > 0$

Case III:

$$\text{If } c = 0, \text{ then } g(c) = g(0) = 0$$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} (x) = g(0)$$

Therefore,  $g$  is continuous at  $x = 0$

From the above three observations, it can be concluded that  $g$  is continuous at all points.

$$h(x) = \sin x$$

It is evident that  $h(x) = \sin x$  is defined for every real number.

Let  $c$  be a real number. Put  $x = c + k$

If  $x \rightarrow c$ , then  $k \rightarrow 0$

$$h(c) = \sin c$$

$$h(c) = \sin c$$

$$\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} \sin x$$

$$= \lim_{k \rightarrow 0} \sin(c + k)$$

$$= \lim_{k \rightarrow 0} [\sin c \cos k + \cos c \sin k]$$

$$= \lim_{k \rightarrow 0} (\sin c \cos k) + \lim_{k \rightarrow 0} (\cos c \sin k)$$

$$= \sin c \cos 0 + \cos c \sin 0$$

$$= \sin c + 0$$

$$= \sin c$$

$$\therefore \lim_{x \rightarrow c} h(x) = g(c)$$

Therefore,  $h$  is a continuous function.

It is known that for real valued functions  $g$  and  $h$ , such that  $(g \circ h)$  is defined at  $c$ , if  $g$  is continuous at  $c$  and if  $h$  is continuous at  $g(c)$ , then  $(g \circ h)$  is continuous at  $c$ .

Therefore,  $f(x) = (g \circ h)(x) = g(h(x)) = g(\sin x) = |\sin x|$  is a continuous function.

#### Question 34:

Find all the points of discontinuity of  $f$  defined by  $f(x) = |x| - |x+1|$ .

Answer

The given function is  $f(x) = |x| - |x+1|$

The two functions,  $g$  and  $h$ , are defined as

$$g(x) = |x| \text{ and } h(x) = |x+1|$$

Then,  $f = g - h$

The continuity of  $g$  and  $h$  is examined first.

$g(x) = |x|$  can be written as

$$g(x) = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases}$$

Clearly,  $g$  is defined for all real numbers.

Let  $c$  be a real number.

Case I:

If  $c < 0$ , then  $g(c) = -c$  and  $\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} (-x) = -c$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x < 0$

Case II:

If  $c > 0$ , then  $g(c) = c$  and  $\lim_{x \rightarrow c} g(x) = \lim_{x \rightarrow c} x = c$

$$\therefore \lim_{x \rightarrow c} g(x) = g(c)$$

Therefore,  $g$  is continuous at all points  $x$ , such that  $x > 0$

Case III:

If  $c = 0$ , then  $g(c) = g(0) = 0$

$$\lim_{x \rightarrow 0^-} g(x) = \lim_{x \rightarrow 0^-} (-x) = 0$$

$$\lim_{x \rightarrow 0^+} g(x) = \lim_{x \rightarrow 0^+} (x) = 0$$

$$\therefore \lim_{x \rightarrow 0} g(x) = \lim_{x \rightarrow 0} (x) = g(0)$$

Therefore,  $g$  is continuous at  $x = 0$

From the above three observations, it can be concluded that  $g$  is continuous at all points.

$h(x) = |x+1|$  can be written as

$$h(x) = \begin{cases} -(x+1), & \text{if } x < -1 \\ x+1, & \text{if } x \geq -1 \end{cases}$$

Clearly,  $h$  is defined for every real number.

Let  $c$  be a real number.

Case I:

If  $c < -1$ , then  $h(c) = -(c+1)$  and  $\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} [-(x+1)] = -(c+1)$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h$  is continuous at all points  $x$ , such that  $x < -1$

Case II:

If  $c > -1$ , then  $h(c) = c+1$  and  $\lim_{x \rightarrow c} h(x) = \lim_{x \rightarrow c} (x+1) = c+1$

$$\therefore \lim_{x \rightarrow c} h(x) = h(c)$$

Therefore,  $h$  is continuous at all points  $x$ , such that  $x > -1$

Case III:

If  $c = -1$ , then  $h(c) = h(-1) = -1+1 = 0$

$$\lim_{x \rightarrow -1^-} h(x) = \lim_{x \rightarrow -1^-} [-(x+1)] = -(-1+1) = 0$$

$$\lim_{x \rightarrow -1^+} h(x) = \lim_{x \rightarrow -1^+} (x+1) = (-1+1) = 0$$

$$\therefore \lim_{x \rightarrow -1^-} h(x) = \lim_{h \rightarrow -1^+} h(x) = h(-1)$$

Therefore,  $h$  is continuous at  $x = -1$

From the above three observations, it can be concluded that  $h$  is continuous at all points of the real line.

$g$  and  $h$  are continuous functions. Therefore,  $f = g - h$  is also a continuous function.

Therefore,  $f$  has no point of discontinuity.