

NEET Revision Notes

Biology

Excretory Products and their Elimination

Introduction

- As a consequence of multiple metabolic reactions, the body produces a variety of harmful substances. Among other things, it includes urea, ammonia, uric acid, carbon dioxide, ions, and water.
- Urea, ammonia, and uric acid are the three main types of nitrogenous wastes present in higher animals. Ammonia is among the most hazardous nitrogenous wastes.

Types of Organisms based on their Excretory Products:

1. Ammonotelic animals:

- They include most bony fishes but also aquatic amphibians, which eject ammonia as nitrogenous waste.
- Diffusion is how ammonia is excreted.

2. Ureotelic organisms:

- They secrete urea as nitrogenous waste.
- Examples include mammals, amphibians, and other ureotelic organisms.

3. Uricotelic organisms:

- Uricotelic animals such as reptiles and birds, detoxify uric acid as nitrogenous waste.

Types of Excretory Organs:

- Different organisms have different excretory structures.
- *Amoeba* and *Paramecium* possess **contractile vacuoles** for excretion.
- The excretory system in **sponges** is identified as the **canal system**. Sponges expel waste via a canal system.
- *Hydra* has the excretory cells called **coelenteron**.
- **Flame cells** are found in **Platyhelminthes**.
- **Nephridia** is found in Annelids such as **earthworms**.
- **Green excretory glands** are found in **prawns**.

- **Malpighian tubules** are the excretory system of **insects**.

Human Excretory System

- Organs including **two kidneys, two ureters, a urinary bladder, and a urethra** comprise the human excretory system.




Image: Human Excretory System

- The **kidneys** are bean-shaped organs located in the abdomen. The right kidney is located a bit lower in the body as compared to the left.
- The kidney's inner side is concave, while the outside is convex.




Image: Kidney

- The **hilum** is a notch located near the centre of the kidney's inner concave surface. It is the point where the ureter, blood vessels, as well as nerves, all enter.
- A funnel-shaped site inside the hilum with calyces projections is referred to as the **renal pelvis**.
- The kidneys are split into two parts: the outer kidney and the inner kidney.
- The **cortex** is the outer portion of the kidney, and the **medulla** is the inner portion.
- The medulla is partitioned into conical masses known as **medullary pyramids**.
- **Bertini's column** is an extended form of the renal cortex that connects the pyramids.

Structure of Nephrons

- The **nephron** is the kidney's basic function as well as a structural unit.
- The **renal corpuscle** and **renal tubule** are two main components of the nephron.




Image: Nephron of human kidney

Renal Corpuscle/ Malpighian Body:




Image: Renal corpuscle

- A **glomerulus** is formed by a tuft of capillaries.
- The **afferent arteriole** carries blood into the glomerulus, whereas the **efferent arteriole** transports blood out.
- The glomerulus is held together by **Bowman's capsule**, which is a cup-shaped structure.
- Bowman's capsule, including the glomerulus, is termed the **Malpighian body or renal corpuscles**.

Renal Tubule:

- The **proximal convoluted tubule (PCT)** is an elongated tube-like structure of the Malpighian body that is strongly coiled.
- The tubule's next section is **Henle's loop**. Henle's loop has two limbs, one **ascending** and one **descending**.
- A **DCT i.e. distal convoluted tubule** is formed by the ascending limb. After that, the DCT links to the **collecting duct**.

Types of Nephrons:

- Cortical nephrons and medullary nephrons are the two major categories of nephrons.
- **Cortical Nephrons:** Cortical nephrons are created when only a tiny fraction of the Henle loop is found in the medulla due to its short length.
- **Medullary Nephrons** are created when the loop of Henle becomes too long and reaches the medulla.




Image: Types of nephrons (Cortical and Juxtamedullary)

Urine Production

- The process of urine formation completes in three steps which are:
 1. Glomerular Filtration
 2. Tubular reabsorption
 3. Urine Secretio

1. Glomerular Filtration:

- Glomerular filtration starts when blood travels the glomerulus by an afferent arteriole.
- Water and nitrogenous waste pass through the glomerulus as well as blood cells, whereas proteins exit via the efferent arteriole.
- On average, the kidney filters 1100 mL to 1200 mL of blood per minute.
- Blood is filtered through three separate layers due to glomerular capillary blood pressure.
- The **endothelium**, which encloses the glomerular blood vessels, is the first layer.
- **Bowman's capsule epithelium** is the second layer, separated by a **basement membrane**.
- Podocytes are Bowman's capsule epithelial cells that are positioned in an organised pattern to leave some tiny openings known as slit pores as well as **filtration slits**.
- The **glomerular filtration rate** is the quantity of filtrate manufactured by the kidneys per minute.

2. Tubular Reabsorption:



- Tubular reabsorption is the procedure of soaking up necessary molecules such as glucose, amino acids, and so on, or ions such as sodium ions, and so on.
- Some substances are actively absorbed, while others are passively absorbed.
- Water is passively (without energy) absorbed, whereas glucose and amino acids are actively (using energy) absorbed.

3. Urine Secretion:

- Urine secretion is the last step in the production of urine.
- To keep the ionic and acid balance of the body fluids, ammonia, potassium ions, hydrogen ions, etc, are released.

Functions of Nephron Tubules

1. PCT (Proximal Convolute Tubules):

- The proximal convolute tubules are lined with simple cuboidal brush border epithelium.
- This type of epithelium has a bigger surface area for reabsorption.
- The majority of electrolytes, as well as water, are reabsorbed during PCT.
- It helps to maintain the pH and ionic stability of body fluids by secreting ions like hydrogen, potassium, and ammonium into the filtrate.

2. Henle's Loop:

- Henle's loop is very useful in maintaining the osmolarity of the fluid.
- There is hardly any reabsorption in its ascending limb. It is impermeable to water but permeable to electrolytes.
- The major part of the water is absorbed by the descending limb, which concentrates the filtrate. All electrolytes are nearly impermeable to the descending limb.
- As a result, Henle's loop absorbs differently in different parts.

3. Distal Convolute Tubule (DCT):

- The distal convolute tubule soaks up water, sodium ions, as well as bicarbonate ions while removing potassium ions and hydrogen ions to maintain fluid ionic balance.

4. Collecting duct:

- To concentrate the urine, the collecting duct reabsorbs lots of water.

- Both Hydrogen ions and potassium ions are secreted by the collecting duct.
- It maintains the ionic balance and pH of the blood.




Image: Functions of different parts of Nephron

Regulation of Kidney functions

- Kidney regulation is aided by hormones secreted by the **hypothalamus**, **juxtaglomerular apparatus (JGA)**, and the **heart**.
- Any change in blood density or ionic balance initiates the osmoreceptors in the body.
- As a result, the **hypothalamus** secretes antidiuretic hormone (ADH)/ **vasopressin**.
- This aids in the reabsorption of water from the tubules.
- This causes an increase in blood volume and, as a result of a negative feedback loop, brings off the osmoreceptors.
- The **juxtaglomerular apparatus** is initiated when glomerular blood pressure falls.
- Juxtaglomerular cells secrete an enzyme called **Renin**, which transforms **angiotensinogen** in the blood to **angiotensin I**, which will then be converted to **angiotensin II**.

- Angiotensin II tends to raise glomerular blood pressure as it is an effective vasoconstrictor.
- Angiotensin II triggers the adrenal cortex to generate aldosterone, which raises the reabsorption of sodium ions and water by distal tubules.
- The renin-angiotensin-aldosterone system (RAAS) is the name given to this procedure.

Renin-Angiotensin-Aldosterone- System (RAAS):




Image: Renin-Angiotensin-Aldosterone- System (RAAS)

- The Renin-Angiotensin-Aldosterone System (RAAS) controls kidney function.
- The heart produces a polypeptide hormone known as an **atrial natriuretic factor** (ANF), which behaves as a vasodilator and hence lowers blood pressure.
- It acts as a renin-angiotensin system negative feedback mechanism.

Micturition:

- Micturition is the activity by which urine is excreted or released from the body. The CNS regulates it through a wide range of neural mechanisms.
- **Other Excretory Organs:** Wastes including carbon dioxide, harmful chemicals, urea, and so on are eliminated by the lungs, liver, kidneys, and skin working together.

Dysfunctions of the Excretory System

- **Uremia** is a buildup of urea in the blood caused by kidney failure.
- **Acute renal failure** happens when one or both kidneys are damaged to filter urine and become unable to act for a range of reasons.
- **Renal calculi** include kidney stones caused by the buildup of insoluble crystals including oxalates.
- **Glomerulonephritis** is an inflammation of the glomerulus.

Hemodialysis




Image: Procedure of Dialysis

- Hemodialysis is a method of cleaning and purifying human blood.
- It could be used to eliminate extra urea from the blood in cases of kidney failure.
- To clean the blood, a dialysis device and a special filter known as an artificial kidney are used.
- Blood is drawn from the body through cellophane tubules but also dialyzed against an isotonic fluid to extract wastes before being reintroduced.
- It is prescribed for people whose kidneys are not functioning normally.
- Hemodialysis necessitates a minor procedure, generally in the arm.

Points to remember:

- Many nitrogen-containing compounds, ions, carbon dioxide, water, and so on must be eliminated from the body.



- The type of nitrogenous wastes created and their excretion differs between animals, owing primarily to habitat and availability of water. The main nitrogenous wastes excreted are ammonia, urea, and uric acid.
- Animal excretory organs include protonephridia, malpighian tubules, nephridia, green glands, and the kidneys. They not only help to remove nitrogenous wastes, but also to maintain the ionic and acid-base equilibrium of body fluids.
- The excretory system in humans consists of one kidney, two ureters, a urinary bladder, and a urethra. Each kidney contains millions of tubular structures known as nephrons.
- The functional unit of the kidney is the nephron, which has two parts: the glomerulus and the renal tubule.
- Glomerulus is a capillary tuft formed by afferent arterioles, which are fine branches of the renal artery.
- The renal tubule begins with a double-walled Bowman's capsule and develops into a proximal convoluted tubule (PCT), Henle's loop (HL), and distal convoluted tubule (DCT).
- Many nephrons' DCTs connect to a common collecting duct, many of which eventually enter into the renal pelvis via the medullary pyramids.
- Bowman's capsule surrounds the glomerulus, forming the Malpighian or renal corpuscle.
- Urine formation consists of three major processes: filtration, reabsorption, and secretion.
- Filtration is a non-selective process that the glomerulus performs using glomerular capillary blood pressure.
- The glomerulus filters approximately 1200 ml of blood per minute, producing 125 ml of filtrate in the Bowman's capsule per minute.
- JGA, a specialised component of the nephrons, is important in the regulation of Glomerular Filtration Rate (GFR).
- Almost all of the filtrate is reabsorption through various parts of the nephrons.
- The primary site of reabsorption and preferential secretion is PCT.
- The primary function of HL is to maintain the osmolar gradient (300 mOsmolL⁻¹ -1200 mOsmolL⁻¹) in the kidney interstitium.
- The DCT and collecting duct allow for extensive water and electrolyte reabsorption, which aids in osmoregulation: H⁺, K⁺, and NH₃ could be

produced into the filtrate by the tubules to preserve the ionic balance as well as pH of body fluids.

- A countercurrent mechanism exists between the two limbs of the Henle loop and those of the vasa recta, a capillary connected to the Henle loop. As it runs down the descending limb, the filtrate becomes more concentrated, but it is diluted along the ascending limb. This arrangement keeps electrolytes and urea in the interstitium.
- The DCT and collecting duct concentrate the filtrate four times, from 300 mOsmolL⁻¹ to 1200 mOsmolL⁻¹ which is an outstanding water conservation mechanism.
- Urine is deposited in the urinary bladder until a voluntary signal from the CNS causes it to be released through the urethra, a process known as micturition. The skin, lungs, and liver all help with excretion.