

NEET 2025 Paper-(PCB)

Instructions:

46. The ratio of the wavelengths of the light absorbed by a Hydrogen atom when it undergoes $n = 2 \rightarrow n = 3$ and $n = 4 \rightarrow n = 6$ transitions, respectively, is

- (A) $\frac{1}{36}$ (B) $\frac{1}{16}$
(C) $\frac{1}{9}$ (D) $\frac{1}{4}$

NEET 2025

Correct Option: D

Solution:

$$\Delta E = \frac{hc}{\lambda} = E_{final} - E_{initial} \left(E_n = \frac{-R_H}{n^2} \right)$$

$$\Delta E_{2 \rightarrow 3} = \frac{hc}{\lambda_{2 \rightarrow 3}} = E_3 - E_2 = \frac{-R_H}{3^2} - \left(\frac{-R_H}{2^2} \right)$$

$$= R_H \left(\frac{1}{4} - \frac{1}{9} \right)$$

$$= R_H \times \frac{5}{36}$$

$$\therefore \lambda_{2 \rightarrow 3} = \frac{hc \cdot 36}{R_H \cdot 5}$$

$$\Delta E_{4 \rightarrow 6} = E_6 - E_4 = \frac{-R_H}{36} + \frac{R_H}{16} = \frac{R_H \times 20}{36 \times 16}$$

$$\frac{hc}{\lambda_{4 \rightarrow 6}} = \frac{R_H \times 20}{36 \times 16}$$

$$\lambda_{4 \rightarrow 6} = \frac{hc \times 36 \times 16}{R_H \cdot 20}$$

$$\frac{\lambda_{2 \rightarrow 3}}{\lambda_{4 \rightarrow 6}} = \frac{\frac{hc \cdot 36}{R_H \cdot 5}}{\frac{hc \times 36 \times 16}{R_H \cdot 20}}$$

$$= \frac{1}{4}$$

47. Which of the following statements are true?
- A. Unlike Ga that has a very high melting point, Cs has a very low melting point.
- B. On Pauling scale, the electronegativity values of N and Cl are not the same.
- C. Ar, K^+ , Cl^- , Ca^{2+} , and S^{2-} are all isoelectronic species.
- D. The correct order of the first ionization enthalpies of Na, Mg, Al, and Si is $Si > Al > Mg > Na$.
- E. The atomic radius of Cs is greater than that of Li and Rb.

Choose the correct answer from the options given below

- (A) A, B and E only (B) C and E only

(C) C and D only

(D) A, C and E only

NEET 20

Correct Option: B

Solution:

(C) Ca^{2+} , K^+ , Ar, Cl^- , S^{2-} all species have 18 electron that's why these are isoelectronic species.

(E) Radius order $Cs > Rb > K > Na > Li$ (4) $Li < Be$
 $N < B < C$

48. Match List I with List II

List I (Ion):		List II (Group Number in Cation Analysis):	
A.	Co^{2+}	I.	Group-I
B.	Mg^{2+}	II.	Group-III
C.	Pb^{2+}	III.	Group-IV
D.	Al^{3+}	IV.	Group-VI

Choose the correct answer from the option given below

- (A) A-III, B-IV, C-II, D-I (B) A-III, B-IV, C-I, D-
(C) A-III, B-II, C-IV, D-I (D) A-III, B-II, C-I, D-


NEET 20





Correct Option: B

Solution:


List I (Ion):		List II (Group Number in Cation Analysis):	
A.	Co^{2+}	I.	Group-IV
B.	Mg^{2+}	II.	Group-VI
C.	Pb^{2+}	III.	Group-I
D.	Al^{3+}	IV.	Group-III

49. Predict the major product 'P' in the following sequence of reactions-





NEET 2025

Correct Option: A**Solution:**

50. Energy and radius of first Bohr orbit of He^+ and Li^{2+} are
[Given $R_H = 2.18 \times 10^{-18} \text{ J}$, $a_0 = 52.9 \text{ pm}$]

(A) $E_n(\text{Li}^{2+}) = -19.62 \times 10^{-18} \text{ J}$;
 $r_n(\text{Li}^{2+}) = 17.6 \text{ pm}$
 $E_n(\text{He}^+) = -8.72 \times 10^{-18} \text{ J}$;
 $r_n(\text{He}^+) = 26.4 \text{ pm}$

(B) $E_n(\text{Li}^{2+}) = -8.72 \times 10^{-18} \text{ J}$;
 $r_n(\text{Li}^{2+}) = 26.4 \text{ pm}$
 $E_n(\text{He}^+) = -19.62 \times 10^{-18} \text{ J}$;
 $r_n(\text{He}^+) = 17.6 \text{ pm}$

(C) $E_n(\text{Li}^{2+}) = -19.62 \times 10^{-16} \text{ J}$;
 $r_n(\text{Li}^{2+}) = 17.6 \text{ pm}$
 $E_n(\text{He}^+) = -8.72 \times 10^{-16} \text{ J}$;
 $r_n(\text{He}^+) = 26.4 \text{ pm}$

(D) $E_n(\text{Li}^{2+}) = -8.72 \times 10^{-16} \text{ J}$;
 $r_n(\text{Li}^{2+}) = 17.6 \text{ pm}$
 $E_n(\text{He}^+) = -19.62 \times 10^{-16} \text{ J}$;
 $r_n(\text{He}^+) = 17.6 \text{ pm}$

NEET 20

Correct Option: A**Solution:**

$$E_n = \frac{-2.18 \times 10^{-18} \times z^2}{n^2} \text{ J}; r_n = \frac{52.9 \times n^2}{z} \text{ pm}$$

For He^+

$$E_{\text{He}^+} = -2.18 \times 10^{-18} \times 4 = -8.72 \times 10^{-18} \text{ J}$$

$$r_{\text{He}^+} = \frac{52.9 \times 1}{2} = 26.45 \text{ pm}$$

For Li^{2+}

$$E_{\text{Li}^{2+}} = -2.18 \times 10^{-18} \times 9 = -19.62 \times 10^{-18} \text{ J}$$

$$r_{\text{Li}^{2+}} = \frac{52.9 \times 1}{3} = 17.63 \text{ pm}$$

51. Which of the following are paramagnetic?

- A. $[\text{NiCl}_4]^{2-}$
 B. $\text{Ni}(\text{CO})_4$
 C. $[\text{Ni}(\text{CN})_4]^{2-}$
 D. $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$
 E. $\text{Ni}(\text{PPh}_3)_4$

Choose the correct answer from the options given below

- (A) A and C only (B) B and E only
 (C) A and D only (D) A, D and E only

NEET 20

Correct Option: C**Solution:**

A. $[\text{NiCl}_4]^{2-}$; Ni^{+2} ; $3d^8$; sp^3 hybridisation; 2 unpaired electrons; paramagnetic

B. $\text{Ni}(\text{CO})_4$; Ni ; $3d^8 4s^2$; sp^3 hybridisation; Zero unpaired electron; diamagnetic

C. $[\text{Ni}(\text{CN})_4]^{2-}$; Ni^{+2} ; $3d^8$; dsp^2 hybridisation; Zero unpaired electron; diamagnetic

D. $[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$; Ni^{2+} ; $3d^8$; $sp^3 d^2$ hybridisation; Two unpaired electron; paramagnetic

E. $\text{Ni}(\text{PPh}_3)_4$; Ni ; $3d^8 4s^2$; sp^3 hybridisation; zero unpaired electron; Diamagnetic

52. Given below are two statements:

Statement I : Like nitrogen that can form ammoniacal can form arsine.



Statement II : Antimony cannot form antimony pentoxide.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (A) Both Statement I and Statement II are correct
 (B) Both Statement I and Statement II are incorrect
 (C) Statement I is correct but Statement II is incorrect
 (D) Statement I is incorrect but Statement II is correct

NEET 2025

Correct Option: C

Solution:

$N = NH_3$ $As \rightarrow AsH_3$ are possible correct statement.

All elements of N-family form two types of E_2O_3 and E_2O_5 oxides.

Statement is wrong.

Statement-I is correct & Statement-II is incorrect.

53. Which among the following electronic configurations belong to main group elements (s & p-Block) ?

- A. $[Ne]3s^1$
 B. $[Ar]3d^34s^2$
 C. $[Kr]4d^{10}5s^25p^5$
 D. $[Ar]3d^{10}4s^1$
 E. $[Rn]5f^06d^27s^2$

- (A) B and E only (B) A and C only
 (C) D and E only (D) A, C and D only

NEET 2025

Correct Option: B

Solution:

Main group elements are s & p-block elements.

$A \rightarrow s\text{-block}$

$C \rightarrow p\text{-block}$

54. Dalton's Atomic theory could not explain which of the following?

- (A) Law of conservation of mass (B) Law of constant proportion
 (C) Law of multiple proportion (D) Law of gaseous volume

NEET 20

Correct Option: D

Solution:

Dalton's atomic theory could not explain law of gaseous volume.

55. Consider the following compounds:

KO_2 , H_2O_2 and H_2SO_4 .

The oxidation states of the underlined elements in are, respectively,

- (A) +1, -1, and +6 (B) +2, -2, and +6
 (C) +1, -2, and +4 (D) +4, -4, and +6

NEET 20

Correct Option: A

Solution:

KO_2 , H_2O_2
 $\downarrow$ $\downarrow$

Alkali metals $2(+1) + 2x = 0$

always show +1 $x = -1$

O.S. in its

compounds

H_2SO_4
 $\downarrow$

$2(+1) + x + 4(-2) = 0$

$x = +6$

56. If the half-life ($t_{1/2}$) for a first order reaction is 1 min then the time required for 99.9 % completion of the reaction is closest to:

- (A) 2 minutes (B) 4 minutes
 (C) 5 minutes (D) 10 minutes

NEET 20

Correct Option: D

Solution:

For 1st order reactions : $t_{99.9\%} = 10 \times t_{1/2}$

$t_{99.9\%} = 10 \times 1$
 $= 10 \text{ minute}$

57. The correct order of the wavelength of light absorbed by the following complexes is,

- A. $[\text{Co}(\text{NH}_3)_6]^{3+}$
 B. $[\text{Co}(\text{CN})_6]^{3-}$
 C. $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$
 D. $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$

Choose the correct answer from the options given below:

- (A) $B < D < A < C$
 (B) $B < A < D < C$
 (C) $C < D < A < B$
 (D) $C < A < D < B$

NEET 2025

Correct Option: B

Solution:

$$\lambda \propto \frac{1}{\text{strength of ligand}}$$

$$\lambda \propto \frac{1}{\text{splitting}}$$

- A. $[\text{Co}(\text{NH}_3)_6]^{3+}$; 475 nm
 B. $[\text{Co}(\text{CN})_6]^{3-}$; 310 nm
 C. $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$; 600 nm
 D. $[\text{Ti}(\text{H}_2\text{O})_6]^{3+}$; 498 nm

Order of $\lambda = C > D > A > B$

58. Which one of the following compounds can exist as cis-trans isomers?

- (A) Pent-1-ene (B) 2-Methylhex-2-ene
 (C) 1, 1-Dimethylcyclopropane (D) 1, 2-Dimethylcyclohexane

NEET 2025

Correct Option: D


Solution:

Cis-trans isomers shown by :

Condition: Restricted rotation around double bond

Or

Different group around double bond



59. Phosphoric acid ionizes in three steps with their ionization constant values K_{a1} , K_{a2} and K_{a3} , respectively, K is the overall ionization constant. Which of the following statements are true?

- A. $\log K = \log K_{a1} + \log K_{a2} + \log K_{a3}$
 B. H_3PO_4 is a stronger acid than H_2PO_4^- and HPO_4^{2-}
 C. $K_{a1} > K_{a2} > K_{a3}$
 D. $K_{a1} = \frac{K_{a2} + K_{a3}}{2}$

Choose the correct answer from the options given below:

- (A) A and B only (B) A and C only
 (C) B, C and D only (D) A, B and C only

NEET 2025

Correct Option: D

Solution:

H_3PO_4 is a stronger acid than H_2PO_4^- and HPO_4^{2-} .

$\text{H}_3\text{PO}_4(\text{aq}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{H}_2\text{PO}_4^-(\text{aq})$ $K_{a1} = 7.5 \times 10^{-3}$

$\text{H}_2\text{PO}_4^-(\text{aq}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{HPO}_4^{2-}(\text{aq})$ $K_{a2} = 6.2 \times 10^{-8}$


$\text{HPO}_4^{2-}(\text{aq}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{PO}_4^{3-}(\text{aq})$ $K_{a3} = 1.7 \times 10^{-13}$


$K_{a1} > K_{a2} > K_{a3}$

$\log K = \log K_{a1} + \log K_{a2} + \log K_{a3}$

Ans. (A), (B) and (C) only

60. Which one of the following reactions does NOT produce benzene as the product?







NEET 2025

Correct Option: D**Solution:**


(1)




(2)



(3)



(4)



61. If the molar conductivity (Λ_m) of a 0.050 mol L^{-1} solution of a monobasic weak acid is $90 \text{ Scm}^2 \text{ mol}^{-1}$, its extent (degree) of dissociation will be [Assume $\Lambda_+^\circ = 349.6 \text{ Scm}^2 \text{ mol}^{-1}$ and $\Lambda_-^\circ = 50.4 \text{ Scm}^2 \text{ mol}^{-1}$.]

(A) 0.115 (B) 0.125 (C) 0.225 (D) 0.215

NEET 2025

Correct Option: C**Solution:**

$$\begin{aligned} \text{Degree of dissociation } (\alpha) &= \frac{\Lambda_m}{\Lambda_m^\circ} \\ \Lambda_m^\circ &= \Lambda_+^\circ + \Lambda_-^\circ \\ &= 349.6 + 50.4 = 400 \text{ Scm}^2 \text{ mol}^{-1} \\ \therefore \alpha &= \frac{\Lambda_m}{\Lambda_m^\circ} = \frac{90}{400} = 0.225 \end{aligned}$$

62. Given below are two statements :


Statement I : A hypothetical diatomic molecule with bond order zero is quite stable.

Statement II : As bond order increases, the bond length increases.

In the light of the above statements, choose the most appropriate answer from the options given below:

- (A) Both Statement I and Statement II are true
(B) Both Statement I and Statement II are false
(C) Statement I is true but Statement II are false
(D) Statement I is false but Statement II are true

NEET 20

Correct Option: B**Solution:**

$$\text{Bond order} \propto \frac{1}{\text{Bond length}}$$

Statement II is false

63. Out of the following complex compounds, which of the following compound will be having the minimum conductance solution?

- (A) $[\text{Co}(\text{NH}_3)_3 \text{Cl}_3]$
(B) $[\text{Co}(\text{NH}_3)_4 \text{Cl}_2]$
(C) $[\text{Co}(\text{NH}_3)_6] \text{Cl}_3$
(D) $[\text{Co}(\text{NH}_3)_5 \text{Cl}] \text{Cl}$


NEET 20

Correct Option: A, B**Solution:**

Conductance of any complex depends on the following factor.

- (1) Number of ions produced by complex.
(2) If number of ions are same then we will check charge on complex unit.
- $\left. \begin{array}{l} (1) [\text{Co}^{+3}(\text{NH}_3)_3 \text{Cl}_3] \\ (2) [\text{Co}^{+2}(\text{NH}_3)_4 \text{Cl}_2] \end{array} \right\} \text{Both complex units have no charge. Therefore both complex units have same conductance.}$





64. Match List-I with List-II

List-I		List-II	
(A)	XeO_3	(I)	$\text{sp}^3 \text{d}$; linear
(B)	XeF_2	(II)	$\text{sp}^3 \text{d}^3$; pyramidal
(C)	XeOF_4	(III)	$\text{sp}^3 \text{d}^2$; distorted octahedral
(D)	XeF_6	(IV)	$\text{sp}^3 \text{d}^2$; square pyramidal




Choose the correct answer from the options given below :


- (A) A-II, B-I, C-IV, D-III (B) A-II, B-I, C-III, D-IV
 (C) A-IV, B-II, C-III, D-I (D) A-IV, B-II, C-I, D-III

NEET 2025





Correct Option: A

Solution:

Molecule	Hybridization	Shape
	sp^3	Pyramidal
	$\text{sp}^3 \text{d}$	Linear
	$\text{sp}^3 \text{d}^2$	Square pyramidal
	$\text{sp}^3 \text{d}^3$	Distorted Octahedral



Which of the following diagrams gives an accurate representation of the above reaction? [R $\rightarrow$ reactants; P $\rightarrow$ products]




NEET 20

Correct Option: A

Solution:

$\Delta H = -74.8 \text{ kJ mol}^{-1}$, it is an exothermic reaction. So, accurate representation is



66. Match List-I with List-II:

List-I (Example)		List-II (Type of Solution)	
(A)	Humidity	(I)	Solid in solid
(B)	Alloys	(II)	Liquid in gas
(C)	Amalgams	(III)	Solid in gas
(D)	Smoke	(IV)	Liquid in solid

Choose the correct answer from the options given below

- (A) A-II, B-IV, C-I, D-III (B) A-II, B-I, C-IV, D-I
 (C) A-III, B-I, C-IV, D-II (D) A-III, B-II, C-I, D-III

NEET 20

Correct Option: B

Solution:

- Humidity is a solution of liquid in gas

- Alloy is a solution of solid in solid
- Amalgam is a solution of liquid in solid
- Smoke is a solution of solid in gas


67. The correct order of decreasing basic strength of the given amines is:
- (A) N-methylaniline > benzenamine > ethanamine > N-ethylethanamine
- (B) N-ethylethanamine > ethanamine > benzenamine > N-methylaniline
- (C) N-ethylethanamine > ethanamine > N-methylaniline > benzenamine
- (D) benzenamine > ethanamine > N-methylaniline > N-ethylethanamine

NEET 2025

Correct Option: C**Solution:**

Basic strength order

N-ethylethanamine > Ethanamine > N-methylaniline > Benzenamine



68. Among the following choose the ones with equal number of atoms.
- A. 212 g of Na_2CO_3 (s) [molar mass = 106 g]
- B. 248 g of Na_2O (s) [molar mass = 62 g]
- C. 240 g of NaOH (s) [molar mass = 40 g]
- D. 12 g of H_2 (g) [molar mass = 2 g]
- E. 220 g of CO_2 (g) [molar mass = 44 g]
- Choose the correct answer from the options given below:
- (A) A, B and C only (B) A, B and D only
- (C) B, C and D only (D) B, D and E only

NEET 2025

Correct Option: B**Solution:**

$$\text{No. of atoms} = \text{Atomicity} \times \text{mole} \times N_A$$

- (A) No. of atoms = $6 \times \frac{212}{106} \times N_A = 12N_A$
- (B) No. of atoms = $3 \times \frac{248}{62} \times N_A = 12N_A$
- (C) No. of atoms = $3 \times \frac{240}{40} \times N_A = 18N_A$
- (D) No. of atoms = $2 \times \frac{12}{2} \times N_A = 12N_A$
- (E) No. of atoms = $3 \times \frac{220}{44} \times N_A = 15N_A$

69. Match List-I with List-II.

List-I (Name of Vitamin)	(Name of)	List-II (Deficiency disease)	(Deficiency)
(A)	Vitamin B12	(I)	Cheilosis
(B)	Vitamin D	(II)	Convulsions
(C)	Vitamin B2	(III)	Rickets
(D)	Vitamin B6	(IV)	Pernicious anaemia

Choose the correct answer from the options given below

- (A) A-IV, B-III, C-I, D-II (B) A-I, B-III, C-II, D-III
- (C) A-II, B-III, C-I, D-IV (D) A-IV, B-III, C-II, D-III

NEET 2019

Correct Option: A**Solution:**


List-I (Name of Vitamin)	List-II (Deficiency disease)
(A) Vitamin B12	Pernicious anaemia
(B) Vitamin D	Rickets
(C) Vitamin B2	Cheilosis
(D) Vitamin B6	Convulsions

70. The correct order of decreasing acidity of the following aliphatic acids is :-
- (A) $(\text{CH}_3)_3\text{CCOOH} > (\text{CH}_3)_2\text{CHCOOH} > \text{CH}_3\text{COOH} > \text{HCOOH}$
- (B) $\text{CH}_3\text{COOH} > (\text{CH}_3)_2\text{CHCOOH} > (\text{CH}_3)_3\text{CCOOH} > \text{HCOOH}$
- (C) $\text{HCOOH} > \text{CH}_3\text{COOH} > (\text{CH}_3)_2\text{CHCOOH} > (\text{CH}_3)_3\text{CCOOH}$
- (D) $\text{HCOOH} > (\text{CH}_3)_3\text{CCOOH} > (\text{CH}_3)_2\text{CHCOOH} > \text{CH}_3\text{COOH}$

NEET 2019

Correct Option: C**Solution:**Acidic strength $\propto -1$ effect

$$\propto \frac{1}{I_{\text{effect}}}$$



71. Given below are two statements :

Statement I : Ferromagnetism is considered as an extreme form of paramagnetism.**Statement II :** The number of unpaired electrons in a Cr^{2+} ion ($Z = 24$) is the same as that of a Nd^{3+} ion ($Z = 60$).



In the light of the above statements, choose the correct answer from the options given below:

- (A) Both Statements I and Statement II are true
 (B) Both Statement I and Statements II are false
 (C) Statement I is true but Statement II is false
 (D) Statement I is false but Statements II is true



NEET 2025**Correct Option: C****Solution:**

Ferromagnetism > Paramagnetism.

Statement-I is true.



4 unpaired electrons.



3 unpaired electrons.

Statement-II is false.

72. Match List I with List II

List-I (Mixture)	List-II (Method of separation)
------------------	--------------------------------

(A)	$\text{CHCl}_3 + \text{C}_6\text{H}_5\text{NH}_2$	(I)	Distillation under reduced pressure
(B)	Crude oil in petroleum industry	(II)	Steam distillation
(C)	Glycerol from spent-lye	(III)	Fractional distillation
(D)	Aniline - water	(IV)	Simple distillation

Choose the correct answer from the options given below:

- (A) A-IV, B-III, C-I, D-II (B) A-IV, B-III, C-II, D-III
 (C) A-III, B-IV, C-I, D-II (D) A-III, B-IV, C-II, D-III

NEET 20**Correct Option: A****Solution:**

List-I (Mixture)	List-II (Method of separation)
(A) $\text{CHCl}_3 + \text{C}_6\text{H}_5\text{NH}_2$	Simple distillation
(B) Crude oil in petroleum industry	Fractional distillation
(C) Glycerol from spent-lye	Distillation under reduced pressure
(D) Aniline - water	Steam Distillation

73. For the reaction $\text{A}(\text{g}) \rightleftharpoons 2 \text{B}(\text{g})$, the backward reaction rate constant is higher than the forward reaction rate constant by a factor of 2500, at 1000 K.[Given : $R = 0.0831 \text{ L atm mol}^{-1} \text{ K}^{-1}$] K_p for the reaction at 1000 K is

- (A) 83.1 (B) 2.077×10^5
 (C) 0.033 (D) 0.021

NEET 20**Correct Option: C****Solution:**

$$\begin{aligned}
 K_C &= \frac{K_f}{K_b} \\
 &= \frac{K_f}{2500 K_f} \\
 &= \frac{1}{2500} \\
 K_p &= K_C (RT)^{\Delta n_g} \\
 &= \frac{1}{2500} (0.0831 \times 1000)^1 \\
 K_p &= 0.033
 \end{aligned}$$

74. Given below are two statements :

Statement I : Benzenediazonium salt is prepared by the reaction of aniline with nitrous acid at 273 – 278 K. It decomposes easily in the dry state.

Statement II : Insertion of iodine into the benzene ring is difficult and hence iodobenzene is prepared through the reaction of benzenediazonium salt with KI.

In the light of the above statements, choose the most appropriate answer from the options given below :


- (A) Both Statement I and Statement II are correct
 (B) Both Statement I and Statement II are incorrect
 (C) Statement I is correct but Statement II is incorrect
 (D) Statement I is incorrect but Statement II is correct

NEET 2025


Correct Option: A

Solution:

Benzenediazonium chloride is a colourless crystalline solid. It is readily soluble in water and is stable in cold but reacts with water when warmed. It decomposes easily in the dry state.



75. How many products (including stereoisomers) are expected from monochlorination of the following compound?




- (A) 2 (B) 3 (C) 5 (D) 6

NEET 20

Correct Option: D


Solution:

For the monochlorination of



the total number of products including stereoisomers is

76. Among the given compound I-III, the correct bond dissociation energy of C – H bond marked with :-



- (A) II > I > III (B) I > II > III
 (C) III > II > I (D) II > III > I


NEET 20

Correct Option: A

Solution:

Bond dissociation energy $\propto$




$$\frac{1}{\text{Stability of free radical from after C-H bond fission homolytically}}$$




Stability order III > I > II





Bond dissociation energy order $\Rightarrow$ II > I > III.

77. Which one of the following compounds does not colorize bromine water?

- (A) 
- (B) 
- (C) 








NEET 2025


Correct Option: A**Solution:**

$\therefore$ Bromine water solution is decolourised in reaction (2), (3), (4) but not in (1)

78. The major product of the following reaction is



NEET 20

Correct Option: B**Solution:**

79. Which of the following aqueous solution will exhibit highest boiling point?

- (A) 0.01 M Urea (B) 0.01 M KNO₃
(C) 0.01 M Na₂SO₄ (D) 0.015 M C₆H₁₂O₆

NEET 20

Correct Option: C**Solution:**

$$\Delta T_b = i K_b \times m$$

$$\Delta T_b \propto i \times m$$

By considering molarity same as molality

- (1) 0.01 M Urea: $i \times m = 1 \times 0.01 = 0.01$
(2) 0.01 M KNO₃: $i \times m = 2 \times 0.01 = 0.02$
(3) 0.01 M Na₂SO₄: $i \times m = 3 \times 0.01 = 0.03$
(4) 0.015 M C₆H₁₂O₆: $i \times m = 1 \times 0.015 = 0.015$

$$T'_b = T_b^* + \Delta T_b$$

Higher the value of ($i \times m$) more will be the boiling point

80. Match List-I with List-II

List I		List II	
A.	Haber process	I.	Fe catalyst
B.	Wacker oxidation	II.	PdCl ₂

C.	Wilkinson catalyst	III.	$[(PPh_3)_3 RhCl]$
D.	Ziegler catalyst	IV.	$TiCl_4$ with $Al(CH_3)_3$

Choose the correct answer from the options given below :

(A) A – I, B – II, C – IV, D – III (B) A – II, B – III, C – I, D – IV

(C) A – I, B – II, C – III, D – IV (D) A – I, B – IV, C – III, D – II

NEET 2025

Correct Option: C

Solution:

81. 5 moles of liquid X and 10 moles of liquid Y make a solution having a vapour pressure of 70 torr. The vapour pressures of pure X and Y are 63 torr and 78 torr respectively. Which of the following is true regarding the described solution?

- (A) The solution shows positive deviation.
 (B) The solution shows negative deviation.
 (C) The solution is ideal.
 (D) The solution has volume greater than the sum of individual volumes.

NEET 2025

Correct Option: B

Solution:

$$P_{total} = X_x P_x^\circ + X_y P_y^\circ$$

$$= \frac{5}{15} \times 63 + \frac{10}{15} \times 78$$

$$= 21 + 52$$

$$= 73 \text{ torr}$$

Observed total pressure of solution is 70 torr.

It is less than calculated total pressure.

Hence, it shows negative deviation.

82. Sugar 'X'

- A. is found in honey
 B. is a keto sugar
 C. exists in α and β - anomeric forms.
 D. Is laevorotatory.

'X' is :


- (A) D-Glucose (B) D-Fructose
 (C) Maltose (D) Sucrose

NEET 20

Correct Option: B


Solution:

D-Fructose is found in honey and is a keto sugar.



D – (–) – Fructose

83. Identify the suitable reagent for the following conversion



- (A) (i) $LiAlH_4$, (ii) H^+ / H_2O
 (B) (i) $AlH(iBu)_2$ (ii) H_2O
 (C) (i) $NaBH_4$, (ii) H^+ / H_2O
 (D) $H_2 / Pd - BaSO_4$

NEET 20

Correct Option: B

Solution:

The suitable reagent for the conversion is

(i) $AlH(iBu)_2$, (ii) H_2O .

84. Given below are two statements: one is labelled as Assertion (A) and the other is labelled as Reason (R) :



undergoes S_N2 reaction faster than



Reason (R) : Iodine is a better leaving group because of its large size.

In the light of the above statements, choose the correct answer from the options given below:

- (A) Both A and R are true and R is the correct explanation of A.
 (B) Both A and R are true but R is not the correct explanation of A.

(C) A is true but R is false

(D) A is false but R is true

NEET 2025

Correct Option: A**Solution:**

Hence







is most reactive

Rate of $S_N2 \propto$ Leaving tendency of leaving group. Iodine is better L.G. than chlorine due to its bigger size.

85. The standard heat of formation, in kcal/mol of Ba^{2+} is :
 [Given : standard heat of formation of SO_4^{2-} ion (aq) = -216kcal/mol , standard heat of crystallisation of $\text{BaSO}_4(\text{s}) = -4.5\text{kcal/mol}$, standard heat of formation of $\text{BaSO}_4(\text{s}) = -349\text{kcal/mol}$]

(A) -128.5 (B) -133.0 (C) +133.0 (D) +220.5

NEET 2025

Correct Option: A**Solution:**

$$\Delta H_f = -216\text{kcal/mol} \quad \dots(1)$$

$$\Delta H_{\text{crystallisation}} = -4.5\text{kcal/mol} \quad \dots(2)$$

$$\Delta H_f(\text{BaSO}_4) = -349\text{kcal/mol} \quad \dots(3)$$

$$\dots(4)$$

From equation (1), (2) and

$$\text{So, } -349 - (-4.5) - (-216)$$

$$\text{So } -349 + 4.5 + 216$$

$$= -349 + 220.5$$

$$= -128.5\text{kcal/mol}$$






86. Total number of possible isomers (both structural as well as stereoisomers) of cyclic ethers of molecular formula $\text{C}_4\text{H}_8\text{O}$ is :

(A) 6 (B) 8 (C) 10 (D) 11

NEET 2025

Correct Option: C**Solution:**

For cyclic ethers O should be in ring * carbon here is chiral



Total number of isomers = 2 + 1 + 1 + 1 + 2 + 1 + 1 : 10

87. Identify the correct orders against the pro mentioned:

A. $\text{H}_2\text{O} > \text{NH}_3 > \text{CHCl}_3$ - dipole momentB. $\text{XeF}_4 > \text{XeO}_3 > \text{XeF}_2$ - number of lone pairs on tral atomC. $\text{O}-\text{H} > \text{C}-\text{H} > \text{N}-\text{O}$ - bond lengthD. $\text{N}_2 > \text{O}_2 > \text{H}_2$ - bond enthalpy

Choose the correct answer from the options given below

(A) A, D only (B) B, D only (C) A, C only (D) B, C

NEET 20

Correct Option: A**Solution:**

	$\mu(\text{D})$
A. H_2O	1.85
NH_3	1.47
CHCl_3	1.04

B. XeF_4 : 2 lone pairs of electron XeO_3 : 1 lone pair of electron XeF_2 : 3 lone pairs of electronC. Order of Bond length :- $\text{N}-\text{O} > \text{C}-\text{H} > \text{O}-\text{H}$ D. N_2 Bond order is 3 H_2 Bond order is 1 O_2 Bond order is 2

88. Higher yield of NO in $\text{N}_2(\text{g}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{NO}(\text{g})$ can be obtained at

[ΔH of the reaction = $+180.7 \text{ kJ mol}^{-1}$]

- A. Higher temperature
B. Lower temperature
C. Higher concentration of N_2
D. Higher concentration of O_2

Choose the correct answer from the options given below :

- (A) A, D only (B) B, C only
(C) B, C, D only (D) A, C, D only

NEET 2025

Correct Option: D

Solution:

Yield of the product generally depends on

- Temperature
- Concentration of reactant(s) and product(s)
- Pressure

As this is an endothermic reaction ($\Delta H = +180.7 \text{ kJ mol}^{-1}$), so, increase in temperature will shift equilibrium in forward direction to increase yield of NO.

Increase in concentration of reactants (N_2 and O_2) also shifts the equilibrium in forward direction and increase the yield of NO.

Hence, (A), (C) and (D) only will increase yield of NO.

89. If the rate constant of a reaction is 0.03 s^{-1} , how much time does it take for 7.2 mol L^{-1} concentration of the reactant to get reduced to 0.9 mol L^{-1} ?

(Given: $\log 2 = 0.301$)

- (A) 69.3 s (B) 23.1 s (C) 210 s (D) 21.0 s

NEET 20

Correct Option: A

Solution:

$$k = 0.03 \text{ s}^{-1}$$

$$\begin{aligned} t &= \frac{2.303}{k} \log \frac{a}{a-x} \\ &= \frac{2.303}{0.03} \log \frac{7.2}{0.9} \\ &= \frac{2.303}{0.03} \log 8 \\ &= \frac{2.303}{0.03} \times 3 \times \log 2 \\ &= \frac{2.303}{0.03} \times 3 \times 0.301 \\ &= 69.3 \text{ s} \end{aligned}$$

90. Which one of the following reactions does NOT belong to "Lassaigne's test"?


- (A) $\text{Na} + \text{C} + \text{N} \xrightarrow{\Delta} \text{NaCN}$
(B) $2\text{Na} + \text{S} \xrightarrow{\Delta} \text{Na}_2\text{S}$
(C) $\text{Na} + \text{X} \xrightarrow{\Delta} \text{NaX}$
(D) $2\text{CuO} + \text{C} \xrightarrow{\Delta} 2\text{Cu} + \text{CO}_2$

NEET 20

Correct Option: D

Solution:

Nitrogen, sulphur, halogens and phosphorus present in an organic compound are detected by "Lassaigne's test".



NEET 2025 Paper-(PCB) - ANSWER KEY

Que.	46	47	48	49	50	51	52	53	54	55
Ans.	D	B	B	A	A	C	C	B	D	A
Que.	56	57	58	59	60	61	62	63	64	65
Ans.	D	B	D	D	D	C	B	A, B	A	A
Que.	66	67	68	69	70	71	72	73	74	75
Ans.	B	C	B	A	C	C	A	C	A	D
Que.	76	77	78	79	80	81	82	83	84	85
Ans.	A	A	B	C	C	B	B	B	A	A
Que.	86	87	88	89	90					
Ans.	C	A	D	A	D					